



Adventures in Bayesian Structural Time Series

Part 3: Analyzing SST Data

Andrew Bates, Josh Gloyd, Tyler Tucker





⊠ SST Data



- ⊠ SST Data
- ⊠ Use **bsts** for



- ⊠ SST Data
- ⊠ Use **bsts** for
 - ⊠ Fit



- ⊠ SST Data
- ⊠ Use **bsts** for
 - ⊠ Fit
 - ⊠ local level



- ⊠ SST Data
- ⊠ Use **bsts** for
 - ⊠ Fit
 - ⊠ local level
 - ⊠ local linear trend model



- ❖ SST Data
- ❖ Use **bsts** for
 - ❖ Fit
 - ❖ local level
 - ❖ local linear trend model
 - ❖ local trend with seasonality



- ⊠ SST Data
- ⊠ Use **bsts** for
 - ⊠ Fit
 - ⊠ local level
 - ⊠ local linear trend model
 - ⊠ local trend with seasonality
 - ⊠ Posterior distribution



- ⊠ SST Data
- ⊠ Use **bsts** for
 - ⊠ Fit
 - ⊠ local level
 - ⊠ local linear trend model
 - ⊠ local trend with seasonality
 - ⊠ Posterior distribution
 - ⊠ Forecast



- ⊠ SST Data
- ⊠ Use **bsts** for
 - ⊠ Fit
 - ⊠ local level
 - ⊠ local linear trend model
 - ⊠ local trend with seasonality
 - ⊠ Posterior distribution
 - ⊠ Forecast
 - ⊠ Model Comparison



⊗ Sea Surface Temperature near Gibraltar



- ⊠ Sea Surface Temperature near Gibraltar
- ⊠ Aggregated every 12 days




- ⊠ Sea Surface Temperature near Gibraltar
- ⊠ Aggregated every 12 days
- ⊠ January 2004 to November 2017



- ⊠ Sea Surface Temperature near Gibraltar
- ⊠ Aggregated every 12 days
- ⊠ January 2004 to November 2017
- ⊠ Obtained from [Argovis](#)



SST of Gilbralter region





Setup

```
library(readr)
library(bsts)
# bsts also loads BoomSpikeSlab, Boom, MASS, zoo, xts

gilbralter <- read_csv("data/gilbraltersimple.csv")
gilt <- ts(gilbralter$tempMean, start=c(2004,1,13),
          end=c(2017, 11, 25), frequency=30)
plot(gilt, main='SST of Gilbralter region',
     xlab='Date',
     ylab='Temperature (C)')
```



Local Level Model


$$y_t = \mu_t + \varepsilon_t \quad \varepsilon_t \sim N(0, \sigma_\varepsilon^2)$$

$$\mu_{t+1} = \mu_t + \xi_t \quad \xi_t \sim N(0, \sigma_\xi^2)$$

```
ll_ss <- list()
ll_ss <- AddLocalLevel(state.specification = ll_ss,
                       y = gilt)
ll_fit <- bsts(gilt, state.specification = ll_ss,
               niter = 1e3)
```





```
plot(ll_fit)
```






```
plot(ll_fit, 'components')
```





```
plot(ll_fit, 'residuals')
```





Forecasting

```
ll_pred <- predict(ll_fit, horizon = 30)  
plot(ll_pred, plot.original = 90)
```

