





Quantum Computing Electronic Structure

Professor Stephen Schnetzer
Eesh Gupta
July 20, 2020



Why Electronic Structure?



- 400 °C
- 200 atm

~2% of World's
total energy
supply




Nitrogenase

- 25 °C
- 1 atm

Why Electronic Structure?

- **Catalytic mechanism** i.e.
“how it accelerates the reaction” is not well understood.
- Crux: Composed of transition metal compounds that are **strongly correlated**.
- Classically: Intractable
- Quantum Computing: 150-200 *logical* qubits


FeMo-co:
Iron, Molybdenum,
Carbon, Hydrogen ,
Oxygen



Why is it so hard?

- Exponential Problem
- Correlated motion
- Quantum properties like interference

**Each box or hole represents a spin orbital - half of a molecular orbital.*




What are Chemists doing? Hartree Fock

- Treat effects of other electrons "**on average**"
- Uses Variational procedure to assign **n** electrons to **n** spin orbitals.
- Recovers **99%** of total energy



What are Chemists Doing? Failures of Hartree Fock

Limited Correlation due to **pigeonholing** of electrons into spin orbitals.



What are Chemists doing? Configuration Interaction

- Recognizes contribution of excited states to **correlation energy**
- If given a complete basis set, provides **exact solution**



$$\Psi_{\text{CI}} = \Phi + \sum_S t_s \Phi_s + \sum_D t_d \Phi_d + \dots = \sum_{\infty} t_i \Phi_i$$

↓
 $T_1 \Phi$

↓
 $T_2 \Phi$

Φ = Hartree Fock State

How many configurations do you see?



*** Remember, spin flips here are not allowed. So a spin up electron cannot excite to a spin down orbital!

What are Chemists Doing? Truncated CI and CC

Configuration
Interaction
Singles and
Doubles



$$\Psi_{\text{CISD}} = (1 + T_1 + T_2)\Phi = \Phi + T_1\Phi + T_2\Phi$$

$$e^x = 1 + x + x^2 + x^3 + \dots$$

Coupled
Cluster **S**ingles
and **D**oubles



$$\Psi_{\text{CCSD}} = e^{T_1+T_2}\Phi = \Phi + T_1\Phi + (T_1^2 + T_2)\Phi + (T_1^3 + T_1T_2)\Phi + \dots$$



Single
Excitations



Double
Excitations




Triple
Excitations

* Ignoring coefficients here

What are Chemists doing? Coupled Cluster

- Approximating higher excitations using smaller excitations
- This is done **without increasing** number of parameters.
- But this method breaks down for larger molecules.



$$T_1^3 + T_1 T_2 \approx T_3$$

Diagram illustrating the Coupled Cluster method equation:

$T_1^3 + T_1 T_2 \approx T_3$

Annotations:


- A blue arrow points from T_1^3 to t_s, t_d (with a green smiley face).
- A blue arrow points from T_3 to t_t (with a green frowny face).

What we desire?


- Most classical methods either require **too many resources** or account for **too little correlation**.
- Goal: a **feasible** way of getting the to **exact** energy.
- Quantum Computers can help!

What is Quantum Computing?


Bit




Qubit



Intel's 49 qubit
superconducting
"quantum chip"



Fuchs, Franz Georg, and Franz Georg FuchsEpost. "Diving Deep into Quantum Computing." *#SINTEFblog*, 1 Apr. 2019, blog.sintef.com/digital-en/diving-deep-into-quantum-computing/.



Dilution refrigerator that houses IBM's quantum computer

<https://phys.org/news/2017-05-ibm-powerful-universal-quantum-processors.html>

Why Quantum Computing?

Myth : Quantum computer is a **faster** version of classical computer

Fact: Only efficient for **certain types** of problems and gate speed is almost 100 times slower than that of classical logic gates

Myth: Quantum Computing is **powerful** because it tries all the possible solutions in parallel.

Fact: Amplitudes leading to wrong answer “interfere destructively” with each other and the amplitudes leading to the right answer “interfere constructively”

Quantum Encoding Jordan Wigner

- How do we **upload** electrons and spin orbitals onto quantum computers ?
- One possibility is to represent **every spin orbital** by a **qubit**.
- This is the Jordan Wigner encoding, one of the many quantum encodings in the literature.



Quantum Encoding Jordan Wigner




Suppose I add in an electron into a spin orbital. How would you represent that situation on a quantum computer?

Variational Quantum Eigensolver

Variational Quantum Eigensolver Simulation

- Using simulators (ideal quantum computers) to calculate energy of H₂ molecule as we pull the 2 hydrogen atom apart.
- In this simulation, we are working with
 - 4 qubits
 - 4 spin orbitals (1s, 1s*)
 - 2 electrons



Variation Quantum Eigensolver

Find the eigenvalue E

$$H\psi = E\psi$$




Find Parameters $\vec{t} = \{t_i\}$ such that
we minimize

$$\langle \psi(\vec{t}) | H | \psi(\vec{t}) \rangle$$

1. **Prepare** a state with a given a set of parameters
2. **Act** on the state with the Hamiltonian.
3. **Measure** the expectation value.
4. **Find** *better* parameters for the next iteration.

QC
CC



State Preparation

Reference State

- Reference state is generally chosen as the **hartree fock state** because of its high overlap with the ground state.



State Preparation UCCSD


- Since quantum gates are reversible and hence unitary operators, we need the **coupled cluster** operator be unitary.

$$\Psi = e^{T-T^\dagger} \Phi$$



Measurement Hamiltonian


What are some **interactions** between the particles that may contribute to the **total energy** of the system?



Measurement Break It Up

- We will **break** up Hamiltonian into smaller, local hamiltonians , **measure** expectation value of each and **combine** these values to get the energy.

$$\text{Hamiltonian} = K_n + P_{n,n} + K_e + P_{n,e} + P_{e,e}$$



n= nucleon
e= electron
K = Kinetic Energy
P = Potential Energy

Optimization


- a) “Robustness” against noise
- b) Number of Function Evaluations




Direct Search Algorithms




Gradient Descent Algorithms





Quantum Error Problem



Quantum Error

- **Environment** can change the state of qubits, resulting in errors to our computations.
- These errors are usually a combination of phase flip and bit flip.
- Example: Thermal Relaxation error

Bit Flip

$$|1\rangle \rightarrow |0\rangle$$

Phase Flip

$$|0\rangle + |1\rangle \rightarrow |0\rangle - |1\rangle$$

Quantum Error Correction

- Fixing quantum errors using additional *ancilla* qubits
- May need up to 1000 helper qubits to correct for 1 qubit
- Near term quantum computers (NISQ devices) will only contain ~ 50-100 qubits.

Example:

$$|0\rangle \Rightarrow |000\rangle$$

$$|1\rangle \Rightarrow |111\rangle$$


$$|000\rangle \rightsquigarrow |100\rangle \downarrow |0\rangle$$

Quantum Error Mitigation

- For low depth (small) circuits, techniques like
 - Extrapolation
 - Probabilistic Error Cancellation
 - Quantum Subspace Expansion


require a multiplicative overhead in number of measurements to mitigate errors.

- Hence, error mitigation does not require as many resources as error correction



<https://phys.org/news/2017-11-ion-qubits-early-glimpse-quantum.html>


Extrapolation What?





Deliberately make noise worse in order to improve results.

Extrapolation Noise Amplification

- Every gate operation on the IBMQ devices is executed using **microwave pulses**.
- Stretching these pulses will “**slow down**” the gate operations, giving qubits more time to talk to the environment.
- Hence, stretching pulses amplifies noise.



Extrapolation In Action on H2



Future for Chemistry Simulations

- Collaboration between chemists, physicists and computer scientists to improve upon
 - **State Preparation** like UCCSD ansatz
 - **Error Mitigation** techniques to reduce noise burden
 - **Systems and observables** more resistant to noise than others. (Eg: Dipole moment and charge density)

Thank You !



“It’s like the first day we see a plane flying, and we want to go to the moon.” - Marco De Vivo, theoretical

chemist at Genoa, on quantum computing and drug discoveries

References

[arXiv:1001.3855v3](#) [quant-ph]

[arXiv:1808.10402v3](#) [quant-ph]

[arXiv:1701.02691v2](#) [quant-ph]

QuTech Academy. “Quantum error correction codes | QuTech Academy” Online video clip. Youtube. Posted Dec 8, 2018.
Accessed May 15, 2020

Jarrod McClean. “Quantum Computation for the Discovery of New Materials” Online video clip. Youtube. Posted Jul 5, 2017.
Accessed May 15, 2020

If any questions/concerns, contact eag190@scarletmail.rutgers.edu

More Detailed Discussion on Quantum Encoding

I omitted this section because it concerns notational circus which can be distracting for students.

Why Second Quantization?

First Quantization Mess

Name- Tagging Electrons:

Electron 1 occupies Spin Orbital i .

Electron 2 occupies Spin Orbital j .



Exchange Symmetry Trap

What if I swap the 2 electrons?



$$\frac{1}{\sqrt{2}} \det \begin{pmatrix} \chi_i(\mathbf{x}_1) & \chi_j(\mathbf{x}_1) \\ \chi_i(\mathbf{x}_2) & \chi_j(\mathbf{x}_2) \end{pmatrix} = \frac{1}{\sqrt{2}} (\chi_i(\mathbf{x}_1)\chi_j(\mathbf{x}_2) - \chi_j(\mathbf{x}_1)\chi_i(\mathbf{x}_2))$$

Second Quantization

Better Bookkeeping

Avoid Name Tagging

“Spin Orbital i and Spin Orbital j both
have 1 electron.”



$|11\rangle$

Exchange Symmetry Trap

What if I swap the 2 electrons?

Second Quantization

Fermionic Operators


- With elegant notation, come elegant operators.

Creation
Operator

$$a^\dagger |0\rangle = |1\rangle$$

Anihilation
Operator

$$a |1\rangle = |0\rangle$$



$$T_1 = \sum_{\substack{i \in \text{occ} \\ a \in \text{virt}}} t_a^i a_a^\dagger a_i$$

Second Quantization

Anticommutation Relations

Exchange symmetry is satisfied by the following anticommutation relations:

$$a_i a_j + a_j a_i = 0$$

$$a_i^\dagger a_j^\dagger + a_j^\dagger a_i^\dagger = 0$$

$$a_i a_j^\dagger + a_j^\dagger a_i = \delta_{ij}$$

Second Quantization

Quantum Encoding Methods

- Makes mapping from the fermionic **Fock Space** to **Hilbert space** of qubits **efficient**.
- Second Quantized methods include **Jordan Wigner**, **Parity basis** and **Bravyi Kitaev** encodings.
- Methods differ on number of qubit operations to realise fermionic operator

Jordan Wigner

$$\begin{bmatrix} o_0 \\ o_1 \\ o_2 \\ o_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

Parity Basis

$$\begin{bmatrix} o_0 \mod 2 \\ o_0 + o_1 \mod 2 \\ o_0 + o_1 + o_2 \mod 2 \\ o_0 + o_1 + o_2 + o_3 \mod 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix}$$

Quantum Encoding

Mapping Operators

- Under Jordan Wigner encoding, qubits represent spin orbitals.
- To add and remove electrons from them, we act on qubits with combinations of pauli gates.

$$a^\dagger = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} = \frac{\sigma^x - i\sigma^y}{2}$$

$$a = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \frac{\sigma^x + i\sigma^y}{2}$$

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$
$$\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$
$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Pauli
Matrices