Landmark & Updated ICU Trials

What each trial showed and what changed practice · updated to 2025 · page 1/3

1. Sepsis: Resuscitation Strategy

✔ **Rivers EGDT** 2001
Protocolised early goal-directed therapy (CVP, MAP, $ScvO_2$) ↓ mortality in a single centre.

= **ProCESS · ARISE · ProMISe** 2014–15
EGDT vs usual care: no mortality difference. EGDT no longer required.

= **ANDROMEDA-SHOCK** 2019
CRT-targeted vs lactate-targeted resuscitation: no significant 28-day mortality difference (trend favours CRT).

✔ **ANDROMEDA-SHOCK-2** 2025
Personalised CRT-targeted resuscitation improved a hierarchical composite (death, organ support, length of stay).

= **CLASSIC** 2022
Restrictive vs standard IV fluids in septic shock: no difference in 90-day mortality.

= **CLOVERS** 2023
Restrictive (early vasopressor) vs liberal fluids in sepsis hypotension: no difference in 90-day mortality.

Bottom line: Give an initial bolus, then reassess (CRT, lactate, fluid responsiveness). Protocolised EGDT and fixed large volumes are not needed.

2. Fluid Type

= **SAFE** 2004
4% albumin vs saline: no mortality difference; harm in traumatic brain injury (SAFE-TBI).

= **ALBIOS** 2014
Albumin to keep ≥ 30 g/L in severe sepsis: no 90-day mortality benefit.

✖ **6S · CHEST** 2012
Hydroxyethyl starch ↑ RRT (both) and ↑ mortality (6S). Avoid HES.

✔ **SMART** 2018
Balanced crystalloids vs saline ↓ major adverse kidney events (MAKE30).

= **BaSICS · PLUS** 2021–22
Balanced solution vs saline: no difference in 90-day mortality.

Bottom line: Balanced crystalloids are first choice; avoid starches; avoid albumin and hypotonic fluids in TBI.

3. Vasopressors, MAP Target & Adjuncts

± **SOAP II** 2010
Dopamine vs norepinephrine: no mortality difference, more arrhythmias; worse in cardiogenic shock subgroup.

= **VASST · VANISH** 2008 · 2016
Adding vasopressin (or early vasopressin): no mortality benefit; spares norepinephrine.

✔ **ATHOS-3** 2017
Angiotensin II raised MAP in vasodilatory shock on high-dose vasopressors.

± **SEPSISPAM** 2014
MAP 80–85 vs 65–70: no mortality difference; less RRT in chronic hypertension, more AF.

= **65 trial** 2020
Permissive MAP 60–65 in patients ≥ 65 y: no significant mortality difference, less vasopressor.

✖ **VITAMINS · LOVIT** 2020 · 2022
IV vitamin C: no benefit; LOVIT ↑ death or persistent organ dysfunction.

Bottom line: Norepinephrine first, MAP ≥ 65 (lower is acceptable in elderly), add vasopressin as second agent. No vitamin C.

4. Corticosteroids

± **ADRENAL** 2018
Hydrocortisone 200 mg/day in septic shock: no 90-day mortality difference; faster shock reversal, fewer transfusions.

✔ **APROCCHSS** 2018
Hydrocortisone + fludrocortisone ↓ 90-day mortality in septic shock.

✔ **CAPE COD** 2023
Hydrocortisone in severe community-acquired pneumonia ↓ 28-day mortality (excluded influenza and shock).

✔ **DEXA-ARDS** 2020
Dexamethasone in moderate–severe ARDS ↑ ventilator-free days and ↓ 60-day mortality.

✔ **RECOVERY (dexamethasone)** 2021
COVID-19 on oxygen or ventilation: ↓ 28-day mortality; possible harm without oxygen.

Bottom line: Steroids for septic shock needing ongoing vasopressors, severe CAP, ARDS and COVID-19 on oxygen.

5. ARDS & Ventilation

✔ **ARMA (ARDSNet)** 2000
VT 6 vs 12 mL/kg PBW, Pplat ≤ 30: ↓ mortality (31% vs 40%).

✔ **PROSEVA** 2013
Prone ≥ 16 h/day if P/F < 150: ↓ 28-day mortality (16% vs 33%).

± **ACURASYS · ROSE** 2010 · 2019
Early 48 h cisatracurium: benefit in ACURASYS, not confirmed in ROSE (lighter sedation in controls).

✖ **ART** 2017
Aggressive recruitment + PEEP titration ↑ mortality.

✖ **OSCILLATE · OSCAR** 2013
HFOV: no benefit (OSCAR) or ↑ mortality (OSCILLATE).

± **FACTT** 2006
Conservative fluids ↑ ventilator-free days; no mortality difference.

± **EOLIA** 2018
VV-ECMO in very severe ARDS: 60-day mortality 35% vs 46% (not significant; 28% crossover). CESAR (2009): ECMO-centre referral helped.

Bottom line: Low VT, plateau ≤ 30, prone early in severe ARDS, conservative fluids once stable, ECMO in experienced centres for refractory cases.

6. Airway & Intubation

✔ **PREVENT** 2019
Bag-mask ventilation between induction and laryngoscopy ↓ severe hypoxaemia.

✔ **PREOXI** 2024
Pre-oxygenation with NIV vs face mask ↓ hypoxaemia during intubation.

✔ **DEVICE** 2023
Video vs direct laryngoscopy ↑ first-attempt success (85% vs 71%).

= **BOUGIE** 2021
Bougie vs stylet: no improvement in first-attempt success.

= **PREPARE II** 2022
Fluid bolus before intubation did not ↓ cardiovascular collapse.

Bottom line: Pre-oxygenate with NIV when possible, ventilate during apnoea, and use video laryngoscopy first.

= no difference ✔ benefit ✖ harm ± mixed

Results summarised for board revision; check the original papers and current guidelines.

Landmark & Updated ICU Trials

What each trial showed and what changed practice · updated to 2025 · page 2/3

7. Oxygen Therapy & Non-Invasive Support

= **ICU-ROX · HOT-ICU** 2020 · 2021
Conservative vs usual oxygen (HOT-ICU: PaO_2 60 vs 90 mmHg): no 90-day mortality difference.

= **PILOT** 2022
SpO_2 88–92 vs 92–96 vs 96–100%: no difference in ventilator-free days.

= **UK-ROX** 2025
Conservative oxygen ($SpO_2 \approx 90\%$) in ventilated adults: no difference in 90-day mortality.

± **FLORALI** 2015
HFNC vs NIV/face mask in hypoxaemic failure: no difference in intubation overall; ↓ 90-day mortality with HFNC.

✓ **Helmet NIV (Patel)** 2016
Helmet vs face-mask NIV in ARDS ↓ intubation (single centre).

✓ **HIGH-WEAN** 2019
After extubation in high-risk patients: NIV alternating with HFNC ↓ reintubation vs HFNC alone.

Bottom line: Avoid hyperoxia and hypoxaemia: target $SpO_2 \approx 92$–96%. HFNC first line for hypoxaemic failure; prophylactic NIV after high-risk extubation.

8. Kidney & Renal Replacement

= **AKIKI · IDEAL-ICU** 2016 · 2018
Early vs delayed RRT: no mortality difference; many delayed patients never needed RRT.

✗ **STARRT-AKI** 2020
Accelerated RRT: no mortality benefit, ↑ RRT dependence at 90 days.

✗ **AKIKI-2** 2021
Waiting even longer (more-delayed) beyond the usual criteria was harmful.

= **ATN · RENAL** 2008 · 2009
Higher-intensity RRT: no benefit (aim effluent 20–25 mL/kg/h).

± **BICAR-ICU** 2018
Bicarbonate for severe metabolic acidosis: no overall benefit; ↓ mortality in AKI (AKIN 2–3) subgroup.

Bottom line: Start RRT for urgent indications, not by a timer. Dose 20–25 mL/kg/h. Bicarbonate for severe acidaemia with AKI.

9. Weaning, Sedation & Delirium

✓ **ABC trial** 2008
Paired spontaneous awakening + breathing trials ↓ 1-year mortality.

✓ **Subirà** 2019
30-min pressure-support SBT vs 2-h T-piece ↑ successful extubation.

± **SPICE III** 2019
Early dexmedetomidine vs usual care: no mortality difference; more bradycardia/hypotension (possible harm < 65 y).

= **MENDS2** 2021
Dexmedetomidine vs propofol in sepsis: no difference in delirium-free days or mortality.

= **MIND-USA · AID-ICU** 2018 · 2022
Haloperidol (± ziprasidone) for delirium: no effect on delirium duration or mortality.

= **NONSEDA** 2020
No sedation vs light sedation: no mortality difference.

Bottom line: Light sedation, daily SAT + SBT, short PSV SBT, no routine antipsychotics for delirium.

10. Neurocritical Care

± **DECRA · RESCUEicp** 2011 · 2016
Decompressive craniectomy in TBI: DECRA worse function; RESCUEicp ↓ mortality but more severe disability.

= **ESETT** 2019
Levetiracetam, fosphenytoin, valproate in established status: equal (~ 47% stop seizures).

✓ **INTERACT3** 2023
ICH care bundle (BP, glucose, temperature, anticoagulation reversal) ↑ functional outcome.

✓ **SELECT2 · ANGEL-ASPECT** 2023
Thrombectomy benefits even large ischaemic core strokes.

✗ **SAFE-TBI** 2007
Albumin in TBI ↑ mortality.

Bottom line: Second-line antiseizure drugs are interchangeable; bundle care in ICH; thrombectomy for large-vessel occlusion including large core.

11. Transfusion & Bleeding

✓ **TRICC · TRISS** 1999 · 2014
Hb trigger 7 vs 9–10 g/dL: restrictive is safe in ICU and septic shock.

± **MINT** 2023
Acute MI: restrictive (Hb < 8) vs liberal (Hb < 10): no significant difference; trend favours liberal.

✓ **TRAIN** 2024
Acute brain injury: liberal (Hb < 9) ↓ unfavourable neurological outcome.

= **HEMOTION · SAHaRA** 2024–25
TBI and SAH: liberal transfusion did not significantly improve neurological outcome.

= **PROPPR** 2015
Plasma:platelets:RBC 1:1:1 vs 1:1:2 in trauma: no mortality difference; better haemostasis.

✓ **CRASH-2 · CRASH-3** 2010 · 2019
Tranexamic acid within 3 h ↓ bleeding death in trauma and in mild–moderate TBI.

✗ **HALT-IT** 2020
TXA in GI bleeding: no benefit, ↑ venous thrombosis.

Bottom line: Hb 7 for most ICU patients; consider higher in acute MI. TXA early in trauma, not in GI bleeding.

12. Pulmonary Embolism & VTE

± **PEITHO** 2014
Tenecteplase in intermediate-high-risk PE ↓ haemodynamic decompensation, ↑ major bleeding and stroke.

= **PREVENT (IPC)** 2019
Adding pneumatic compression to drug prophylaxis: no ↓ in proximal DVT.

Bottom line: Thrombolysis for high-risk (shocked) PE; monitor intermediate-high risk and rescue if it deteriorates.

= no difference ✓ benefit ✗ harm ± mixed

Results summarised for board revision; check the original papers and current guidelines.

Landmark & Updated ICU Trials

What each trial showed and what changed practice · updated to 2025 · page 3/3

13. Glucose, Nutrition & GI Prophylaxis

✖ **NICE-SUGAR** 2009
Tight glucose 4.5–6 mmol/L ↑ mortality; target 8–10 mmol/L (144–180 mg/dL).

= **TGC-fast** 2023
Tight glucose control without early parenteral nutrition: no benefit.

✔ **EPaNIC** 2011
Late (day 8) vs early parenteral nutrition: faster recovery, fewer infections.

= **CALORIES · NUTRIREA-2** 2014 · 2018
Enteral vs parenteral route: no mortality difference; EN in shock had more GI complications.

✔ **NUTRIREA-3** 2023
Low calorie/protein in the first week: faster readiness for discharge, fewer GI complications.

± **EFFORT Protein** 2023
Higher protein: no benefit; possible harm in AKI and high organ failure.

± **SUP-ICU · REVISE** 2018 · 2024
PPI prophylaxis ↓ clinically important GI bleeding (REVISE); no mortality difference.

Bottom line: Glucose 8–10 mmol/L, cautious enteral feeding in week 1, delay PN, PPI prophylaxis for ventilated high-risk patients.

14. Cardiogenic Shock

= **IABP-SHOCK II** 2012
IABP in MI cardiogenic shock: no mortality benefit.

✔ **CULPRIT-SHOCK** 2017
Culprit-lesion-only PCI ↓ 30-day death or RRT vs multivessel PCI.

= **DOREMI** 2021
Milrinone vs dobutamine: no difference in outcomes.

= **ECLS-SHOCK** 2023
Routine VA-ECMO in MI shock: no mortality benefit, more bleeding and vascular complications.

✔ **DanGer Shock** 2024
Impella CP in STEMI shock ↓ 180-day mortality (46% vs 59%), more complications.

Bottom line: Revascularise the culprit only; no routine IABP or VA-ECMO; microaxial pump for selected STEMI shock.

15. Cardiac Arrest & Post-Resuscitation

= **TTM · TTM2** 2013 · 2021
33 °C vs 36 °C, then 33 °C vs normothermia (treat fever ≥ 37.8 °C): no difference in survival or neurology.

✔ **HYPERION** 2019
33 °C after non-shockable arrest ↑ good neurological outcome (modest).

= **BOX** 2022
MAP 63 vs 77 mmHg and restrictive vs liberal oxygen: no difference.

= **TAME** 2023
Mild hypercapnia after arrest: no neurological benefit.

= **COACT · TOMAHAWK** 2019 · 2021
Immediate angiography in arrest without ST elevation: no benefit.

± **PARAMEDIC2** 2018
Adrenaline ↑ 30-day survival but not favourable neurological survival.

Bottom line: Prevent fever (≤ 37.7 °C), avoid hypotension and hypoxaemia, angiography immediately only for STEMI.

16. Infection & Antibiotics

✖ **MERINO** 2018
Pip-tazo vs meropenem for ESBL bacteraemia ↑ 30-day mortality.

✔ **SAPS (procalcitonin)** 2016
PCT-guided stopping ↓ antibiotic duration and mortality.

✔ **ADAPT-Sepsis** 2025
Daily PCT-guided protocol safely ↓ antibiotic duration; CRP-guided did not.

± **BLING III** 2024
Continuous vs intermittent β-lactam infusion: 90-day mortality 24.9% vs 26.8% (not significant).

✔ **BALANCE** 2024
Bloodstream infection: 7 days non-inferior to 14 days of antibiotics.

= **SuDDICU** 2022
Selective decontamination of the digestive tract: no significant mortality difference.

Bottom line: Carbapenem for ESBL bacteraemia, shorter courses (7 days for most bacteraemia), PCT to guide stopping, consider prolonged β-lactam infusion.

= no difference ✔ benefit ✖ harm ± mixed

Results summarised for board revision; check the original papers and current guidelines.

1. Sepsis: Resuscitation Strategy

✔ **Rivers EGDT** 2001
Protocolised early goal-directed therapy (CVP, MAP, $ScvO_2$) ↓ mortality in a single centre.

= **ProCESS · ARISE · ProMISe** 2014–15
EGDT vs usual care: no mortality difference. EGDT no longer required.

= **ANDROMEDA-SHOCK** 2019
CRT-targeted vs lactate-targeted resuscitation: no significant 28-day mortality difference (trend favours CRT).

✔ **ANDROMEDA-SHOCK-2** 2025
Personalised CRT-targeted resuscitation improved a hierarchical composite (death, organ support, length of stay).

= **CLASSIC** 2022
Restrictive vs standard IV fluids in septic shock: no difference in 90-day mortality.

= **CLOVERS** 2023
Restrictive (early vasopressor) vs liberal fluids in sepsis hypotension: no difference in 90-day mortality.

Bottom line: Give an initial bolus, then reassess (CRT, lactate, fluid responsiveness). Protocolised EGDT and fixed large volumes are not needed.

2. Fluid Type

= **SAFE** 2004
4% albumin vs saline: no mortality difference; harm in traumatic brain injury (SAFE-TBI).

= **ALBIOS** 2014
Albumin to keep ≥ 30 g/L in severe sepsis: no 90-day mortality benefit.

✖ **6S · CHEST** 2012
Hydroxyethyl starch ↑ RRT (both) and ↑ mortality (6S). Avoid HES.

✔ **SMART** 2018
Balanced crystalloids vs saline ↓ major adverse kidney events (MAKE30).

= **BaSICS · PLUS** 2021–22
Balanced solution vs saline: no difference in 90-day mortality.

Bottom line: Balanced crystalloids are first choice; avoid starches; avoid albumin and hypotonic fluids in TBI.

3. Vasopressors, MAP Target & Adjuncts

± **SOAP II** 2010
Dopamine vs norepinephrine: no mortality difference, more arrhythmias; worse in cardiogenic shock subgroup.

= **VASST · VANISH** 2008 · 2016
Adding vasopressin (or early vasopressin): no mortality benefit; spares norepinephrine.

✓ **ATHOS-3** 2017
Angiotensin II raised MAP in vasodilatory shock on high-dose vasopressors.

± **SEPSISPAM** 2014
MAP 80–85 vs 65–70: no mortality difference; less RRT in chronic hypertension, more AF.

= **65 trial** 2020
Permissive MAP 60–65 in patients ≥ 65 y: no significant mortality difference, less vasopressor.

✖ **VITAMINS · LOVIT** 2020 · 2022
IV vitamin C: no benefit; LOVIT ↑ death or persistent organ dysfunction.

Bottom line: Norepinephrine first, MAP ≥ 65 (lower is acceptable in elderly), add vasopressin as second agent. No vitamin C.

4. Corticosteroids

± **ADRENAL** 2018
Hydrocortisone 200 mg/day in septic shock: no 90-day mortality difference; faster shock reversal, fewer transfusions.

✔ **APROCCHSS** 2018
Hydrocortisone + fludrocortisone ↓ 90-day mortality in septic shock.

✔ **CAPE COD** 2023
Hydrocortisone in severe community-acquired pneumonia ↓ 28-day mortality (excluded influenza and shock).

✔ **DEXA-ARDS** 2020
Dexamethasone in moderate–severe ARDS ↑ ventilator-free days and ↓ 60-day mortality.

✔ **RECOVERY (dexamethasone)** 2021
COVID-19 on oxygen or ventilation: ↓ 28-day mortality; possible harm without oxygen.

Bottom line: Steroids for septic shock needing ongoing vasopressors, severe CAP, ARDS and COVID-19 on oxygen.

5. ARDS & Ventilation

✔ **ARMA (ARDSNet)** 2000
VT 6 vs 12 mL/kg PBW, Pplat ≤ 30: ↓ mortality (31% vs 40%).

✔ **PROSEVA** 2013
Prone ≥ 16 h/day if P/F < 150: ↓ 28-day mortality (16% vs 33%).

± **ACURASYS · ROSE** 2010 · 2019
Early 48 h cisatracurium: benefit in ACURASYS, not confirmed in ROSE (lighter sedation in controls).

✖ **ART** 2017
Aggressive recruitment + PEEP titration ↑ mortality.

✖ **OSCILLATE · OSCAR** 2013
HFOV: no benefit (OSCAR) or ↑ mortality (OSCILLATE).

± **FACTT** 2006
Conservative fluids ↑ ventilator-free days; no mortality difference.

± **EOLIA** 2018
VV-ECMO in very severe ARDS: 60-day mortality 35% vs 46% (not significant; 28% crossover). CESAR (2009): ECMO-centre referral helped.

Bottom line: Low VT, plateau ≤ 30, prone early in severe ARDS, conservative fluids once stable, ECMO in experienced centres for refractory cases.

6. Airway & Intubation

✔ **PREVENT** 2019
Bag-mask ventilation between induction and laryngoscopy ↓ severe hypoxaemia.

✔ **PREOXI** 2024
Pre-oxygenation with NIV vs face mask ↓ hypoxaemia during intubation.

✔ **DEVICE** 2023
Video vs direct laryngoscopy ↑ first-attempt success (85% vs 71%).

= **BOUGIE** 2021
Bougie vs stylet: no improvement in first-attempt success.

= **PREPARE II** 2022
Fluid bolus before intubation did not ↓ cardiovascular collapse.

Bottom line: Pre-oxygenate with NIV when possible, ventilate during apnoea, and use video laryngoscopy first.

7. Oxygen Therapy & Non-Invasive Support

= **ICU-ROX · HOT-ICU** 2020 · 2021
Conservative vs usual oxygen (HOT-ICU: PaO_2 60 vs 90 mmHg): no 90-day mortality difference.

= **PILOT** 2022
SpO_2 88–92 vs 92–96 vs 96–100%: no difference in ventilator-free days.

= **UK-ROX** 2025
Conservative oxygen ($SpO_2 \approx 90\%$) in ventilated adults: no difference in 90-day mortality.

± **FLORALI** 2015
HFNC vs NIV/face mask in hypoxaemic failure: no difference in intubation overall; ↓ 90-day mortality with HFNC.

✓ **Helmet NIV (Patel)** 2016
Helmet vs face-mask NIV in ARDS ↓ intubation (single centre).

✓ **HIGH-WEAN** 2019
After extubation in high-risk patients: NIV alternating with HFNC ↓ reintubation vs HFNC alone.

Bottom line: Avoid hyperoxia and hypoxaemia: target $SpO_2 \approx 92$–96%. HFNC first line for hypoxaemic failure; prophylactic NIV after high-risk extubation.

8. Kidney & Renal Replacement

= **AKIKI · IDEAL-ICU** 2016 · 2018
Early vs delayed RRT: no mortality difference; many delayed patients never needed RRT.

✖ **STARRT-AKI** 2020
Accelerated RRT: no mortality benefit, ↑ RRT dependence at 90 days.

✖ **AKIKI-2** 2021
Waiting even longer (more-delayed) beyond the usual criteria was harmful.

= **ATN · RENAL** 2008 · 2009
Higher-intensity RRT: no benefit (aim effluent 20–25 mL/kg/h).

± **BICAR-ICU** 2018
Bicarbonate for severe metabolic acidosis: no overall benefit; ↓ mortality in AKI (AKIN 2–3) subgroup.

Bottom line: Start RRT for urgent indications, not by a timer. Dose 20–25 mL/kg/h. Bicarbonate for severe acidaemia with AKI.

9. Weaning, Sedation & Delirium

✔ **ABC trial** 2008
Paired spontaneous awakening + breathing trials ↓ 1-year mortality.

✔ **Subirà** 2019
30-min pressure-support SBT vs 2-h T-piece ↑ successful extubation.

± **SPICE III** 2019
Early dexmedetomidine vs usual care: no mortality difference; more bradycardia/hypotension (possible harm < 65 y).

= **MENDS2** 2021
Dexmedetomidine vs propofol in sepsis: no difference in delirium-free days or mortality.

= **MIND-USA · AID-ICU** 2018 · 2022
Haloperidol (± ziprasidone) for delirium: no effect on delirium duration or mortality.

= **NONSEDA** 2020
No sedation vs light sedation: no mortality difference.

Bottom line: Light sedation, daily SAT + SBT, short PSV SBT, no routine antipsychotics for delirium.

10. Neurocritical Care

± **DECRA · RESCUEicp** 2011 · 2016
Decompressive craniectomy in TBI: DECRA worse function; RESCUEicp ↓ mortality but more severe disability.

= **ESETT** 2019
Levetiracetam, fosphenytoin, valproate in established status: equal (~ 47% stop seizures).

✔ **INTERACT3** 2023
ICH care bundle (BP, glucose, temperature, anticoagulation reversal) ↑ functional outcome.

✔ **SELECT2 · ANGEL-ASPECT** 2023
Thrombectomy benefits even large ischaemic core strokes.

✖ **SAFE-TBI** 2007
Albumin in TBI ↑ mortality.

Bottom line: Second-line antiseizure drugs are interchangeable; bundle care in ICH; thrombectomy for large-vessel occlusion including large core.

11. Transfusion & Bleeding

✔ **TRICC · TRISS** 1999 · 2014
Hb trigger 7 vs 9–10 g/dL: restrictive is safe in ICU and septic shock.

± **MINT** 2023
Acute MI: restrictive (Hb < 8) vs liberal (Hb < 10): no significant difference; trend favours liberal.

✔ **TRAIN** 2024
Acute brain injury: liberal (Hb < 9) ↓ unfavourable neurological outcome.

= **HEMOTION · SAHaRA** 2024–25
TBI and SAH: liberal transfusion did not significantly improve neurological outcome.

= **PROPPR** 2015
Plasma:platelets:RBC 1:1:1 vs 1:1:2 in trauma: no mortality difference; better haemostasis.

✔ **CRASH-2 · CRASH-3** 2010 · 2019
Tranexamic acid within 3 h ↓ bleeding death in trauma and in mild–moderate TBI.

✖ **HALT-IT** 2020
TXA in GI bleeding: no benefit, ↑ venous thrombosis.

Bottom line: Hb 7 for most ICU patients; consider higher in acute MI. TXA early in trauma, not in GI bleeding.

12. Pulmonary Embolism & VTE

± **PEITHO** 2014

Tenecteplase in intermediate-high-risk PE ↓ haemodynamic decompensation, ↑ major bleeding and stroke.

= **PREVENT (IPC)** 2019

Adding pneumatic compression to drug prophylaxis: no ↓ in proximal DVT.

Bottom line: Thrombolysis for high-risk (shocked) PE; monitor intermediate-high risk and rescue if it deteriorates.

13. Glucose, Nutrition & GI Prophylaxis

✖ **NICE-SUGAR** 2009
Tight glucose 4.5–6 mmol/L ↑ mortality; target 8–10 mmol/L (144–180 mg/dL).

= **TGC-fast** 2023
Tight glucose control without early parenteral nutrition: no benefit.

✓ **EPaNIC** 2011
Late (day 8) vs early parenteral nutrition: faster recovery, fewer infections.

= **CALORIES · NUTRIREA-2** 2014 · 2018
Enteral vs parenteral route: no mortality difference; EN in shock had more GI complications.

✓ **NUTRIREA-3** 2023
Low calorie/protein in the first week: faster readiness for discharge, fewer GI complications.

± **EFFORT Protein** 2023
Higher protein: no benefit; possible harm in AKI and high organ failure.

± **SUP-ICU · REVISE** 2018 · 2024
PPI prophylaxis ↓ clinically important GI bleeding (REVISE); no mortality difference.

Bottom line: Glucose 8–10 mmol/L, cautious enteral feeding in week 1, delay PN, PPI prophylaxis for ventilated high-risk patients.

14. Cardiogenic Shock

= **IABP-SHOCK II** 2012
IABP in MI cardiogenic shock: no mortality benefit.

✓ **CULPRIT-SHOCK** 2017
Culprit-lesion-only PCI ↓ 30-day death or RRT vs multivessel PCI.

= **DOREMI** 2021
Milrinone vs dobutamine: no difference in outcomes.

= **ECLS-SHOCK** 2023
Routine VA-ECMO in MI shock: no mortality benefit, more bleeding and vascular complications.

✓ **DanGer Shock** 2024
Impella CP in STEMI shock ↓ 180-day mortality (46% vs 59%), more complications.

Bottom line: Revascularise the culprit only; no routine IABP or VA-ECMO; microaxial pump for selected STEMI shock.

15. Cardiac Arrest & Post-Resuscitation

= **TTM · TTM2** 2013 · 2021
33 °C vs 36 °C, then 33 °C vs normothermia (treat fever ≥ 37.8 °C): no difference in survival or neurology.

✓ **HYPERION** 2019
33 °C after non-shockable arrest ↑ good neurological outcome (modest).

= **BOX** 2022
MAP 63 vs 77 mmHg and restrictive vs liberal oxygen: no difference.

= **TAME** 2023
Mild hypercapnia after arrest: no neurological benefit.

= **COACT · TOMAHAWK** 2019 · 2021
Immediate angiography in arrest without ST elevation: no benefit.

± **PARAMEDIC2** 2018
Adrenaline ↑ 30-day survival but not favourable neurological survival.

Bottom line: Prevent fever (≤ 37.7 °C), avoid hypotension and hypoxaemia, angiography immediately only for STEMI.

16. Infection & Antibiotics

✖ **MERINO** 2018
Pip-tazo vs meropenem for ESBL bacteraemia ↑ 30-day mortality.

✔ **SAPS (procalcitonin)** 2016
PCT-guided stopping ↓ antibiotic duration and mortality.

✔ **ADAPT-Sepsis** 2025
Daily PCT-guided protocol safely ↓ antibiotic duration; CRP-guided did not.

± **BLING III** 2024
Continuous vs intermittent β-lactam infusion: 90-day mortality 24.9% vs 26.8% (not significant).

✔ **BALANCE** 2024
Bloodstream infection: 7 days non-inferior to 14 days of antibiotics.

= **SuDDICU** 2022
Selective decontamination of the digestive tract: no significant mortality difference.

Bottom line: Carbapenem for ESBL bacteraemia, shorter courses (7 days for most bacteraemia), PCT to guide stopping, consider prolonged β-lactam infusion.