# Haemodynamic Monitoring – Section Summary

Arterial line • CVP • PA catheter • SVV • Shock profiles

Critical Care Board Review • Section 4 • MCQ-Oriented

## 1. Pressure Measurement

- Discrepant BP → compare **MAP** of cuff vs line; check transducer level and zero
- CVP used **with** clinical exam, not alone

## 2. PA Catheter

- CO underestimated (thermodilution) with **tricuspid regurgitation**
- Cardiogenic: **low CO, high PCWP, high SVR**
- Septic: low SVR (~700 dyn·s/cm$^5$)
- PAOP waveform: large v waves → MR; step-up → VSD

## 3. Fluid Responsiveness

- Low SVV → **stop fluids**, reassess
- SVV/PPV valid only in controlled ventilation, sinus rhythm

## 4. High-Yield MCQ Pearls

- ☑ TR → wrong thermodilution CO
- ☑ Cardiogenic: high PCWP + SVR

## 1. Pressure Measurement

- Discrepant BP → compare **MAP** of cuff vs line; check transducer level and zero
- CVP used **with** clinical exam, not alone

## 2. PA Catheter

- CO underestimated (thermodilution) with **tricuspid regurgitation**
- Cardiogenic: **low CO, high PCWP, high SVR**
- Septic: low SVR (~700 dyn·s/cm$^5$)
- PAOP waveform: large v waves → MR; step-up → VSD

## 3. Fluid Responsiveness

- Low SVV → **stop fluids**, reassess
- SVV/PPV valid only in controlled ventilation, sinus rhythm

## 4. High-Yield MCQ Pearls

- TR → wrong thermodilution CO
- Cardiogenic: high PCWP + SVR