# Analgesia, Sedation, Delirium & Coma – Section Summary

Sedation choice • Withdrawal • Scores • ABCDEF

Critical Care Board Review • Section 15 • MCQ-Oriented

## 1. Key Points

- Benzodiazepine problems → switch to **fentanyl, propofol, dexmedetomidine**
- Lorazepam/diazepam infusion: **propylene glycol** → hyperosmolar acidosis, haemolysis
- Analgesia-first: **fentanyl** infusion
- Refractory alcohol withdrawal → **propofol** (or phenobarbital)
- SBT pass: **RSBI < 105**
- FOUR score for coma (e.g. 4T); RASS +1 = restless
- ABCDEF: **A = Assess, prevent, manage pain**

## 2. High-Yield MCQ Pearls

- [x] Analgesia first
- [x] Propylene glycol toxicity with lorazepam

# 1. Key Points

- Benzodiazepine problems → switch to **fentanyl, propofol, dexmedetomidine**
- Lorazepam/diazepam infusion: **propylene glycol** → hyperosmolar acidosis, haemolysis
- Analgesia-first: **fentanyl** infusion
- Refractory alcohol withdrawal → **propofol** (or phenobarbital)
- SBT pass: **RSBI < 105**
- FOUR score for coma (e.g. 4T); RASS +1 = restless
- ABCDEF: **A = Assess, prevent, manage pain**

## 2. High-Yield MCQ Pearls

- Analgesia first
- Propylene glycol toxicity with lorazepam