Stroke – Section Summary

Imaging • Thrombolysis • ICH reversal • SAH

Critical Care Board Review • Section 16 • MCQ-Oriented

1. Ischaemic

- Suspected stroke → **non-contrast CT** first
- Beyond window → no tPA (unless imaging selection)
- Within window → **tPA + CTA** for thrombectomy
- Glucose check only lab required before tPA
- Deterioration → repeat CT stat

2. ICH & SAH

- Warfarin ICH → **PCC**
- Thunderclap headache → CT; negative → **LP** (or CTA)
- SAH → **secure aneurysm** (clip/coil)
- Modified Fisher → **vasospasm risk**

3. High-Yield MCQ Pearls

- [x] CT first
- [x] Warfarin ICH → PCC

1. Ischaemic

- Suspected stroke → **non-contrast CT** first
- Beyond window → no tPA (unless imaging selection)
- Within window → **tPA + CTA** for thrombectomy
- Glucose check only lab required before tPA
- Deterioration → repeat CT stat

2. ICH & SAH

- Warfarin ICH → **PCC**
- Thunderclap headache → CT; negative → **LP** (or CTA)
- SAH → **secure aneurysm** (clip/coil)
- Modified Fisher → **vasospasm risk**

3. High-Yield MCQ Pearls

- [x] CT first
- [x] Warfarin ICH → PCC