

MATH 247: Analysis 3

Instructor: Blake Madill

Spring 2021

Table of Contents

Part 1: Topology of Normed Vector Spaces

Week 1

Normed Vector Spaces

Convergence

Completeness

Banach Spaces

Week 2

Closed and Open Sets

Closure and Interior

Examples

More Properties

Week 3

Compactness I

Open Covers

Compactness II

Compactness III

Week 4

Limits

Continuity

Uniform Continuity

Extreme Value Theorem

Week 5

Sequences of Functions

Examples

Theorem A

Theorem B

Part 2: Multivariable Calculus

Week 6

Partial Derivatives

Differentiability

Total Derivatives

Continuous Partials

Appendix

Week 7

Differentiation Rules

The Chain Rule

The Mean Value Theorem

Tangent Hyperplanes

Appendix

Week 8

Higher Order Total Derivatives

Taylor's Theorem

Optimization

Examples

Appendix

Week 9

Inverse Function Theorem

Implicit Function Theorem

Lagrange Multipliers

Appendix

Week 10

Riemann Integration

Characterization Theorem

Jordan Content and Lebesgue Measure

Integrability and Measure

Appendix

Week 11

General Integrability

Fubini's Theorem

Iterated Integrals

Examples

Change of Variables

Week 1 Notes.

Math 247: ① Introduction to Real Analysis
② Multivariable Calculus

Module 1: Normed Vector Spaces.

Definition 1.1.1 (Normed vector space):

A normed vector space (NVS) is a vector space V over \mathbb{R} equipped with a function:

$$\|\cdot\|: V \rightarrow [0, \infty)$$

such that:

- ① $\|v\| = 0$ iff $v = 0$.
- ② For all $\alpha \in \mathbb{R}$, $v \in V$, $\|\alpha v\| = |\alpha| \|v\|$
- ③ (Triangle Inequality) For all $u, v \in V$,

$$\|u+v\| \leq \|u\| + \|v\|.$$

$\|\cdot\|$ is called a norm on V .

Remark 1.1.2: Geometric Motivations:

Connotation) A NVS is represented by the pair $(V, \|\cdot\|)$.

- ① $\|v\|$: "the length of v " or "the distance between v and 0 "
- ② $\|v-w\|$: "the distance between v and w "

Remark 1.1.3:

Real Analysis is the study of approximating objects related to real numbers (e.g.) \mathbb{R} , \mathbb{R}^n , sequences of functions from $\mathbb{R} \rightarrow \mathbb{R}$. Typically one approximates

bad, unruly elements (e.g. \mathbb{R}) with nicely ordered elements (e.g. \mathbb{Q}). The notion of distance in a NVS allows us to talk about approximation.

Example 1.1.4: $(\mathbb{R}, |\cdot|)$ (usual absolute value).

Example 1.1.5: $(\mathbb{R}, \|\cdot\|)$, where $\|\alpha\| = 3|\alpha|$.

Example 1.1.6: $V = \mathbb{R}^n$ (ℓ^p norm)

For $p \geq 1$ and $V = (V_1, \dots, V_n) \in \mathbb{R}^n$

$$\|V\|_p = \left(\sum_{i=1}^n |V_i|^p \right)^{1/p}$$

is a norm on \mathbb{R}^n .

When $p=2$:

$$\|V\|_2 = \left(\sum_{i=1}^n V_i^2 \right)^{1/2}$$

is called the Euclidean norm. This is the usual distance in \mathbb{R}^n .

Remark 1.1.7: We equip \mathbb{R}^n with $\|\cdot\|_2$, unless stated otherwise.

Example 1.1.8: (infinity norm). $V = \mathbb{R}^n$

$$\|V\|_\infty = \max \{ |V_1|, |V_2|, \dots, |V_n| \}.$$

Example 1.1.9: $V = \mathbb{R}^{\mathbb{N}} = \{ \text{can} \}_{n=1}^{\infty} : a_i \in \mathbb{R} \}$.

$V = (V_1, V_2, \dots) \in V$.

For $p \geq 1$,

$$\|V\|_p = \left(\sum_{i=1}^{\infty} |V_i|^p \right)^{1/p}$$

$$\|V\|_\infty = \sup \{ |V_1|, |V_2|, \dots \}.$$

Define:

① $\ell^p = \{v \in \mathbb{R}^{\mathbb{N}} : \|v\|_p < \infty\}$. (ℓ^p can be shown to be a subspace of V). $(\ell^p, \|\cdot\|_p)$ is a NVS.

② $\ell^\infty = \{v \in \mathbb{R}^{\mathbb{N}} : \|v\|_\infty < \infty\}$ (the set of all bounded sequences). $(\ell^\infty, \|\cdot\|_\infty)$ is a NVS.

Example 1.1.10:

$V = C([a, b]) = \{f: [a, b] \rightarrow \mathbb{R} : f \text{ is continuous}\}$.

① $p \geq 1, f \in V$.

$$\|f\|_p \equiv \left(\int_a^b |f(x)|^p dx \right)^{1/p}.$$

is a norm. This is called the integration norm.

② $\|f\|_\infty \equiv \sup \{ |f(x)| : x \in [a, b] \}$

$$\stackrel{\text{EVI}}{=} \max \{ |f(x)| : x \in [a, b] \}$$

is a norm. This is called the uniform norm.

Remark 1.1.11: Unless otherwise stated, we equip $C([a, b])$ with $\|\cdot\|_\infty$.

Module 2: Convergence

Remark 1.2.1: V is a NVS. By a sequence in V we mean $(a_n)_{n=1}^{\infty} = (a_1, a_2, \dots)$, where each $a_i \in V$. We denote this by $(a_n) \subseteq V$.

Definition 1.2.2: V is a NVS. We say (a_n) converges to $a \in V$, written $a_n \rightarrow a$, if for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that $\|a_n - a\| < \varepsilon$ for all $n \geq N$.

Definition 1.2.3: For $(a_n) \subseteq V$, if for all $a \in V$, $a_n \not\rightarrow a$, we say (a_n) diverges.

Example 1.2.4: $V = \ell^{\infty}$, $(a_n) \subseteq V$.

$$a_n = (1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, 0, 0, \dots)$$

Claim: $a_n \rightarrow a$, where

$$a = (1, \frac{1}{2}, \frac{1}{3}, \dots)$$

Proof:

Let $\varepsilon > 0$ be given. Choose $N \in \mathbb{N}$ such that

$\frac{1}{N} < \varepsilon$. For $n \geq N$, we have:

$$\begin{aligned} & \|a_n - a\|_{\infty} \\ &= \|(0, 0, \dots, \frac{1}{n+1}, \frac{1}{n+2}, \dots)\|_{\infty} \\ &= \sup \{0, |\frac{1}{n+1}|, |\frac{1}{n+2}|, \dots\} \\ &= \frac{1}{n+1} < \frac{1}{n} \leq \frac{1}{N} < \varepsilon. \end{aligned}$$

□

Example 1.2.5: $V = \mathbb{L}^\infty$, $(a_n) \subseteq V$ with

$$a_n = (1, \underbrace{1, \dots, 1}_{n \text{ times}}, 0, 0, \dots)$$

Claim: $a_n \rightarrow a$ with

$$a = (1, 1, 1, \dots)$$

Proof:

For any $n \in \mathbb{N}$,

$$\|a_n - a\|_\infty = 1.$$

□

Fact: (a_n) diverges

Definition 1.2.6: V is a NVS, $(a_n) \subseteq V$, $A \subseteq V$.

① We say A is bounded if there exists $M > 0$ such that $\|a\| \leq M$ for all $a \in A$.

② We say (a_n) is bounded if $\{a_1, a_2, \dots\}$ is bounded. In other words, $\exists M > 0$, $\|a_n\| \leq M$ for all n .

Theorem 1.2.7 (convergence implies bounded): if $(a_n) \subseteq V$ is convergent, then (a_n) is bounded.

Proof:

Suppose $a_n \rightarrow a \in V$. So, there exists $N \in \mathbb{N}$ such that if $n \geq N$, then

$$\|a_n - a\| < 1.$$

$$\begin{aligned} \text{For } n \geq N, \|a_n\| &= \| (a_n - a) + a \| \leq \|a_n - a\| + \|a\| \\ &< 1 + \|a\|. \end{aligned}$$

$$\text{Take } M = \max \{ \|a_1\|, \dots, \|a_{N-1}\|, 1 + \|a\| \}.$$

We have that $\|a_n\| \leq M$ for all $n \in \mathbb{N}$ \square

Note: the converse is not true.

Proposition 1.2.8: for $(a_n), (b_n) \subseteq V$, $a_n \rightarrow a$, $b_n \rightarrow b$,

① $a_n + b_n \rightarrow a + b$

② if $\alpha \in \mathbb{R}$ then $\alpha a_n \rightarrow \alpha a$.

Module 3. Completeness.

Definition 1.3.1: We say $(a_n) \subseteq V$ is a Cauchy sequence if for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that:

$$\|a_n - a_m\| < \varepsilon$$

for all $n, m \geq N$.

Proposition 1.3.2: (Convergence implies Cauchy). If $(a_n) \subseteq V$ is convergent then (a_n) is Cauchy.

Proof:

Let $\varepsilon > 0$ be given. There exists $N \in \mathbb{N}$ and $a \in V$ such that $\|a_n - a\| < \frac{\varepsilon}{2}$ for all $n \geq N$.

For $n, m \geq N$,

$$\begin{aligned} \|a_n - a_m\| &\leq \|a_n - a\| + \|a_m - a\| \\ &< \varepsilon. \end{aligned}$$

□

Example 1.3.3 (being Cauchy does not imply convergence).

$$C_00 = \{x_n \in \ell^\infty : \exists N, \forall n \geq N, x_n = 0\}$$

$(C_00, \|\cdot\|_\infty)$ is a NVS

Let $a_n = (1, \frac{1}{2}, \dots, \frac{1}{n}, 0, 0, \dots)$.

$$a = (1, \frac{1}{2}, \frac{1}{3}, \dots) \notin C_00.$$

Then, $a_n \rightarrow a$ in $\ell^\infty \Rightarrow (a_n) \subseteq \ell^\infty$ is Cauchy.

$\Rightarrow (a_n) \subseteq C_00$ is Cauchy.

Since $a_n \rightarrow a \notin C_00$ and limits are unique, we have $(a_n) \subseteq C_00$ diverges (see definition of divergence). So we are motivated to study the NVS that Cauchy and convergence do agree.

Definition 1.3.4. Let V be a NVS.

- ① We say $A \subseteq V$ is complete if whenever $(a_n) \subseteq A$ is Cauchy then $\exists a \in A$ such that $a_n \rightarrow a$.
- ② If V is complete, we call V a Banach space.

Example 1.3.5: $C(C_0, \| \cdot \|_\infty)$ is not a Banach space.

Module 4. Banach Spaces.

Example 1.4.1: $(\mathbb{C}^n, \|\cdot\|)$ is a Banach space.

Remark 1.4.2:

$$v = (v_1, v_2, \dots, v_n) \in \mathbb{C}^n, 1 \leq p < \infty.$$

$$\textcircled{1} \|v\|_p^p = |v_1|^p + \dots + |v_n|^p \leq n \|v\|_\infty^p$$

$$\textcircled{2} \|v\|_\infty^p \leq |v_1|^p + \dots + |v_n|^p = \|v\|_p^p$$

$$\textcircled{3} \|v\|_p \leq \sqrt[p]{n} \|v\|_\infty.$$

$$\|v\|_\infty \leq \|v\|_p.$$

\textcircled{4} So it suffices to show that $(\mathbb{C}^n, \|\cdot\|_\infty)$ is a Banach space to prove that $(\mathbb{C}^n, \|\cdot\|_p)$ are Banach spaces for all $1 \leq p \leq \infty$.

Proof:

Assume $(\mathbb{C}^n, \|\cdot\|_\infty)$ is a Banach space.

Let $1 \leq p < \infty$ and let $(a_k) \subseteq \mathbb{C}^n$ be Cauchy with respect to $\|\cdot\|_p$. So, (a_k) is Cauchy with respect to $\|\cdot\|_\infty$ (Remark 1.4.2 \textcircled{3}) $\Rightarrow a_k \rightarrow a \in \mathbb{C}^n$ with respect to $\|\cdot\|_\infty$. $\Rightarrow a_k \rightarrow a$ with respect to $\|\cdot\|_p$ (Remark 1.4.2 \textcircled{3}) $\Rightarrow (\mathbb{C}^n, \|\cdot\|_p)$ is a Banach Space.

Proposition 1.4.3: $(\mathbb{C}^n, \|\cdot\|_\infty)$ is a Banach Space.

Proof:

Suppose $(a_k) \subseteq \mathbb{C}^n$ is Cauchy. Say

$a_k = (a_k^{(1)}, a_k^{(2)}, \dots, a_k^{(n)})$.

for all $k \in \mathbb{N}$, where $a_i^{(j)} \in \mathbb{R}$.

Let $\varepsilon > 0$ be given. There exists $N \in \mathbb{N}$ such that $\|a_k - a_l\|_\infty < \varepsilon$ for all $k, l \geq N$. For $k, l \geq N$, and $1 \leq i \leq n$,

$$|a_k^{(i)} - a_l^{(i)}| \leq \|a_k - a_l\|_\infty < \varepsilon.$$

$\Rightarrow (a_k^{(i)})_{k=1}^\infty \subseteq \mathbb{R}$ is Cauchy for all $1 \leq i \leq n$.

Since \mathbb{R} is complete.

$$a_k^{(i)} \rightarrow b_i \in \mathbb{R}$$

for all $1 \leq i \leq n$.

We claim $a_k \rightarrow (b_1, b_2, \dots, b_n)$.

Let $\varepsilon > 0$ be given. Fix $1 \leq i \leq n$. There exists $N_i \in \mathbb{N}$ such that $|a_k^{(i)} - b_i| < \varepsilon$ for all $k \geq N_i$.

Let $N = \max \{N_1, \dots, N_n\}$. For $k \geq N$,

$$\begin{aligned} & \|a_k - (b_1, \dots, b_n)\|_\infty \\ &= \max \{ |a_k^{(i)} - b_i| : 1 \leq i \leq n \} < \varepsilon. \end{aligned} \quad \square$$

Proposition 1-4-4: ℓ^∞ is a Banach space.

Proof:

Let $(a_n) \subseteq \ell^\infty$ be Cauchy. For all $n \in \mathbb{N}$, we may write

$$a_n = (a_n^{(1)}, a_n^{(2)}, \dots)$$

with each $a_n^{(i)} \in \mathbb{R}$.

We claim For $i \in \mathbb{N}$, $(a_n^{(i)})$ is Cauchy.

Let $\varepsilon > 0$ be given.

There exists $N \in \mathbb{N}$ such that

$$\|a_n - a_m\|_\infty < \varepsilon$$

for $n, m \geq N$.

Fix $i \in \mathbb{N}$. For $n, m \geq N$,

$$\begin{aligned} |a_n^{(i)} - a_m^{(i)}| &\leq \sup \{ |a_n^{(i)} - a_m^{(i)}| : i \in \mathbb{N} \} \\ &= \|a_n - a_m\|_\infty < \varepsilon. \end{aligned}$$

By completeness of \mathbb{R} ,

$$a_n^{(i)} \rightarrow b_i \quad (n \rightarrow \infty)$$

for all $i \in \mathbb{N}$.

Let $b = (b_1, b_2, \dots)$. we claim that $a_n \rightarrow b$.

Let $\varepsilon > 0$ be given. There exists $N \in \mathbb{N}$ such that

$$\|a_n - a_m\|_\infty < \frac{\varepsilon}{2}.$$

for all $n, m \geq N$.

We have that.

$$|a_n^{(i)} - a_m^{(i)}| < \frac{\varepsilon}{2}.$$

for all $n, m \geq N$ and all $i \in \mathbb{N}$.

Take $m \rightarrow \infty$, for $n \geq N$.

$$|a_n^{(i)} - b_i| \leq \frac{\varepsilon}{2}.$$

for all $i \in \mathbb{N}$.

So, $\|a_n - b\|_\infty \leq \frac{\varepsilon}{2} < \varepsilon$ for all $n \geq N$. \square

Week 2 Notes.

Module 1. Closed and Open Sets.

Remark 2.1.1: Big idea.

For a NVS, V , some subsets $A \subseteq V$ work nicer with limits/convergence than others. By understanding these "nice" subsets helps us understand convergence in V .

Definition 2.1.2: Let V be NVS.

1. A subset $C \subseteq V$ is closed whenever $(c_n) \subseteq C$, such that $c_n \rightarrow a \in V$, then $a \in C$.

2. We say $U \subseteq V$ is open if $V \setminus U$ is closed.

3. The collection

$$\{U \subseteq V : U \text{ is open}\}$$

is called the Topology on V . The study of open/closed sets on a space is called Topology.

Example 2.1.3: $\emptyset, V \subseteq V$ are open and closed.

Example 2.1.4: $[0, 1] \subseteq \mathbb{R}$ is neither open nor closed.

Example 2.1.5: for $r > 0$, $a \in V$.

The closed ball of radius r , centered at a ,

$$\overline{B_r(a)} := \{x \in V : \|a - x\| \leq r\}$$

is closed.

Proof:

Let $(a_n) \subseteq \overline{B_r(a)}$ such that $a_n \rightarrow b \in V$.

By definition:

$$\|a_n - a\| \leq r$$

for all $n \in \mathbb{N}$. Since $a_n \rightarrow b$,

$$\|a_n - a\| \rightarrow \|b - a\|$$

$\leq r$ (limit preserves order).

$\Rightarrow b \in \overline{Br(a)}$. Hence, $\overline{Br(a)}$ is closed \square

Example 2.1.6 $\{x \in V : \|x - a\| \geq r\}$ is closed.

Example 2.1.7. The open ball.

$$Br(a) := \{x \in V : \|x - a\| < r\}.$$

is open.

Example 2.1.8: $V = \ell^\infty$.

$$C_0 := \{(x_n) \in \ell^\infty : x_n \rightarrow 0\}$$

is closed.

Proof:

Let $(a_n) \subseteq C_0$ such that $a_n \rightarrow a \in \ell^\infty$.

Let

$$a_n = (a_n^{(1)}, a_n^{(2)}, \dots),$$

for all $n \in \mathbb{N}$.

Hence, $\lim_{k \rightarrow \infty} a_n^{(k)} = 0$ for all $n \in \mathbb{N}$.

Say $a = (b_1, b_2, \dots)$.

Let $\varepsilon > 0$ be given. We may find $N_1, N_2 \in \mathbb{N}$.

$$\textcircled{1} \quad \|a_n - a\|_\infty < \frac{\varepsilon}{2} \text{ for } n \geq N_1.$$

②. $|a_{N_1}^{(k)}| < \frac{\varepsilon}{2}$ for $k \geq N_2$.

For $k \geq N_2$

$$\begin{aligned} |b_k| &= |a_{N_1}^{(k)} - b_k - a_{N_1}^k| \leq |a_{N_1}^{(k)} - b_k| + |a_{N_1}^k| \\ &\leq \|a_{N_1} - a\|_\infty + |a_{N_1}^{(k)}| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

This shows $b_k \rightarrow 0$ and so $a = (b_1, b_2, \dots) \in C_0$.



Hence, C_0 is closed. \square

Proposition 2.1.9. Let V be MVS, $U \subseteq V$.

The following are equivalent:

- ① U is open.
- ② For all $a \in U$, there exists $r > 0$ such that $B_r(a) \subseteq U$.

Example 2.1.10: In \mathbb{R}^2



Proof for proposition 2.1.9:

(\Rightarrow) Assume V is open. Hence, $V \setminus V$ is closed.

For contradiction, assume $\exists a \in U$ such that $\nexists r > 0$ with $B_r(a) \subseteq V$.

For all $n \in \mathbb{N}$, we may find $a_n \in B_{\frac{1}{n}}(a)$ such that $a_n \notin V$.

Note:

1. $\|a_n - a\| < \frac{1}{n} \rightarrow 0 \Rightarrow a_n \rightarrow a$.

2. $(a_n) \subseteq V \setminus V$. \leftarrow closed.

Therefore: $a \in V \setminus V$, contradicting $a \in U$.

(\Leftarrow) Assume for all $a \in U$, $\exists r > 0$ such that $B_r(a) \subseteq V$.

Claim: $V \setminus V$ is closed.

Let $(a_n) \subseteq V \setminus V$ such that $a_n \rightarrow a \in V$.

For contradiction, assume $a \in U$. we may find $r > 0$ such that $B_r(a) \subseteq V$. Since $a_n \rightarrow a$, we can find $N \in \mathbb{N}$ such that $\|a_N - a\| < r \Rightarrow$

$a_N \in B_r(a) \subseteq V \Rightarrow a_N \in U$, contradicting $a_N \in V \setminus V$.

Module 2: Closure and Interior.

Remark 2.2.1: Big Idea.

Given $A \subseteq V$, there are "naturally close" sets.

$U \subseteq A \subseteq C$ such that U is open and C is closed.

Proposition 2.2.2: Let V be MVS.

1. If $\{U_\alpha\}_{\alpha \in I}$ is a collection of open sets in V , then $U = \bigcup_{\alpha \in I} U_\alpha$ is open.

2. If $\{C_\alpha\}_{\alpha \in I}$ is a collection of closed sets in V . Then, $C = \bigcap_{\alpha \in I} C_\alpha$ is closed.

3. If $U_1, \dots, U_n \subseteq V$ are open, then

$$\bigcap_{i=1}^n U_i$$

is open.

4. If $C_1, \dots, C_n \subseteq V$ are closed, then

$$\bigcup_{i=1}^n C_i$$

is closed.

Proof Sketch.

1. Take $a \in \bigcup_{\alpha \in I} U_\alpha \Rightarrow \exists \alpha, a \in U_\alpha \Rightarrow \exists r > 0, B_r(a) \subseteq U_\alpha \Rightarrow \exists r > 0, B_r(a) \subseteq \bigcup_{\alpha \in I} U_\alpha$.

2. Consider $V \setminus \bigcap_{\alpha \in I} C_\alpha = \bigcup_{\alpha \in I} (V \setminus C_\alpha) \rightarrow$ so this is open.
 $\Rightarrow \bigcap_{\alpha \in I} C_\alpha$ is closed.

3. Take $a \in \bigcap_{i=1}^n U_i \Rightarrow \forall 1 \leq i \leq n, \exists r_i > 0$ such

That $\text{Br}(a) \subseteq U_i$. Take $r = \min\{r_1, \dots, r_n\}$
 $\Rightarrow \text{Br}(a) \subseteq \bigcap_{i=1}^n U_i$. So $\bigcap_{i=1}^n U_i$ is open.)

This is where finite
 intersection comes from.

4. $V \setminus \bigcup_{i=1}^n C_i = \bigcap_{i=1}^n (V \setminus C_i) \rightarrow$ so this is open.
 open.

So, $\bigcup_{i=1}^n C_i$ is closed.

Example 2.2.3. $\bigcap_{n=1}^{\infty} \left(\frac{1}{n}, \frac{1}{n} \right) = \{0\}$.
 open not open.

Example 2.2.4. $\bigcup_{n=1}^{\infty} \left[0, 1 - \frac{1}{n} \right] = [0, 1)$.
 closed not closed.

Definition 2.2.5: Let $A \subseteq V$.

1. The closure of A is.

$\bar{A} := \bigcap_{\substack{A \subseteq C \\ C \text{ closed}}} C$. (this is closed as well)

2. The interior of A is:

$\text{Int}(A) := \bigcup_{\substack{U \subseteq A \\ U \text{ open}}} U$ (this is open as well)

Remarks 2.2.6:

1. \bar{A} is the smallest closed set containing A .


2. $\text{Int}(A)$ is the largest open set contained in A .

Definition 2.2.7: Let V be MVS.

1. If $A \subseteq V$, then a is called a limit point of A if $\exists (a_n) \subseteq A$ with $a_n \rightarrow a$.

2. If $A \subseteq V$, then an interior point of A is $a \in A$ such that $\exists r > 0$, $B_r(a) \subseteq A$.

Example 2.2.8: In \mathbb{R}^2 .



Proposition 2.2.9: $\forall A \subseteq V$.

1. $\bar{A} = \{ \text{limit points of } A \}$.

2. $\text{Int}(A) = \{ \text{Interior points of } A \}$.

Proof.

② Piazza.

①. Let $X = \{ \text{limit points of } A \}$.

Claim: X is closed.

Let $(a_n) \subseteq X$ such that $a_n \rightarrow a \in V$.

For all $n \in \mathbb{N}$, $\exists b_n \in A$ such that $|a_n - b_n| < \frac{1}{n}$.

$\Rightarrow b_n = b_n - a_n + a_n \rightarrow 0 + a = a$.

$\Rightarrow a \in X$ since a is a limit point of A .

$\Rightarrow X$ is closed.

By definition, $\bar{A} \subseteq X$.

Let $x \in X$, so that $\exists (a_n) \subseteq A$ with $a_n \rightarrow x$.

Let $C \subseteq V$ be closed such that $A \subseteq C$.

Then $(A_n) \subseteq C$ and so $X \in C$.

Hence $X \subseteq C$ for all $A \subseteq C$ with C closed.

$$\Rightarrow X \subseteq \bar{A}.$$

$$\Rightarrow X = \bar{A}.$$

□

Remark 2.2.10. Summary

① A closed $\Leftrightarrow A$ contains all its limit points.


$$\Leftrightarrow A = \bar{A}.$$

② A open \Leftrightarrow All points in A are interior points

$$\Leftrightarrow A = \text{Int}(A).$$


Module 3: Examples

Example 2.3.1: In \mathbb{R}^2 :




A

$\text{Int}(A)$:



\bar{A} :



Example 2.3.2: $A = \{(a_n) \in \ell^1 : a_n \in \mathbb{Q}\}$.

Claim: $\bar{A} = \ell^1$.

Remark 2.3.3: Let $x \in \ell^1$.

Assume $\forall \varepsilon > 0$, $\exists a \in A$, $\|x - a\|_1 < \varepsilon$.

Then, $x \in \bar{A}$.

$\forall n \in \mathbb{N}$, $\exists a_n \in A$, $\|x - a_n\|_1 < \frac{1}{n}$.

Then, $(a_n) \subseteq A$ and $a_n \rightarrow x \Rightarrow x \in \bar{A}$.

Proof for Example 2.3.2:

Let $x = (x_1, x_2, \dots) \in \ell^1$ and let $\varepsilon > 0$ be given.

By the density of rationals, $\forall n \in \mathbb{N}$, $\exists y_n \in \mathbb{Q}$ such that $|x_n - y_n| < \frac{\varepsilon}{2n}$.

Consider $y = (y_1, y_2, \dots)$

Then,

$$\|x-y\|_1 = \sum_{n=1}^{\infty} |x_n - y_n| < \sum_{n=1}^{\infty} \frac{\epsilon}{2^n} = \epsilon.$$

Therefore, $x \in \overline{A}$, so $\mathcal{J}' \subseteq \overline{A}$, $\mathcal{J}' = \overline{A}$. \square

Example 2.3.4. $V = \ell^\infty$.

Claim: $\overline{C_0} = C_0$.

Proof:

We know that $C_0 \subseteq \overline{C_0}$ and C_0 is closed.

So, $\overline{C_0} \subseteq C_0$.

Let $x = (x_1, x_2, \dots) \in C_0$ and let $\epsilon > 0$ be given. Since $x_n \rightarrow 0$, $\exists N \in \mathbb{N}$ such that $|x_n| < \frac{\epsilon}{2}$ for all $n \geq N$.

Let $y = (x_1, \dots, x_{N-1}, 0, 0, \dots) \in C_0$.

$$\begin{aligned} \text{So, } \|x-y\|_\infty &= \|(0, \dots, 0, x_N, x_{N+1}, \dots)\|_\infty \\ &= \sup \{ |x_k| : k \geq N \} \leq \frac{\epsilon}{2} < \epsilon. \end{aligned}$$

So, $x \in \overline{C_0}$ and so $\overline{C_0} = C_0$.

(here we used Remark 2.3.3) \square

Module 4: More Properties.

Proposition 2.4.1: Let V be NVS.

$$1. \overline{A \cup B} = \overline{A} \cup \overline{B}$$

$$2. \text{Int}(A \cap B) = \text{Int}(A) \cap \text{Int}(B).$$

$$3. \overline{A \cap B} \subseteq \overline{A} \cap \overline{B}.$$

$$4. \text{Int}(A) \cup \text{Int}(B) \subseteq \text{Int}(A \cup B).$$

Proof:

①. Since $\overline{A} \cup \overline{B}$ is closed and $A \cup B \subseteq \overline{A} \cup \overline{B}$,
 $\overline{A \cup B} \subseteq \overline{\overline{A} \cup \overline{B}}$.

Since $A, B \subseteq A \cup B$, we have $\overline{A}, \overline{B} \subseteq \overline{A \cup B}$,

$$\text{so } \overline{A} \cup \overline{B} \subseteq \overline{A \cup B}.$$

□

Example 2.4.2: $A = (0, 1), B = (1, 2)$.

$$\overline{A \cap B} = \overline{\emptyset} = \emptyset.$$

$$\overline{A} \cap \overline{B} = [0, 1] \cap [1, 2] = \{1\}.$$

Example 2.4.3: $A = [0, 1], B = [1, 2]$.

$$\text{Int}(A \cup B) = (0, 2)$$

$$\text{Int}(A) \cup \text{Int}(B) = (0, 1) \cup (1, 2)$$

Proposition 2.4.4: Let $A \subseteq V$.

$$1. \text{Int}(V \setminus A) = V \setminus \overline{A}.$$

$$2. \overline{V \setminus A} = V \setminus \text{Int}(A).$$

Proof ①.

Since $V \setminus \overline{A} \subseteq V \setminus A$, and $V \setminus A$ is open, we have $V \setminus \overline{A} \subseteq \text{Int}(V \setminus A)$.

Observe that, $\underbrace{V \setminus \text{Int}(V \setminus A)}_{\text{closed}} \supseteq V \setminus (V \setminus A) = A$.

and so $\overline{A} \subseteq V \setminus \text{Int}(V \setminus A)$
 $\Rightarrow V \setminus \overline{A} \supseteq \text{Int}(V \setminus A)$.

Replace A with $V \setminus A$ to prove ② \square

Definition 2.4.5. Let $A \subseteq V$.

The boundary of A is:

$$\partial(A) := \overline{A} \setminus \text{Int}(A).$$

Proposition 2.4.6: Let $A \subseteq V$.

1. $\partial(A)$ is closed.
2. A is closed iff $\partial(A) \subseteq A$.

Proof Sketch:

$$1. \partial(A) = \overline{A} \setminus \text{Int}(A) = \underbrace{\overline{A}}_{\text{closed}} \cap \underbrace{(V \setminus \text{Int}(A))}_{\text{closed}}.$$

2. (\Rightarrow) A is closed.

$$\partial(A) \subseteq \overline{A} = A.$$

$$(\Leftarrow) . \partial(A) \subseteq A$$

$$\partial(A) = \overline{A} \setminus \text{Int}(A)$$

$$\Rightarrow \bar{A} = \partial(A) \cup \text{Int}(A)$$
$$\subseteq A \quad \subseteq A.$$

$$\Rightarrow A \subseteq \bar{A} \subseteq A \Rightarrow A = \bar{A} \Rightarrow A \text{ is closed.}$$

Week 3 Notes.

Module 1. Compactness 1.

Remark 3.1.1: Big Idea.

Compact sets $C \subseteq V$ have nice "finiteness" topological properties. They may be infinite, but in some sense are "small".

Definition 3.1.2: Let V be NVS.

We say $C \subseteq V$ is compact if every $(a_n) \subseteq C$ has a subsequence $a_{n_k} \rightarrow a \in C$.

Example 3.1.3: $A \subseteq \mathbb{R}^n$ closed and bounded.

Claim: A is compact.

Let $(a_k) \subseteq A$.

Since A is bounded, (a_k) is bounded.

By assignment, $\exists (a_{k_l})$ such that $a_{k_l} \rightarrow a \in \mathbb{R}^n$.

Since A is closed, $a \in A$. \square .

Example 3.1.4: $A = \{(1, 0, 0, \dots), (0, 1, 0, \dots), \dots\} \subseteq \ell^\infty$
 $(a_n) \subseteq A$ has no convergent subsequence.

For $n \neq m$, $\|a_n - a_m\|_\infty = 1$.

So, A is not compact.

Example 3.1.5: $\overline{B_1(0)} \subseteq \ell^\infty$ closed, bounded.

But this is not compact. (See the sequence above)

Proposition 3.1.6: $C \subseteq V$ is compact Then C is closed and bounded.

Proof:

Let $C \subseteq V$ be compact.

① Claim: C is closed.

Let $(a_n) \subseteq C$ such that $a_n \rightarrow a \in V$.

$\exists (a_{n_k})$ such that $a_{n_k} \rightarrow b \in C$.

However, we must have $a = b \in C$ (subsequences of a convergent sequence converge to the same limit).

② Claim: C is bounded.

Suppose C is not bounded.

For all $n \in \mathbb{N}$ we may find $a_n \in C$ such that $\|a_n\| \geq n$. Consider $(a_n) \subseteq C$. Every subsequence of (a_n) is unbounded \Rightarrow divergent, contradicting.

C is compact. \square

Theorem 3.1.7 [Heine-Borel Theorem].

A set $C \subseteq \mathbb{R}^n$ is compact iff it is closed and bounded. (already proved).

Proposition 3.1.8: Let $C \subseteq V$ be compact.

If $A \subseteq C$ is closed, then A is compact.

Proof sketch:

$(a_n) \subseteq A \subseteq C \Rightarrow \exists (a_{n_k})$ converges in C .

Since A is closed, the limit belongs to A .

Module 2. Open Covers.

Remark 3.2.1: Goal:

Give an alternate description of compactness which exhibits the finiteness motivation from Module 1.

Definition 3.2.2: Let V be NVS, $A \subseteq V$.

① An open cover of A is a collection of open sets $\{U_\alpha : \alpha \in I\}$ such that

$$A \subseteq \bigcup_{\alpha \in I} U_\alpha.$$

It is called finite if $|I| < \infty$.

② A subset of an open cover of A , $\{U_\alpha : \alpha \in I\}$, which is also an open cover of A , is called a subcover of $\{U_\alpha : \alpha \in I\}$.

Example 3.2.3: $V = \mathbb{R}$, $A = [0, 1]$

An open cover of A :

$$A \subseteq \bigcup_{\alpha \in [0, 1] \cap \mathbb{Q}} (\alpha - \frac{1}{4}, \alpha + \frac{1}{4}).$$

A finite subcover:

$$A \subseteq (-\frac{1}{4}, \frac{1}{4}) \cup (0, \frac{1}{2}) \cup (\frac{1}{4}, \frac{3}{4}) \cup (\frac{1}{2}, 1) \cup (\frac{3}{4}, \frac{5}{4}).$$

Example 3.2.4: $V = \mathbb{R}^2$, $A = \mathbb{Z} \times \mathbb{Z}$.

$$A \subseteq \bigcup_{\alpha \in \mathbb{Z} \times \mathbb{Z}} B_{\frac{1}{2}}(\alpha).$$

There is no finite subcover.

Example 3.2.6: $V = \mathbb{R}$, $A = [0, 1]$

$$A \subseteq \bigcup_{n \in \mathbb{N}} \left(\frac{1}{n}, 2\right).$$

There is no finite subcover.

Theorem 3.2.7: Let V be NVS, $A \subseteq V$.

Then, $A \subseteq V$ is compact iff every open cover of A has a finite subcover.

Will be proved in later modules.

Module 3. Compactness 2.

Lemma 3.3.1. Let V be NVS, $A \subseteq V$ be compact. Let $A \subseteq \bigcup_{\alpha \in I} U_\alpha$ be an open cover of A . There exists $R > 0$ such that for all $a \in A$, $B_R(a) \subseteq U_\alpha$ for some $\alpha \in I$.

Proof:

Suppose no such $R > 0$ exists. In particular, for all $n \in \mathbb{N}$, $\exists a_n \in A$ such that $B_{\frac{1}{n}}(a_n) \not\subseteq U_\alpha$ for all $\alpha \in I$.

Since $(a_n) \subseteq A$ and A is compact, $\exists a_{n_k} \rightarrow a \in A$.
Say $a \in U_{\alpha_0}$, $\alpha_0 \in I$.

Pick $M \in \mathbb{N}$ such that $B_{\frac{2}{M}}(a) \subseteq U_{\alpha_0}$.

Moreover, since $a_{n_k} \rightarrow a$, we may find $N \in \mathbb{N}$ such that $a_{n_k} \in B_{\frac{1}{M}}(a)$ for $k \geq N$.

Then, for $k \geq N$, such that $n_k \geq M$, take
 $x \in B_{\frac{1}{M}}(a_{n_k}) \Rightarrow \|x - a\| = \|x - a_{n_k} + a_{n_k} - a\|$
 $\leq \|x - a_{n_k}\| + \|a_{n_k} - a\|$
 $< \frac{1}{M} + \frac{1}{M} = \frac{2}{M}$
 $\Rightarrow x \in B_{\frac{2}{M}}(a)$.

Therefore, $B_{\frac{1}{M}}(a_{n_k}) \subseteq B_{\frac{2}{M}}(a) \subseteq U_{\alpha_0}$.

Since $n_k > M$, $B_{\frac{1}{n_k}}(a_{n_k}) \subseteq B_{\frac{1}{M}}(a_{n_k}) \subseteq U_{\alpha_0}$.

Contradiction. □

Remark 3.3.2: R is called the Lebesgue number.

Proposition 3.3.3 [Part 1] Let V be NVS.

If $A \subseteq V$ is compact, then every open cover of A has a finite subcover.

Proof:

Suppose $A \subseteq V$ is compact. Let $A \subseteq \bigcup_{\alpha \in I} U_\alpha$ be an open cover of A .

We may find $R > 0$ as in the lemma.

If $\exists a_1, \dots, a_n$ such that $A \subseteq B_R(a_1) \cup \dots \cup B_R(a_n)$ we are done.

So, suppose there is no such covering. Take:

$$a_1 \in A.$$

$$a_2 \in A, a_2 \notin B_R(a_1).$$

$$a_3 \in A, a_3 \in B_R(a_1) \cup B_R(a_2).$$

⋮

Since $(a_n) \subseteq A$ and A is compact, (a_n) has a convergent subsequence. However, for $n < m$,

$$a_m \notin B_R(a_n).$$

$$\Rightarrow \|a_m - a_n\| \geq R.$$

$\Rightarrow (a_n)$ has no Cauchy subsequence.

$\Rightarrow (a_n)$ has no convergent subsequence.

Contradiction. \square .

Module 4. Compactness 3.

Proposition 3.4.1: Let V be NVS. $A \subseteq V$. If every open cover of A has a finite subcover, then A is compact.

Lemma 3.4.2. Let V be NVS, $A \subseteq V$. Suppose every open cover of A has a finite subcover.

If $A \subseteq \bigcup_{\alpha \in I} U_\alpha$, where each U_α is relatively open in A , then $\exists \alpha_1, \alpha_2, \dots, \alpha_n \in I$ such that $A \subseteq U_{\alpha_1} \cup \dots \cup U_{\alpha_n}$.

Proof.

Suppose $A \subseteq \bigcup_{\alpha \in I} U_\alpha$, where $U_\alpha = A \cap O_\alpha$, ($O_\alpha \subseteq V$ is open).

$$\Rightarrow A \subseteq \bigcup_{\alpha \in I} (A \cap O_\alpha) = A \cap \bigcup_{\alpha \in I} O_\alpha \subseteq \bigcup_{\alpha \in I} O_\alpha.$$

$$\Rightarrow A \subseteq O_{\alpha_1} \cup \dots \cup O_{\alpha_n} \quad (\text{by assumption}).$$

$$\Rightarrow A \subseteq U_{\alpha_1} \cup \dots \cup U_{\alpha_n}.$$

□

Proof of Proposition 3.4.1:

Suppose $A \subseteq V$ such that every open cover of A has a finite subcover.

Consider $(A_n) \subseteq \overline{A}$.

For $k \in \mathbb{N}$, consider $C_k = \overline{\{A_n : n \geq k\}} \cap A$.

We want to show $\bigcap_{k=1}^{\infty} C_k \neq \emptyset$.

Each C_k is relatively closed in A . Hence, every $U_k = A \setminus C_k$ is relatively open in A .

For contradiction, assume $\bigcap_{k=1}^{\infty} C_k = \emptyset$.

$$A = A \setminus \emptyset = A \setminus (C_1 \cap C_2 \cap \dots) = \bigcup_{k=1}^{\infty} (A \setminus C_k) = \bigcup_{k=1}^{\infty} U_k.$$

By the lemma, there exists $\exists i_1 < i_2 < \dots < i_l$ such that $A \subseteq U_{i_1} \cup U_{i_2} \cup \dots \cup U_{i_l}$.

Since $C_1 \supseteq C_2 \supseteq \dots$, we have $U_1 \subseteq U_2 \subseteq \dots$.

So, $A \subseteq U_{i_l} \subseteq A$ (relatively open).

$$\Rightarrow A = U_{i_l}.$$

$$\Rightarrow C_{i_l} = A \setminus U_{i_l} = \emptyset,$$

However, $U_{i_l} \cap C_{i_l} = \emptyset$, contradiction.

Thus, we may find at $\bigcap_{k=1}^{\infty} C_k$.

[Piazza]

So, we may find $n_1 < n_2 < n_3 < \dots$ such that

$$\|a_{n_k} - a\| < \frac{1}{k} \text{ for every } k \in \mathbb{N}.$$

Hence, $(a_{n_k}) \subseteq A$ with $a_{n_k} \rightarrow a \in A$. \square

Week 4 Notes.

Module 1 Limits.

Remark 4.1.1: Goal:

Give a quick overview of limits of functions
 $f: A \rightarrow W$, $A \subseteq V$, where V and W are NVS.

Definition 4.1.2: $f: A \rightarrow W$, $A \subseteq V$, $a \in V$.

We say the limit of $f(x)$ as x approaches a is $w \in W$ if

① $a \in \overline{A \setminus \{a\}}$ and

② $\forall \varepsilon > 0$, $\exists \delta > 0$ such that if $x \in A$ with
 $0 < \|x - a\| < \delta$, then $\|f(x) - w\| < \varepsilon$.

We write

$$\lim_{x \rightarrow a} f(x) = w.$$

Note: w is unique.

Remark 4.1.3: The condition $a \in \overline{A \setminus \{a\}}$.

If $a \notin \overline{A \setminus \{a\}}$, then we call $a \in V$ an isolated point with respect to A .

If $a \notin \overline{A \setminus \{a\}}$, then $\exists r > 0$ such that $B_r(a) \cap A = \{a\}$ or \emptyset .

Proof sketch: suppose no such r exist, in particular for every $\frac{1}{n}$, $\exists a_n \in A \setminus \{a\}$ so that $a_n \in B_{\frac{1}{n}}(a)$. $a_n \rightarrow a$, contradicting $a \notin \overline{A \setminus \{a\}}$.

Thus, there is no $x \in A$ with $0 < \|x - a\| < r$.

Proposition 4.1.4: $A \subseteq V$, $f, g, h: A \rightarrow \mathbb{R}$, at $\overline{A \setminus \{a\}}$.

① [limits preserve order].

If $\lim_{x \rightarrow a} f(x)$, $\lim_{x \rightarrow a} g(x)$ exist and $f(x) \leq g(x)$ for all $x \in A$, then $\lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} g(x)$.

② [Squeeze Theorem].

If $f(x) \leq g(x) \leq h(x)$ and $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$ exist, then $\lim_{x \rightarrow a} g(x) = L$.

Remark 4.1.5:

We will use all limit laws freely. The proofs are nearly identical to $f: A \rightarrow \mathbb{R}$, $A \subseteq \mathbb{R}$ case.

Example 4.1.6:

$$\lim_{(x,y,z) \rightarrow (0,0,0)} \frac{xy^2 + x^2z + xyz}{\sqrt{x^2 + y^2 + z^2}}$$

Solution: If $x \neq 0$.

$$\begin{aligned} 0 &\leq \left| \frac{xy^2 + x^2z + xyz}{\sqrt{x^2 + y^2 + z^2}} \right| \leq \frac{|xy^2 + x^2z + xyz|}{\sqrt{x^2}} \\ &\leq \frac{|xy^2| + |x^2z| + |xyz|}{|x|} \leq \frac{|x|y^2 + |x|^2|z| + |x||y||z|}{|x|} \\ &= y^2 + |x||z| + |y||z|. \end{aligned}$$

Note: If $x=0$, then $f(x,y,z) = 0$.

$$\text{Since } \lim_{(x,y,z) \rightarrow (0,0,0)} y^2 + |x||z| + |y||z| = 0.$$

We know the original limit is 0 by the squeeze theorem.

Example 4.1.7

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2+y^4}.$$

As $(\frac{1}{n}, 0) \rightarrow (0, 0)$, $f(\frac{1}{n}, 0) = 0 \rightarrow 0$.

As $(\frac{1}{n^2}, \frac{1}{n}) \rightarrow (0, 0)$, $f(\frac{1}{n^2}, \frac{1}{n}) = \frac{1}{n^4} / (\frac{1}{n^4} + \frac{1}{n^4}) = \frac{1}{2} \rightarrow \frac{1}{2}$.

This limit does not exist.

Module 2 Continuity.

Remark 4.2.1: Notations.

Unless stated otherwise, we only consider

$f: A \rightarrow W$, $A \subseteq V$ where V, W are NVS.

Definition 4.2.2:

We say that f is continuous at $a \in A$ if for all $\epsilon > 0$, there exists $\delta > 0$ such that if $x \in A$ with $\|x - a\| < \delta$, then $\|f(x) - f(a)\| < \epsilon$.

Remark 4.2.3:

① If $a \in \overline{A \setminus \{a\}}$, then f is continuous at a iff $\lim_{x \rightarrow a} f(x) = f(a)$

② If $a \notin \overline{A \setminus \{a\}}$, then f is continuous at a .

Proof Sketch: $\exists r > 0$, $B_r(a) \cap A = \{a\}$. Let $\epsilon > 0$ be given. Choose $\delta = r$. If $x \in A$ and $\|x - a\| < \delta$, then $x = a$. So, $\|f(x) - f(a)\| = \|f(a) - f(a)\| = 0 < \epsilon$.

Definition 4.2.4:

If f is continuous at all $a \in A$, then we say f is continuous.

Proposition 4.2.5: The following are equivalent:

① f is continuous.

② f preserves convergence.

③ \forall open $U \subseteq W$ such that $f^{-1}(U)$ is relatively

open in A .

Proof:

② \Leftrightarrow ③ from assignment.

① \Leftrightarrow ②

Suppose f is continuous and let $(a_n) \subseteq A$ such that $a_n \rightarrow a \in A$. Let $\varepsilon > 0$ be given.

There exists $\delta > 0$ such that if $x \in A$ and $\|x - a\| < \delta$, then $|f(x) - f(a)| < \varepsilon$.

Take $N \in \mathbb{N}$ such that $\|a_n - a\| < \delta$ for all $n \geq N$.

Then for $n \geq N$,

$$|f(a_n) - f(a)| < \varepsilon.$$

which shows $f(a_n) \rightarrow f(a)$.

Assume f preserves convergence.

For contradiction, suppose f is discontinuous at a . There exists $\varepsilon > 0$ and $(a_n) \subseteq A$ such that $\|a_n - a\| < \delta$ but $|f(a_n) - f(a)| \geq \varepsilon$. Then $a_n \rightarrow a$ but $f(a_n) \not\rightarrow f(a)$. Contradiction. \square

Example 4.2.6: $P_i: \mathbb{R}^n \rightarrow \mathbb{R}$, $1 \leq i \leq n$.

$P_i(x_1, x_2, \dots, x_n) = x_i$ is continuous.

Proof:

Let $(a_k) \subseteq \mathbb{R}^n$ such that $a_k \rightarrow a \in \mathbb{R}^n$.

Say:

$$a_k = (a_k^{(1)}, a_k^{(2)}, \dots, a_k^{(n)})$$

$$a = (b_1, b_2, \dots, b_n).$$

We know (by assignment), $\alpha_k(x_i) \rightarrow b_i$.

$\Rightarrow p_i(\alpha_k) \rightarrow p_i(b_i)$.

So, p_i is continuous.

Proposition 4.2.7:

① If $f+g: A \rightarrow W$ are continuous, then $f+g$ and αf ($\alpha \in \mathbb{R}$) are continuous.

② If $f: A \rightarrow W$ is continuous and $g: B \rightarrow W_2$, $B \subseteq W_1$, is continuous, then $g \circ f$ is continuous.

Proof sketch:

① Let $(a_n) \subseteq A$ such that $a_n \rightarrow a$. Since f, g are continuous,

$$f(a_n) \rightarrow f(a).$$
$$g(a_n) \rightarrow g(a).$$

By the limit laws:

$$f(a_n) + g(a_n) \rightarrow f(a) + g(a).$$
$$\alpha f(a_n) \rightarrow \alpha f(a).$$

So, $f+g, \alpha f$ are continuous.

② Let $(a_n) \subseteq A$ such that $a_n \rightarrow a$.

Since f is continuous, $f(a_n) \rightarrow f(a)$.

Since g is continuous,

$$g(f(a_n)) \rightarrow g(f(a)).$$

So, $g \circ f$ is continuous.

Module 3: Uniform Continuity.

Definition 4.3.1: $A \subseteq V, f: A \rightarrow W$.

We say f is uniformly continuous if $\forall \varepsilon > 0, \exists \delta > 0$ such that if $x, a \in A$ with $\|x-a\| < \delta$, then $\|f(x) - f(a)\| < \varepsilon$.

Remark 4.3.2: Big idea.

The same δ works uniformly for all $a \in A$.

Note: uniformly continuous \rightarrow continuous.

Example 4.3.3.

$f: A \rightarrow W$ is Lipschitz if $\exists M > 0$ such that

$$\|f(a) - f(b)\| \leq M \|a - b\|$$

for all $a, b \in A$.

Claim: Lipschitz \Rightarrow uniformly continuous.

Proof:

Let $\varepsilon > 0$ be given. Choose $\delta = \frac{\varepsilon}{M}$

If $a, b \in A$ with $\|a - b\| < \delta$, then $\|f(a) - f(b)\| \leq M \|a - b\| < M \delta = \varepsilon$.

□

Example 4.3.4: $f: [0, \infty) \rightarrow \mathbb{R}$.

$$f(x) = x^2$$

Claim: f is not uniformly continuous

Proof:

Let $a_n = n + \frac{1}{n}$, $b_n = n$, $n \in \mathbb{N}$.


$$|a_n - b_n| = \frac{1}{n} \rightarrow 0.$$

$$|f(a_n) - f(b_n)| = \left(n + \frac{1}{n}\right)^2 - n^2 = 2 + \frac{1}{n^2} \geq 2.$$

□.

Example 4.3.5: $f: (0, 1] \rightarrow \mathbb{R}$.

$$f(x) = \ln(x)$$



Proof:

$$\text{Let } a_n = \frac{1}{n}, \quad b_n = \frac{1}{n^2}, \quad n \in \mathbb{N}.$$

$$|a_n - b_n| = \frac{1}{n} - \frac{1}{n^2} = \frac{n-1}{n^2} \rightarrow 0.$$

$$|f(a_n) - f(b_n)| = \ln \frac{1}{n} - \ln \frac{1}{n^2} = \ln n \rightarrow \infty. \quad \square.$$

Theorem 4.3.6:

If $C \subseteq V$ is compact and $f: C \rightarrow W$ is continuous, then f is uniformly continuous.

Proof Sketch:

Suppose f is not uniformly continuous.

$$\exists \varepsilon > 0, (a_n), (b_n) \subseteq C, \|a_n - b_n\| < \frac{1}{n},$$

$$\|f(a_n) - f(b_n)\| \geq \varepsilon.$$

By compactness, $\exists a_{n_k} \rightarrow a \in C$.

$$\Rightarrow \underbrace{b_{n_k} - a_{n_k}}_{\rightarrow 0} + \underbrace{a_{n_k}}_{\rightarrow a} \rightarrow a.$$

By continuity: $f(c_{nk}) \rightarrow f(a)$, $f(b_{nk}) \rightarrow f(a)$.
 $\Rightarrow |f(c_{nk}) - f(b_{nk})| \rightarrow 0$. Contradiction.

Module 4: Extreme Value Theorem.

Proposition 4.4.1:

If $C \subseteq V$ is compact, $f: C \rightarrow W$ is continuous, then $f(C)$ is compact.

Proof sketch:

Take $(f(a_n)) \subseteq f(C)$, $a_n \in C$.

$(a_n) \subseteq C$. $\exists a_n \rightarrow a \in C$.

$\Rightarrow f(a_n) \rightarrow f(a) \in f(C)$.

Lemma 4.4.2.

If $A \subseteq \mathbb{R}$ is bounded and non-empty, then

$\inf A, \sup A \in \bar{A}$.

Proof sketch:

$\forall n \in \mathbb{N}, \exists \sup A - \frac{1}{n} < a_n \leq \sup A$.

By squeeze theorem, $a_n \rightarrow \sup A$, so $\sup A \in \bar{A}$.

Theorem 4.4.3 [Extreme Value Theorem].

If $C \subseteq V$, $C \neq \emptyset$ and C is compact, $f: C \rightarrow \mathbb{R}$ is continuous, then there exists $a, b \in C$ such that

$$f(a) = \min f(C).$$

$$f(b) = \max f(C).$$

Proof:

$f(C)$ is compact by the proposition.

$\Rightarrow f(C)$ is closed and bounded.

$\sup f(C), \inf f(C) \in \overline{f(C)} = f(C)$.

$$f(a) = \inf f(C) = \min f(C).$$

$$f(b) = \sup f(x) = \max f(x).$$

□.

Proposition 4.4.4:

Let $K \subseteq V$ be compact, W be NVS.

$C(K, W) = \{f: K \rightarrow W \text{ continuous}\}$ is a NVS.

When equipped with the uniform norm:

$$\|f\|_{\infty} = \max \{ \|f(x)\| : x \in K \}.$$

Note: $f: K \rightarrow W$ is continuous

$\|\cdot\|: W \rightarrow \mathbb{R}$ is continuous.

$\Rightarrow \|f\|: K \rightarrow \mathbb{R}$ is continuous.

By the Extreme Value Theorem, $\max \{ \|f(x)\| : x \in K \}$ exists.

Week 5 Notes

Module 1. Sequences of Functions.

Remark 5.1.1: Big Idea.

Given $A \subseteq V$ and W , and a sequence of functions $f_n: A \rightarrow W$, how can we make sense of (f_n) "converging" to some function $f: A \rightarrow W$?

Definition 5.1.2: $A \subseteq V$, $f_n: A \rightarrow W$, $f: A \rightarrow W$.

① We say f_n converges to f pointwise if $f_n(x) \rightarrow f(x)$ for all $x \in A$.

② We say f_n converges to f uniformly if for all $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that $\|f_n(x) - f(x)\| < \epsilon$ for all $n \geq N$ and $x \in A$.

Remark 5.1.3: Idea.

If $f_n \rightarrow f$ uniformly, then the same N works uniformly for all $x \in A$.

Remark 5.1.4: Notation.

Let $f_n, f: A \rightarrow W$, $A \subseteq V$.

$$\|f_n - f\|_\infty := \sup \{ \|f_n(x) - f(x)\| : x \in A \}.$$

This is not necessarily a norm, because it could be ∞ .

Note that:

$$f_n \rightarrow f \text{ uniformly} \Leftrightarrow$$

① $\|f_n - f\|_\infty < \infty$ eventually.

② $\|f_n - f\|_\infty \rightarrow 0$.

Example 5.1.5: $f_n: \mathbb{R} \rightarrow \mathbb{R}$.

$f_1(x) = x$, $f_n(x) = 0$ for all $n \geq 2$.

Then $f_n \rightarrow 0$ uniformly, even though $\|f_1 - 0\|_\infty = \infty$.

Module 2. Examples.

Given the following sequences of functions, find the pointwise limit and determine if the convergence is uniform.

Example 5.2.1: $f_n: (0, 1) \rightarrow \mathbb{R}$.

$$f_n(x) = \frac{nx}{1+nx}.$$

For $x \in (0, 1)$, $f_n(x) \rightarrow 1$. Therefore: $f_n \rightarrow 1$ pointwise.

Claim: this convergence is not uniform.

For $n > 1$.

$$|f_n\left(\frac{1}{n}\right) - 1| = \frac{1}{2}.$$

So, $\|f_n - f\|_\infty \not\rightarrow 0$, and so the convergence is not uniform.

Example 5.2.2: $f_n: \mathbb{C} \rightarrow \mathbb{R}$, $f_n(c_{nk}) = a_n$

For $(c_{nk}) \in \mathbb{C}$, $f_n(c_{nk}) = a_n \rightarrow 0$ as $n \rightarrow \infty$.

So, $f_n \rightarrow 0$ pointwise

For $n \in \mathbb{N}$, $|f_n\left(\underbrace{1, 1, 1, \dots, 1}_{n\text{-times}}, 0, 0, 0, \dots\right) - 0| = |1 - 0| = 1$.

$\Rightarrow \|f_n - 0\|_\infty \geq 1 \Rightarrow \|f_n - 0\|_\infty \not\rightarrow 0$, $f_n \not\rightarrow 0$ uniformly.

Example 5.2.3: $f_n: [0, 1] \times [0, 1] \rightarrow \mathbb{R}$.

$$f_n(a, b) = \frac{a^n}{n} + \frac{1}{b+n}$$

For $(a, b) \in [0, 1] \times [0, 1]$

$$f_n(a, b) = \frac{a^n}{n} + \frac{1}{b+n} \rightarrow 0.$$
$$\rightarrow 0 \quad \rightarrow 0$$

So, $f_n \rightarrow 0$ pointwise.

Note:

$$|f_n(a, b) - 0| = \frac{a^n}{n} + \frac{1}{b+n} \leq \frac{1}{n} + \frac{1}{n} = \frac{2}{n}.$$

Therefore: $\|f_n - 0\|_{\infty} \leq \frac{2}{n} \rightarrow 0$, and so $f_n \rightarrow 0$ uniformly.

Module 3: Theorem A.

Example 5.3.1.

$$f_n: [0, 1] \rightarrow \mathbb{R}.$$

$f_n(x) = x^n$ is continuous.

$$f_n \rightarrow \begin{cases} 1 & x=1 \\ 0 & x \neq 1 \end{cases} \text{ pointwise.}$$

Pointwise limit of continuous functions needs not to be continuous!

Theorem 5.3.2. $(f_n), f_n: A \rightarrow W, A \subseteq V$.

If f_n is continuous for all $n \in \mathbb{N}$ and $f_n \rightarrow f$ uniformly, then f is continuous.

Proof:

Let $(a_n) \subseteq A$ such that $a_n \rightarrow a$ and let $\epsilon > 0$ be given.

We may find $N \in \mathbb{N}$ such that

$$\|f_N - f\|_\infty < \frac{\epsilon}{3}$$

Since f_N is continuous, $\exists M \in \mathbb{N}$ such that

$|f_N(a_n) - f_N(a)| < \frac{\epsilon}{3}$ for all $n \geq M$. Then for $n \geq M$,

$$|f(a_n) - f(a)| \leq$$

$$\begin{aligned} & |f(a_n) - f_N(a_n)| + |f_N(a_n) - f_N(a)| + |f_N(a) - f(a)| \\ & \leq \|f - f_N\|_\infty + |f_N(a_n) - f_N(a)| + \|f_N - f\|_\infty \\ & < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon. \end{aligned}$$

Therefore, $f(a_n) \rightarrow f(a)$ and so f is continuous \square

Module 4.

Theorem 5.4.1. Let $A \subseteq V$ be compact, W be a Banach space, then $(C(CA, W), \| \cdot \|_\infty)$ is a Banach space.

Proof:

Let $(f_n) \subseteq C(CA, W)$ be Cauchy. Let $\varepsilon > 0$ be given. There exists $N \in \mathbb{N}$ such that

$$\|f_n - f_m\|_\infty < \varepsilon$$

for all $n, m \geq N$.

For $x \in A$ and $n, m \geq N$,

$$\begin{aligned} & \|f_n(x) - f_m(x)\| \\ & \leq \|f_n - f_m\|_\infty < \varepsilon. \end{aligned} \quad (*)$$

Hence, $(f_n(x)) \subseteq W$ is Cauchy. Since W is a Banach space, we know that

$$f_n(x) \rightarrow f(x) \in W$$

for some $f(x) \in W$.

We have constructed a function $f: A \rightarrow W$, such that $f_n \rightarrow f$ pointwise.

For $x \in A$ and $n \geq N$, $\lim_{m \rightarrow \infty} \|f_n(x) - f_m(x)\| \leq \varepsilon$.

(limits preserve order).

$$\Rightarrow \|f_n(x) - f(x)\| \leq \varepsilon$$

($f_m(x) \rightarrow f(x)$, $\| \cdot \|$ are continuous).

$$\Rightarrow \|f_n - f\|_\infty \leq \varepsilon.$$

So, $f_n \rightarrow f$ uniformly

By previous theorem, $f \in C(A, W)$ and so $f_n \rightarrow f$
in $C(A, W)$. This completes the proof. \square

Week 6 Notes.

Part 2: Multivariable Calculus.

Module 1. Partial Derivatives.

Remark 6.1.1. Idea.

Given a function $f: A \rightarrow \mathbb{R}^m$, $A \subseteq \mathbb{R}^n$, develop a theory of multivariable differentiation using as much single variable as possible.

Definition 6.1.2.

A scalar function is a function of the form $f: A \rightarrow \mathbb{R}$, $A \subseteq \mathbb{R}^n$.

Remark 6.1.3:

If $f: A \rightarrow \mathbb{R}^m$, $A \subseteq \mathbb{R}^n$ is a function, then $\exists f_1, f_2, \dots, f_m$ on A such that $f = (f_1, f_2, \dots, f_m)$.

Example 6.1.4.

$f: \mathbb{R}^3 \rightarrow \mathbb{R}^2$.

$$f(x, y, z) = (\underbrace{xz e^y}, \underbrace{x^2 + z^2}).$$

$f_1(x, y, z) \quad f_2(x, y, z).$

Definition 6.1.5.

$A \subseteq \mathbb{R}^n$, $f: A \rightarrow \mathbb{R}$. Let $\{e_1, e_2, \dots, e_n\}$ be the standard basis for \mathbb{R}^n . For $1 \leq i \leq n$, we define the i^{th} partial derivative of f at $a = (a_1, a_2, \dots, a_n)$ to A by:

$$f_{x_i}(a) = \frac{\partial f}{\partial x_i}(a)$$

$$:= \lim_{h \rightarrow 0} \frac{f(a+h\mathbf{e}_i) - f(a)}{h}$$

provided the limit exists.

Remark 6.1.6: Big picture.

$a \in A \subseteq \mathbb{R}^n$, $f: A \rightarrow \mathbb{R}$, we write $f(x_1, x_2, \dots, x_n)$.

- ① $f_{x_i}(a)$ is the derivative of f at a with respect to the variable x_i (treating $x_j, j \neq i$, as constants).
- ② $f_{x_i}(a)$ is the slope of the tangent line to the surface $y = f(x_1, x_2, \dots, x_n)$ which is parallel to e_i .

Example 6.1.7: $f: \mathbb{R}^2 \rightarrow \mathbb{R}$. $f(x, y)$.

$$a = (a_1, a_2) \in \mathbb{R}^2.$$

$$\begin{aligned} f_x(a) &= \lim_{h \rightarrow 0} \frac{f(a+h\mathbf{e}_1) - f(a)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(a_1+h, a_2) - f(a_1, a_2)}{h} \end{aligned}$$

Remark 6.1.8:

As usual, we can think of $\frac{\partial f}{\partial x_i}$ as a function

We write:

$$f_{x_i}(x_1, x_2, \dots, x_n)$$

or

$$\frac{\partial}{\partial x_i} f(x_1, x_2, \dots, x_n).$$

Example 6.1.9. $f: \mathbb{R}^3 \rightarrow \mathbb{R}$.

$$f(x, y, z) = xy^2z + e^{xy}.$$

$$f_x(x, y, z) = y^2 z + y e^{xy}$$

$$\frac{\partial f(x, y, z)}{\partial y} = 2xyz + xe^{xy}$$

$$f_z(x, y, z) = xy^2$$

Definition 6.1.10.

$A \subseteq \mathbb{R}^n$, $f: A \rightarrow \mathbb{R}^m$, $f = (f_1, f_2, \dots, f_m)$.

For $a \in A$

$$\begin{aligned} \frac{\partial f}{\partial x_i}(a) &= f_{x_i}(a) \\ &:= \left(\frac{\partial f_1}{\partial x_i}(a), \frac{\partial f_2}{\partial x_i}(a), \dots, \frac{\partial f_m}{\partial x_i}(a) \right) \end{aligned}$$

provided it exists.

Example 6.1.11: $f: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

$$f(x, y) = (2x^2y, 4x, e^{xy})$$

$$f_x(x, y) = (4xy, 4, ye^{xy})$$

$$f_y(x, y) = (2x^2, 0, xe^{xy})$$

Module 2. Differentiability

Remark 6.2.1

Recall for single-variable functions.

$A \subseteq \mathbb{R}$, $f: A \rightarrow \mathbb{R}$. f is differentiable at $a \in A$.
iff.

① $a \in \text{Int}(A)$

② $\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = f'(a)$

exists

$$\Leftrightarrow \lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - mh}{h} = 0.$$

for some $m \in \mathbb{R}$.

Remark 6.2.2

Since $a \in \text{Int}(A)$, for small enough h , $a+h \in A$.

The limit above is well-defined.

Remark 6.2.3

$T: \mathbb{R} \rightarrow \mathbb{R}$ is a linear transformation iff

$$T(x) = mx \text{ for some } m \in \mathbb{R}.$$

Remark 6.2.4: Idea

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - mh}{h} \xrightarrow{T(h)} 0.$$

f over $[a, a+h]$ (or $[a+h, a]$) can be approximated arbitrarily well by the line $T(x) = mx$.

Remark 6.2.5. Notation.

$$L(\mathbb{R}^n, \mathbb{R}^m) = \{T: \mathbb{R}^n \rightarrow \mathbb{R}^m : T \text{ is linear}\}.$$

Definition 6.2.6:

$a \in A \subseteq \mathbb{R}^n$, $f: A \rightarrow \mathbb{R}^m$, we say f is differentiable at $a \in A$ if

① $a \in \text{Int}(A)$

② There exists $T \in L(\mathbb{R}^n, \mathbb{R}^m)$ such that.

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - T(h)}{\|h\|} = 0.$$

Remark 6.2.7:

By ①, $f(a+h)$ is defined for small enough h .

Remark 6.2.8:

① Let $T \in L(\mathbb{R}^n, \mathbb{R}^m)$. Let B be the matrix of T relative to the standard ordered base. $T(X) = B(X)$ for all $X \in \mathbb{R}^n$.

② (Assignment question).

Let $A \in M_{m \times n}(\mathbb{R})$.

$$\|A\|_{\text{op}} = \sup \{ \|Ax\| : x \in \mathbb{R}^n, \|x\|=1 \}.$$

is a norm on $M_{m \times n}(\mathbb{R})$. We call it the operator norm.

③ If $A \in M_{m \times n}(\mathbb{R})$ and $X \in \mathbb{R}^n$. Then.

$$\|Ax\| \leq \|A\|_{\text{op}} \cdot \|x\|.$$

(already shown in assignment).

Theorem 6.2.9.

$a \in A \subseteq \mathbb{R}^n$, $f: A \rightarrow \mathbb{R}^m$. If f is differentiable at a , then f is continuous at a .

Proof Sketch:

f is differentiable at a

$\Rightarrow T \in L(\mathbb{R}^n, \mathbb{R}^m)$

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - T(h)}{\|h\|} = 0.$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - Bh}{\|h\|} = 0.$$

\Rightarrow We may find $\delta > 0$ such that.

If $0 < \|h - 0\| = \|h\| < \delta$, then

$$\left\| \frac{f(a+h) - f(a) - Bh}{\|h\|} \right\| < 1.$$

$$\Rightarrow \|f(a+h) - f(a) - Bh\| < \|h\|.$$

$$\Rightarrow \|f(a+h) - f(a)\| - \|Bh\| < \|h\|.$$

$$\Rightarrow \|f(a+h) - f(a)\| < \|Bh\| + \|h\|.$$

$$\leq \|B\|_{op} \|h\| + \|h\|$$

As $h \rightarrow 0$, $\|B\|_{op} \|h\| + \|h\| \rightarrow 0$

Therefore, $\lim_{h \rightarrow 0} f(a+h) = f(a)$.

Letting $x = a+h$. $\lim_{x \rightarrow a} f(x) = f(a)$, so f is continuous.

□

Definition 6.2.10.

Let $U \subseteq \mathbb{R}^n$ be open, $f: U \rightarrow \mathbb{R}^m$.

We say that f is differentiable on U if f is differentiable at every point in U .

Module 3. Total Derivative.

Example 6.3.1. $f: \mathbb{R}^2 \rightarrow \mathbb{R}$.

$$f(x, y) = \begin{cases} (x^2 + y^2) \sin \frac{1}{\sqrt{x^2 + y^2}} & ((x, y) \neq (0, 0)) \\ 0 & ((x, y) = (0, 0)) \end{cases}$$

f is differentiable at $(0, 0)$.

Consider $B = (0, 0)$.

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0) - Bh}{\|h\|} = \lim_{h \rightarrow 0} \frac{f(h)}{\|h\|}$$

$$= \lim_{(x, y) \rightarrow (0, 0)} (x^2 + y^2) \sin \frac{1}{\sqrt{x^2 + y^2}} \cdot \frac{1}{\sqrt{x^2 + y^2}}$$

$$= \lim_{(x, y) \rightarrow (0, 0)} \sqrt{x^2 + y^2} \sin \frac{1}{\sqrt{x^2 + y^2}}$$

$$= 0 \text{ (Squeeze Theorem)}$$

Remark 6.3.2: Problem: How to find the matrix B ?

Remark 6.3.3: Investigation.

$$a \in A \subseteq \mathbb{R}^n, f: A \rightarrow \mathbb{R}^m$$

Suppose f is differentiable at a , so $\exists B \in M_{m \times n}(\mathbb{R})$

$$\text{such that } \lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - Bh}{\|h\|} = 0.$$

Let $\{e_1, \dots, e_n\}$ be the standard ordered basis.

$$\text{As } t \rightarrow 0, te_i \rightarrow 0 \in \mathbb{R}^n.$$

$$\Rightarrow \lim_{t \rightarrow 0} \frac{f(\mathbf{a} + t\mathbf{e}_i) - f(\mathbf{a}) - B\mathbf{e}_i}{|t|} = 0.$$

Therefore,

$$\lim_{t \rightarrow 0^+} \frac{f(\mathbf{a} + t\mathbf{e}_i) - f(\mathbf{a}) - tB\mathbf{e}_i}{t} = 0.$$

and $\lim_{t \rightarrow 0^+} \frac{tB\mathbf{e}_i}{t} = B\mathbf{e}_i$.

$$\Rightarrow \lim_{t \rightarrow 0^+} \frac{f(\mathbf{a} + t\mathbf{e}_i) - f(\mathbf{a})}{t} = B\mathbf{e}_i.$$

$$\lim_{t \rightarrow 0^-} \frac{f(\mathbf{a} + t\mathbf{e}_i) - f(\mathbf{a}) - tB\mathbf{e}_i}{-t} = 0.$$

$$\Rightarrow \lim_{t \rightarrow 0^-} \frac{f(\mathbf{a} + t\mathbf{e}_i) - f(\mathbf{a})}{t} = B\mathbf{e}_i.$$

$$\Rightarrow B\mathbf{e}_i = \lim_{t \rightarrow 0} \frac{f(\mathbf{a} + t\mathbf{e}_i) - f(\mathbf{a})}{t} = \frac{\partial f}{\partial x_i}(\mathbf{a}).$$

$$= \left(\frac{\partial f_1}{\partial x_i}(\mathbf{a}), \frac{\partial f_2}{\partial x_i}(\mathbf{a}), \dots, \frac{\partial f_m}{\partial x_i}(\mathbf{a}) \right).$$

$= (b_{1i}, b_{2i}, \dots, b_{mi})$. (ith column of B).

$$\Rightarrow B = [b_{ij}] \text{ where } b_{ij} = \frac{\partial f_i}{\partial x_j}(\mathbf{a}).$$

Definition 6.3.4 :

$\mathbf{a} \in A \subseteq \mathbb{R}^n$, $f: A \rightarrow \mathbb{R}^m$. We call the matrix

$$Df(\mathbf{a}) = \left[\frac{\partial f_i}{\partial x_j}(\mathbf{a}) \right]_{m \times n}.$$

the total derivative of f at \mathbf{a} , provided it exists.

Theorem 6.3.5:

$a \in A \subseteq \mathbb{R}^n, f: A \rightarrow \mathbb{R}^m$. If f is differentiable at a , then:

① for all $1 \leq j \leq n$. $\frac{\partial f}{\partial x_j}(a)$ exists ($= B_{1j}$).

② $\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - Df(a)h}{\|h\|} = 0$.

Definition 6.3.6: $a \in A \subseteq \mathbb{R}^n, f: A \rightarrow \mathbb{R}$.

We call $Df(a)$ the gradient of f at a and label it by $\nabla f(a)$.

$$\nabla f(a) = \left(\frac{\partial f}{\partial x_1}(a), \frac{\partial f}{\partial x_2}(a), \dots, \frac{\partial f}{\partial x_n}(a) \right).$$

Module 4. Continuous Partials.

Example 6.4.1.

$$f(x,y) = \begin{cases} \frac{xy}{x^2+y^2} & (x,y) \neq (0,0) \\ 0 & (x,y) = (0,0) \end{cases}$$

$$fx(0,0) = 0.$$

$$fy(0,0) = 0 \text{ by symmetry.}$$

$$\nabla f(0,0) = 0.$$

But f is not even continuous, so not differentiable.

$$(\frac{1}{n}, \frac{1}{n}) \rightarrow 0, f(\frac{1}{n}, \frac{1}{n}) = \frac{1}{2} \rightarrow \frac{1}{2}.$$

Theorem 6.4.2.

$U \subseteq \mathbb{R}^n$ open, $f: U \rightarrow \mathbb{R}$

If $a \in U$ and $\forall 1 \leq j \leq n$, if $\frac{\partial f}{\partial x_j}$ exist on U and are continuous at a , then f is differentiable at a .

Proof: Appendix.

Note: On assignment, this will be extended to $f: U \rightarrow \mathbb{R}^m$.

Example 6.4.3. $U = \mathbb{R}^2 \setminus \{(0,0)\}$.

Is $f: U \rightarrow \mathbb{R}$ given by $f(x,y) = \frac{\sin(xy)}{x^2+y^2}$ differentiable on U ?

$$fx(x,y) = \frac{(x^2+y^2)\cos(xy)y - 2x\sin(xy)}{(x^2+y^2)^2}$$

\hookrightarrow exists on U and is continuous on U .

By symmetry, $f_y(x, y)$ exists on U and is continuous.

By the Theorem, f is differentiable on U .

Remark 6-4-4.

Do not use the false converse of this theorem!

If $f: U \rightarrow \mathbb{R}$ is differentiable at $a \in U$, then there may exist a $\frac{\partial f}{\partial x_j}$ which is discontinuous at a .

Example 6-4-5: [Appendix].

$$f(x, y) = \begin{cases} (x^2 + y^2) \sin \frac{1}{\sqrt{x^2 + y^2}} & ((x, y) \neq (0, 0)) \\ 0 & ((x, y) = (0, 0)) \end{cases}$$

Week 6 Appendix

Theorem. Let $a \in U \subseteq \mathbb{R}^n$ be open and let $f : U \rightarrow \mathbb{R}$ be a function. If all of the partial derivatives of f exist on U (ie. the total derivative exists on U) **and are continuous at a** , then f is differentiable at a .

Proof. Suppose every partial derivative of f exists on U and that every partial derivative is continuous at a .

Suppose $a = (a_1, a_2, \dots, a_n)$. Since U is open there exists $r > 0$ such that $B_r(a) \subseteq U$. For any $h = (h_1, h_2, \dots, h_n) \neq 0$ such that $a + h \in B_r(a)$,

$$\begin{aligned}
 f(a + h) - f(a) &= f(a_1 + h_1, \dots, a_n + h_n) - f(a_1, \dots, a_n) \\
 &= f(a_1 + h_1, \dots, a_n + h_n) - f(a_1, a_2 + h_2, \dots, a_n + h_n) \\
 &\quad + f(a_1, a_2 + h_2, \dots, a_n + h_n) - f(a_1, a_2, a_3 + h_3, \dots, a_n + h_n) \\
 &\quad + f(a_1, a_2, a_3 + h_3, \dots, a_n + h_n) - f(a_1, a_2, a_3, a_4 + h_4, \dots, a_n + h_n) \\
 &\quad \vdots \\
 &\quad + f(a_1, \dots, a_{n-1}, a_n + h_n) - f(a_1, a_2, \dots, a_n).
 \end{aligned}$$

However, by the single variable Mean Value Theorem, for every $1 \leq j \leq n$ there exists c_j between a_j and $a_j + h_j$ such that

$$\begin{aligned}
 &\frac{f(a_1, \dots, a_{j-1}, a_j + h_j, \dots, a_n + h_n) - f(a_1, \dots, a_j, a_{j+1} + h_{j+1}, \dots, a_n + h_n)}{a_j + h_j - a_j} \\
 &= \frac{\partial f}{\partial x_j}(a_1, \dots, a_{j-1}, c_j, a_{j+1} + h_{j+1}, \dots, a_n + h_n).
 \end{aligned}$$

Putting all of this mess together,

$$f(a + h) - f(a) = \sum_{j=1}^n h_j \frac{\partial f}{\partial x_j}(a_1, \dots, a_{j-1}, c_j, a_{j+1} + h_{j+1}, \dots, a_n + h_n).$$

Now, for $1 \leq j \leq n$ let

$$\delta_j := \frac{\partial f}{\partial x_j}(a_1, \dots, a_{j-1}, c_j, a_{j+1} + h_{j+1}, \dots, a_n + h_n) - \frac{\partial f}{\partial x_j}(a_1, \dots, a_n),$$

and $\delta = (\delta_1, \dots, \delta_n)$. Then,

$$f(a + h) - f(a) - \nabla f(a) \cdot h = h \cdot \delta.$$

Since all of the partials are continuous at a , as $h \rightarrow 0$, each $\delta_j \rightarrow 0$, and so $\delta \rightarrow 0$ in \mathbb{R}^n . Therefore

$$0 \leq \lim_{h \rightarrow 0} \frac{|f(a + h) - f(a) - \nabla f(a) \cdot h|}{\|h\|} = \lim_{h \rightarrow 0} \frac{|\delta \cdot h|}{\|h\|} \leq \lim_{h \rightarrow 0} \frac{\|\delta\| \cdot \|h\|}{\|h\|} = 0.$$

Note that in the last inequality, we used the Cauchy-Schwarz inequality from linear algebra!. Therefore

$$\lim_{h \rightarrow 0} \frac{|f(a + h) - f(a) - \nabla f(a) \cdot h|}{\|h\|} = 0$$

and so

$$\lim_{h \rightarrow 0} \frac{f(a + h) - f(a) - \nabla f(a) \cdot h}{\|h\|} = 0$$

as well. This exactly means that f is differentiable at a .

□

Example. Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$f(x, y) = \begin{cases} (x^2 + y^2) \sin\left(\frac{1}{\sqrt{x^2 + y^2}}\right) & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

From the module videos, we know that f is differentiable at $(0, 0)$. Observe that

$$f_x(x, y) = 2x \sin\left(\frac{1}{\sqrt{x^2 + y^2}}\right) - \cos\left(\frac{1}{\sqrt{x^2 + y^2}}\right) \frac{x}{\sqrt{x^2 + y^2}},$$

for all $(x, y) \neq (0, 0)$. Now, we see that $(1/n, 0) \rightarrow (0, 0)$ but

$$f_x(1/n, 0) = \frac{2}{n} \sin(n) - \cos(n)$$

diverges. Therefore f_x is not continuous at $(0, 0)$.

Week 7 Notes.

Module 1 Differentiation Rules.

Remark 7.1.1: General Strategy.

$f: U \rightarrow \mathbb{R}^n$, $a \in U \subseteq \mathbb{R}^n$. U is open.

Suppose $\exists A \in M_{m \times n}(\mathbb{R})$ such that

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - Ah}{\|h\|} = 0.$$

Then:

① f is differentiable at $a \in U$.

$$\textcircled{2} \quad \lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - Ah}{\|h\|}$$

$$= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - Df(a)h}{\|h\|}$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{(Df(a) - A)h}{\|h\|} = 0.$$

$$\Rightarrow \lim_{t \rightarrow 0^+} \frac{(Df(a) - A)te_i}{|t|} = 0.$$

$$\Rightarrow (Df(a) - A)e_i = 0.$$

$$\Rightarrow Df(a) - A = 0.$$

$$\Rightarrow Df(a) = A$$

Proposition 7.1.2: [Sum and Scalar Multiplication rule].

$f, g: A \rightarrow \mathbb{R}^m$, differentiable at $a \in A \subseteq \mathbb{R}^n$.

For $\alpha \in \mathbb{R}$, $f + \alpha g$ is differentiable at a and.

$$D(f + \alpha g)(a) = Df(a) + \alpha Dg(a).$$

Proof:

Let $P = f + \alpha g$

Then,

$$\begin{aligned} & \lim_{h \rightarrow 0} \frac{P(a+h) - P(a) - (Df(a) + \alpha Dg(a))h}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - Df(a)h}{\|h\|} + \alpha \lim_{h \rightarrow 0} \frac{g(a+h) - g(a) - Dg(a)h}{\|h\|} \\ &= 0 + \alpha 0 = 0. \end{aligned}$$

Proposition 7.1.3. [Dot product rule].

$f, g: A \rightarrow \mathbb{R}^m$ differentiable at $a \in A \subseteq \mathbb{R}^n$.

Consider $f \cdot g: A \rightarrow \mathbb{R}$ defined by

$$(f \cdot g)(x) = f(x) \cdot g(x).$$

Then, $f \cdot g$ is differentiable at a and

$$D(f \cdot g)(a) = g(a)Df(a) + f(a)Dg(a).$$

Proof: Appendix.

Module 2. Chain Rule.

Theorem 7.2.1.

$A \subseteq \mathbb{R}^n, B \subseteq \mathbb{R}^m$

$f: A \rightarrow \mathbb{R}^m, g: B \rightarrow \mathbb{R}^k, f(A) \subseteq B$

If f is differentiable at $a \in A$, and g is differentiable at $f(a) \in B$, then $g \circ f$ is differentiable at a , with

$$D(g \circ f)(a) = Dg(f(a)) \cdot Df(a).$$

Proof: Appendix.

Example 7.2.2. Let $f(x, y, z)$ be real-valued, differentiable. $x(t_1, t_2), y(t_1, t_2), z(t_1, t_2)$.

Then:

$$p(t_1, t_2) = (x(t_1, t_2), y(t_1, t_2), z(t_1, t_2)).$$

is differentiable.

Moreover:

$$\begin{aligned} D(f \circ p)(t_1, t_2) &= Df(p(t_1, t_2)) Dp(t_1, t_2) \\ \Rightarrow \nabla f(t_1, t_2) &= Df(x, y, z) \cdot Dp(t_1, t_2) \\ \Rightarrow \left[\frac{\partial f}{\partial t_1}, \frac{\partial f}{\partial t_2} \right] &= \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] \begin{bmatrix} \frac{\partial x}{\partial t_1} & \frac{\partial x}{\partial t_2} \\ \frac{\partial y}{\partial t_1} & \frac{\partial y}{\partial t_2} \\ \frac{\partial z}{\partial t_1} & \frac{\partial z}{\partial t_2} \end{bmatrix} \end{aligned}$$

$$\text{Therefore: } \frac{\partial f}{\partial t_1} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial t_1} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial t_1} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial t_1}.$$

$$\frac{\partial f}{\partial t_2} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial t_2} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial t_2} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial t_2}.$$

(This is what is on usual multi-variable calculus book).

Module 3. Mean Value Theorem

Remark 7.3.1.

Recall for single-valued functions:

$f: [a, b] \rightarrow \mathbb{R}$ is continuous. If f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f'(c) = \frac{f(b) - f(a)}{b - a}$.

Remark 7.3.2: Naive approach.

$U \subset \mathbb{R}^n$ is open, $f: U \rightarrow \mathbb{R}^m$ is differentiable. If $a, b \in U$, then there exists $c \in L[a, b] = \{c(1-t)a + tb : t \in [0, 1]\}$ such that

$$f(a) - f(b) = Df(c)(b-a).$$

Example 7.3.3: The above is false, unfortunately.

$$f: \mathbb{R} \rightarrow \mathbb{R}^2$$

$$f(x) = (\cos x, \sin x).$$

$$f(0) = f(2\pi) = (1, 0).$$

$$\text{But, } Df(x) = \begin{bmatrix} -\sin x \\ \cos x \end{bmatrix} \neq 0.$$

$$\text{So, } 0 \neq 2\pi(-\sin x, \cos x)^T.$$

for all $x \in \mathbb{R}$.

Remark 7.3.4: Idea.

We will work "one direction at a time".

for all $x \in \mathbb{R}^m$, find $c \in L(a, b)$ such that

$$x \cdot (f(b) - f(a) - Df(c)(b-a)) = 0.$$

$$\Leftrightarrow x \cdot (f(b) - f(a)) = x \cdot (Df(c)(b-a))$$

Lemma 7.3.5: $a, b \in \mathbb{R}^n$

The function $\varphi: \mathbb{R} \rightarrow \mathbb{R}^n$

$$\varphi(t) = (1-t)a + tb$$

is differentiable with

$$D\varphi(t) = b - a.$$

Proof:

$$\begin{aligned} & \lim_{h \rightarrow 0} \frac{\varphi(t+h) - \varphi(t) - (b-a)h}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{(1-t-h)a + (t+h)b - (1-t)a - tb - (b-a)h}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{-ha + hb - (b-a)h}{\|h\|} \\ &= 0. \end{aligned}$$

Theorem 7.3.6 [Mean Value Theorem].

Let $U \subseteq \mathbb{R}^n$ be open. If $f: U \rightarrow \mathbb{R}^m$ is differentiable and $a, b \in U$ such that $L(a, b) \subseteq U$, then for all $x \in \mathbb{R}^m$, there exists $c \in L(a, b)$ such that

$$x \cdot (f(b) - f(a)) = x \cdot (Df(c)(b-a)).$$


(c depends on the x chosen).

Proof: Let $L(a, b) \subseteq U$, $x \in \mathbb{R}^m$

$$\textcircled{1} \quad \varphi(t) = (1-t)a + tb.$$

$$\varphi([0,1]) = L(a, b) \subseteq U \quad (U \text{ is open}).$$

$\Rightarrow \exists \delta > 0$ such that $\varphi(0-\delta, 1+\delta) \subseteq U$.



\textcircled{2} For $t \in (-\delta, 1+\delta)$,

$$D(f \circ \varphi)(t) = Df(\varphi(t))(b-a)$$

\textcircled{3} $F: (-\delta, 1+\delta) \rightarrow \mathbb{R}$

$$F(t) = \chi \cdot (f \circ \varphi)(t).$$

By Dot product rule:

$$F'(t) = \chi \cdot Df(\varphi(t))(b-a).$$

\textcircled{4} By single value mean value theorem.

$\exists t_0 \in (0, 1)$ such that

$$F(1) - F(0) = F'(t_0)(1-0)$$

$$\Rightarrow \chi \cdot f(b) - \chi \cdot f(a) = \chi \cdot Df(\varphi(t_0))(b-a).$$

$$\Rightarrow \chi \cdot (f(b) - f(a)) = \chi \cdot Df(\varphi(t_0))(b-a).$$

Choose $c = \varphi(t_0)$, we are done. □

Module 4. Tangent Hyperplanes.

Remark 7-4-1: Goal

Understand the geometrical interpretation of the total derivative (gradient) of scalar functions:

$$f: U \rightarrow \mathbb{R}$$

where $U \subseteq \mathbb{R}^n$ is open.

Remark 7-4-2: Motivation.

$n=1$. Let $U \subseteq \mathbb{R}$ be open.

If $f: U \rightarrow \mathbb{R}$ is differentiable at $a \in U$, then $f'(a) = Df(a)$ is the slope of the tangent line to the curve $y = f(x)$ at $x = a$.

When $n=2$, $U \subseteq \mathbb{R}^2$ is open, we want:

If $f: U \rightarrow \mathbb{R}$ is differentiable at $a \in U$, then

$Df(a) = Df(a)$ tells us information about the tangent plane to the surface $z = f(x, y)$ at $(x, y) = a$.

Definition 7-4-3.

A hyperplane in \mathbb{R}^n is a set of the form

$$P = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : a_1x_1 + a_2x_2 + \dots + a_nx_n = d\}$$

for some fixed $a_1, a_2, \dots, a_n \in \mathbb{R}$ (not all zero) and $d \in \mathbb{R}$.

Remark 7-4-4.

$n=2$, hyperplanes are lines.

$n=3$, hyperplanes are planes.

Example 7-4-5: $P = \{(x, y, z) \in \mathbb{R}^3 : 2x + y - 3z = 13\}$

Definition 7-4-6.

Let $P = \{(x_1, \dots, x_n) : a_1 x_1 + \dots + a_n x_n = d\}$ be a hyperplane in \mathbb{R}^n . We call $n = (a_1, a_2, \dots, a_n)$ the normal (vector) of P .

Remark 7-4-7: Geometrically:

Let $b = (b_1, b_2, \dots, b_n) \in P$.

Then, $d = a_1 b_1 + a_2 b_2 + \dots + a_n b_n$

Therefore, $x = (x_1, \dots, x_n) \in P$

$$\Leftrightarrow d = a_1 x_1 + a_2 x_2 + \dots + a_n x_n.$$


$$\Leftrightarrow 0 = d - d$$

$$= a_1(x_1 - b_1) + \dots + a_n(x_n - b_n)$$

$$= n \cdot (x - b).$$

So, $P = \{x \in \mathbb{R}^n : n \cdot (x - b) = 0\}$.

i.e. $x \in P$ iff n is orthogonal/prependicular to $x - b$.



Definition 7-4-8: $A \subseteq \mathbb{R}^n$, $a \in A$.

A hyperplane $P \subseteq \mathbb{R}^n$ with normal n is said to be tangent to A at a if


$$h \cdot \frac{a_k - a}{\|a_k - a\|} \rightarrow 0.$$

for all sequences $(a_k) \subseteq A \setminus \{a\}$ such that $a_k \rightarrow a$.


Remark 7.4.9: Why is this a good definition?

Recall that $a, b \in \mathbb{R}^n$ are orthogonal iff $a \cdot b = 0$.

Then, $h \cdot \frac{a_k - a}{\|a_k - a\|} \rightarrow 0$ says that unit (length 1) vectors in the direction of $a_k - a$:



become closer and closer to being orthogonal to h as $k \rightarrow \infty$.



Theorem 7.4.10:

$U \subseteq \mathbb{R}^n$ open, $a \in U$, $f: U \rightarrow \mathbb{R}$.

If f is differentiable at a , then the surface

$$S = \{(x, z) \in \mathbb{R}^{n+1} : z = f(x), x \in U\}.$$

has a tangent hyperplane at $(a, f(a))$ with the normal

$$n = (\nabla f(a), -1)$$

Proof: Appendix.

Example 7.4.11:

Find the tangent plane to the surface.

$$z = 2x^2 + y^2$$

at $(1, 1, 3)$.

Consider $f: \mathbb{R}^2 \rightarrow \mathbb{R}$

$$f(x, y) = 2x^2 + y^2$$

Note f_x, f_y exist and are continuous on \mathbb{R}^2 .

So, f is differentiable on \mathbb{R}^2 .

$$\nabla f(x, y) = (4x, 2y)$$

$$\nabla f(1, 1) = (4, 2)$$

The normal is:

$$n = (4, 2, -1)$$

The equation of the tangent plane is:

$$4x + 2y - z = d$$

$$\Rightarrow d = 4 \cdot 1 + 2 \cdot 1 - 3 = 3$$

$$\Rightarrow 4x + 2y - z = 3$$

Week 7 Appendix

Theorem (Dot Product Rule). Let $A \subseteq \mathbb{R}^n$ and let f and g be functions from A to \mathbb{R}^m . If f and g are differentiable at $a \in A$ then $f \cdot g$ is differentiable at a and

$$D(f \cdot g)(a) = g(a)Df(a) + f(a)Dg(a).$$

Proof. We must prove that

$$\lim_{h \rightarrow 0} \frac{(f \cdot g)(a + h) - (f \cdot g)(a) - Xh}{\|h\|} = 0,$$

where $X = g(a)Df(a) + f(a)Dg(a)$. To ease notations, let

$$\varepsilon(h) = f(a + h) - f(a) - Df(a)h$$

and

$$\delta(h) = g(a + h) - g(a) - Dg(a)h.$$

Since f and g are differentiable at a we have that

$$\lim_{h \rightarrow 0} \frac{\varepsilon(h)}{\|h\|} = \lim_{h \rightarrow 0} \frac{\delta(h)}{\|h\|} = 0.$$

Then,

$$\begin{aligned} & \lim_{h \rightarrow 0} \frac{(f \cdot g)(a + h) - (f \cdot g)(a) - Xh}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{(f \cdot g)(a + h) - (f \cdot g)(a) - g(a)Df(a)h - f(a)Dg(a)h}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{g(a) \cdot \varepsilon(h) + f(a) \cdot \delta(h)}{\|h\|} \\ &+ \lim_{h \rightarrow 0} \frac{f(a) \cdot g(a) - g(a) \cdot f(a + h) - f(a) \cdot g(a + h) + f(a + h) \cdot g(a + h)}{\|h\|} \\ &= 0 + \lim_{h \rightarrow 0} \frac{f(a) \cdot g(a) - g(a) \cdot f(a + h) - f(a) \cdot g(a + h) + f(a + h) \cdot g(a + h)}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{g(a) \cdot (f(a) - f(a + h)) - g(a + h) \cdot (f(a) - f(a + h))}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{(g(a) - g(a + h)) \cdot (f(a) - f(a + h))}{\|h\|}. \end{aligned}$$

However, by the Cauchy-Schwarz inequality,

$$\frac{|(g(a) - g(a+h)) \cdot (f(a) - f(a+h))|}{\|h\|} \leq \frac{\|g(a) - g(a+h)\| \cdot \|f(a) - f(a+h)\|}{\|h\|}.$$

Therefore,

$$\begin{aligned} 0 &\leq \lim_{h \rightarrow 0} \frac{|(f \cdot g)(a+h) - (f \cdot g)(a) - Xh|}{\|h\|} \\ &\leq \lim_{h \rightarrow 0} \frac{\|g(a) - g(a+h)\| \cdot \|f(a) - f(a+h)\|}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{\|g(a) - g(a+h)\|}{\|h\|} \cdot \frac{\|f(a) - f(a+h)\|}{\|h\|} \|h\| \\ &= \lim_{h \rightarrow 0} \frac{\|g(a) - g(a+h) - Dg(a)h + Dg(a)h\|}{\|h\|} \cdot \frac{\|f(a) - f(a+h) - Df(a)h + Df(a)h\|}{\|h\|} \|h\| \\ &\leq \lim_{h \rightarrow 0} \frac{\|g(a) - g(a+h) - Dg(a)h\| + \|Dg(a)h\|}{\|h\|} \cdot \frac{\|f(a) - f(a+h) - Df(a)h\| + \|Df(a)h\|}{\|h\|} \|h\| \\ &\leq \lim_{h \rightarrow 0} \frac{\|g(a) - g(a+h) - Dg(a)h\| + \|Dg(a)\|_{op} \|h\|}{\|h\|} \cdot \frac{\|f(a) - f(a+h) - Df(a)h\| + \|Df(a)\|_{op} \|h\|}{\|h\|} \|h\| \\ &= \lim_{h \rightarrow 0} (0 + \|Dg(a)\|_{op})(0 + \|Df(a)\|_{op}) \|h\| = 0 \end{aligned}$$

The result follows. \square

Theorem (Chain Rule). Let $A \subseteq \mathbb{R}^n$ and $B \subseteq \mathbb{R}^m$ and consider two functions $f : A \rightarrow \mathbb{R}^m$ and $g : B \rightarrow \mathbb{R}^k$. If f is differentiable at $a \in A$ and g is differentiable at $f(a) \in B$ then $g \circ f$ is differentiable at a with

$$D(g \circ f)(a) = Dg(f(a))Df(a).$$

Proof. We must prove that

$$\lim_{h \rightarrow 0} \frac{(g \circ f)(a + h) - (g \circ f)(a) - Xh}{\|h\|} = 0,$$

where $X = Dg(f(a))Df(a)$. To ease notation, let $b = f(a)$,

$$\varepsilon(h) = f(a + h) - f(a) - Df(a)h,$$

$$\delta(k) = g(b + k) - g(b) - Dg(b)k,$$

so that

$$\lim_{h \rightarrow 0} \frac{\varepsilon(h)}{\|h\|} = 0$$

and

$$\lim_{k \rightarrow 0} \frac{\delta(k)}{\|k\|} = 0.$$

Now, consider $k = f(a + h) - f(a)$. Note that $k \rightarrow 0$ as $h \rightarrow 0$, by continuity of f at a . Then,

$$\begin{aligned} & \lim_{h \rightarrow 0} \frac{(g \circ f)(a + h) - (g \circ f)(a) - Xh}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{(g \circ f)(a + h) - (g \circ f)(a) - Dg(f(a))Df(a)h}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{g(k + b) - g(b) - Dg(b)Df(a)h}{\|h\|} \\ &= \lim_{h \rightarrow 0} \frac{Dg(b)\varepsilon(h) + \delta(k)}{\|h\|} \\ &= \lim_{h \rightarrow 0} Dg(b) \frac{\varepsilon(h)}{\|h\|} + \frac{\delta(k)}{\|h\|}. \end{aligned}$$

Now, since

$$0 \leq \frac{\|Dg(b)\varepsilon(h)\|}{\|h\|} \leq \|Dg(b)\|_{op} \frac{\|\varepsilon(h)\|}{\|h\|} \rightarrow 0,$$

as $h \rightarrow 0$ we see that

$$\lim_{h \rightarrow 0} Dg(b) \frac{\varepsilon(h)}{\|h\|} = 0.$$

Next,

$$\lim_{h \rightarrow 0} \frac{\delta(k)}{\|h\|} = \lim_{h \rightarrow 0} \frac{\delta(k)}{\|k\|} \cdot \frac{\|k\|}{\|h\|}.$$

However,

$$\|k\| = \|Df(a)h + \varepsilon(h)\| \leq \|Df(a)\|_{op} \|h\| + \|\varepsilon(h)\|,$$

from which it follows that

$$\frac{\|k\|}{\|h\|}$$

is bounded. By a squeeze theorem argument,

$$\lim_{h \rightarrow 0} \frac{\delta(k))}{\|h\|} = \lim_{h \rightarrow 0} \frac{\delta(k)}{\|k\|} \cdot \frac{\|k\|}{\|h\|} = 0,$$

as required. □

Theorem. Let a be an element of an open set $U \subseteq \mathbb{R}^n$ and let $f : U \rightarrow \mathbb{R}$ be a function. If f is differentiable at a then the surface

$$S = \{(x, z) \in \mathbb{R}^{n+1} : z = f(x), x \in U\},$$

has a tangent hyperplane at $(a, f(a))$ with normal $n = (\nabla f(a), -1)$.

Proof. Let $(x_k, f(x_k)) \subseteq S \setminus \{(a, f(a))\}$ be a sequence such that $(x_k, f(x_k)) \rightarrow (a, f(a))$. By A1, $x_k \rightarrow a$. We must prove that

$$\lim_{k \rightarrow \infty} n \cdot \frac{(x_k, f(x_k)) - (a, f(a))}{\|(x_k, f(x_k)) - (a, f(a))\|} = 0.$$

Since f is differentiable at a we have that

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - \nabla f(a)h}{\|h\|} = 0.$$

Letting $\varepsilon(h) = f(a+h) - f(a) - \nabla f(a)h$,

$$\lim_{h \rightarrow 0} \frac{\varepsilon(h)}{\|h\|} = 0.$$

Moreover, we see that

$$\|(x_k, f(x_k)) - (a, f(a))\|^2 = \|(x_k - a, f(x_k) - f(a))\|^2 \geq \|x_k - a\|^2.$$

Then, since $x_k - a \rightarrow 0$,

$$\begin{aligned} 0 &\leq \left| \lim_{k \rightarrow \infty} n \cdot \frac{(x_k, f(x_k)) - (a, f(a))}{\|(x_k, f(x_k)) - (a, f(a))\|} \right| \\ &= \lim_{k \rightarrow \infty} \frac{|\nabla f(a)(x_k - a) - (f(x_k) - f(a))|}{\|(x_k, f(x_k)) - (a, f(a))\|} \\ &\leq \lim_{k \rightarrow \infty} \frac{|\nabla f(a)(x_k - a) - (f(x_k) - f(a))|}{\|x_k - a\|} \\ &= \lim_{k \rightarrow \infty} \frac{|\varepsilon(x_k - a)|}{\|x_k - a\|} \\ &= 0. \end{aligned}$$

The result follows. □

Week 8 Notes.

Module 1 Higher Order Total Derivatives.

Remark 8.1.1: Goal.

Come up with a sensible definition of an n^{th} order total derivative of a scalar function.

Remark 8.1.2: Idea.

Use n^{th} order partial derivative.

Remark 8.1.3: Notation

$f: U \rightarrow \mathbb{R}$ is differentiable at $a \in U$. We may think of $\nabla f(a)$ as a function via matrix multiplication:

$$\nabla f(a): \mathbb{R}^n \rightarrow \mathbb{R}.$$

$$\nabla f(a)(h_1, h_2, \dots, h_n) = \frac{\partial f}{\partial x_1}(a)h_1 + \dots + \frac{\partial f}{\partial x_n}(a)h_n.$$

Definition 8.1.4:

Let $U \subseteq \mathbb{R}^n$ be open. $f: U \rightarrow \mathbb{R}$, $k \in \mathbb{N}$. Assume all partials of order less or equal to k exist at $a \in U$. We define the k^{th} order total derivative of f at a by:

$$D^k f(a): \mathbb{R}^n \rightarrow \mathbb{R},$$

$$D^k f(a)(h_1, \dots, h_n) = \sum_{i_1=1}^n \dots \sum_{i_k=1}^n \frac{\partial^k f}{\partial x_{i_1} \dots \partial x_{i_k}}(a)h_{i_1}h_{i_2} \dots h_{i_k}.$$

Example 8.1.5: $f: \mathbb{R}^2 \rightarrow \mathbb{R}$.

$$\begin{aligned} D^2 f(a)(h_1, h_2) = & f_{xx}(a) h_1^2 + f_{xy}(a) h_1 h_2 \\ & + f_{yx}(a) h_2 h_1 + f_{yy}(a) h_2^2. \end{aligned}$$

Module 8.2: Taylor's Theorem

The purpose of this short, written module is to prove that there is a multivariable analogue of Taylor's theorem. This result allows us to approximate a function, under certain circumstances, using higher order total derivatives and Taylor-like series.

Theorem. (Taylor's Theorem) Let $p \in \mathbb{N}$, $U \subseteq \mathbb{R}^n$ be open, and $f \in C^p(U)$. For all $x, a \in U$ with $L(x, a) \subseteq U$, there exists $c \in L(x, a)$ such that

$$f(x) = f(a) + \sum_{k=1}^{p-1} \frac{1}{k!} D^k f(a)(x-a) + \frac{1}{p!} D^p f(c)(x-a).$$

Proof. Let $x, a \in U$ and consider $h = x - a = (h_1, \dots, h_n)$. Since $L(x, a) \subseteq U$ and U is open, there exists $\delta > 0$ such that $a + th \in U$ for all $t \in I := (-\delta, 1 + \delta)$. Now, the function $g : I \rightarrow \mathbb{R}$ given by $g(t) = f(a + th)$ is differentiable by the chain rule and

$$g'(t) = Df(a + th)h = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(a + th)h_i.$$

Moreover, it may be shown by induction that for $1 \leq j \leq p$,

$$g^{(j)}(t) = \sum_{i_1=1}^n \cdots \sum_{i_j=1}^n \frac{\partial^j f}{\partial x_{i_1} \partial x_{i_2} \cdots \partial x_{i_j}}(a + th)h_{i_1} \cdots h_{i_j}.$$

Note that this is the motivation for the definition for the higher-order total derivative! In particular, for $1 \leq j \leq p-1$ we have that

$$g^{(j)}(0) = D^j f(a)(h)$$

and

$$g^{(p)}(t) = D^p f(a + th)(h).$$

Therefore $g : I \rightarrow \mathbb{R}$ is p -times differentiable and so by the 1D version of Taylor's Theorem,

$$g(1) - g(0) = \sum_{k=1}^{p-1} \frac{1}{k!} g^{(k)}(0) + \frac{1}{p!} g^{(p)}(t),$$

for some $0 \leq t \leq 1$. Thus,

$$f(x) - f(a) = f(a + h) - f(a) = \sum_{k=1}^{p-1} \frac{1}{k!} D^k f(a)(h) + \frac{1}{p!} D^p f(a + th)(h),$$

and so we are done by taking $c = a + th$. □

Module 3. Optimization.

Definition 8.3.1.

Let $U \subseteq \mathbb{R}^n$ be open, $f: U \rightarrow \mathbb{R}$. Let $a \in U$.

- ① $f(a)$ is a local maxima (resp. local minima) of f if there exists $r > 0$ such that $f(x) \leq f(a)$ (resp. $f(x) \geq f(a)$) for all $x \in B_r(a)$.
- ② $f(a)$ is a local extrema if it is a local minima or maxima.

Remark 8.3.2.

Let U be open, $f: U \rightarrow \mathbb{R}$ is differentiable at $a \in U$.

Assume $f(a)$ is a local extrema of f , say $a = (a_1, \dots, a_n)$. Then


$$g_i(t) = f(a_1, \dots, a_{i-1}, t, a_{i+1}, \dots, a_n).$$

has a local extrema at $t = a_i$. Hence $g'_i(a_i) = 0$.
i.e. $\frac{\partial f}{\partial x_i}(a) = 0$. This implies $\nabla f(a) = 0$.

Example 8.3.3.

$$f(x, y) = x^2 + y^2. \quad \nabla f(x, y) = (2x, -2y).$$

$\Rightarrow \nabla f(0, 0) = (0, 0)$. But $f(0, 0)$ is not an extrema.



This is something called a saddle point. A saddle point is a point where the gradient is zero but neither a maxima nor a minima.

Theorem 8.3.4 [2nd Derivative Test].

Let $U \subseteq \mathbb{R}^n$ be open, $f \in C^2(U)$, at U . If $Df(a) = 0$, then:

- ① $\forall n \neq 0, D^2f(a)(n) > 0 \Rightarrow f(a)$ is a local minima.
- ② $\forall n \neq 0, D^2f(a)(n) < 0 \Rightarrow f(a)$ is a local maxima.
- ③ $\exists h, k \in \mathbb{R}^n$ such that.

$$D^2f(a)(h) > 0, D^2f(a)(k) < 0.$$

$\Rightarrow a$ is a saddle point.

Proof: Appendix

Lemma 8.3.5: $a, b, c \in \mathbb{R}, D := b^2 - ac$.

Consider $\varphi: \mathbb{R}^2 \rightarrow \mathbb{R}$.

$$\varphi(h, k) = ah^2 + 2bhk + ck^2.$$

- ① If $D < 0$, then $a, \varphi(h, k)$ have the same sign for all $(h, k) \neq 0$.
- ② If $D > 0$, then $\varphi(h, k)$ takes on both positive and negative values.

Proof: Omitted.

Theorem 8.3.6.

Let $U \subseteq \mathbb{R}^2$ be open, $f \in C^2(U)$, $Df(a, b) = 0$.

$$D := f_{xy}(a, b)^2 - f_{xx}(a, b)f_{yy}(a, b)$$

(called the discriminant).

- ① If $D < 0$ and $f_{xx}(a, b) > 0$, then $f(a, b)$ is a local minima.
- ② If $D < 0$ and $f_{xx}(a, b) < 0$, then $f(a, b)$ is a local maxima.
- ③ If $D > 0$, then (a, b) is a saddle point.

Proof Sketch.

Let $a = f_{xx}(a, b)$, $b = f_{xy}(a, b)$, $c = f_{yy}(a, b)$

$\varphi(h, k) = D^2 f(a) (h, k)$ in the Lemma.

Module 4: Examples

Example 8.4.1.

Classify all local extrema and/or saddle points

$$\text{of } f(x,y) = x^4 + y^4 - 4xy + 2.$$

Solution:

$$\nabla f(x,y) = (4x^3 - 4y, 4y^3 - 4x).$$

$$\nabla f(x,y) = 0 \Leftrightarrow \begin{cases} x^3 - y = 0 & \Leftrightarrow x = y^3, y = x^3 \\ y^3 - x = 0 \end{cases}$$

$$\Leftrightarrow x = x^9 \Leftrightarrow x = 0, y = 0 \text{ or } x = 1, 1 \text{ or } x = -1, y = -1.$$

These are the critical points.

$$f_{xx}(x,y) = 12x^2, f_{yy}(x,y) = 12y^2, f(x,y) = -4.$$

① $(0,0)$.

$D = 16 - 0 > 0$. This is a saddle point.

② $(1,1)$.

$$D = 16 - 12 \cdot 12 < 0. f_{xx}(1,1) = 12 > 0.$$

$\Rightarrow f(1,1) = 0$ is a local minima.

③ $(-1,-1)$.

$$D = 16 - 12 \cdot 12 < 0. f_{xx}(-1,-1) = 12 > 0.$$

$\Rightarrow f(-1,-1) = 0$ is a local minima.

Example 8.4.2. $K = \overline{B(0,0)}$.

Find the absolute maxima and minima of $f: K \rightarrow \mathbb{R}$.

$$f(x,y) = 2x^3 + y^4.$$

i.e. $\max f(K), \min f(K)$.

① Critical points of $f: B(0,0) \rightarrow \mathbb{R}$.

$$\nabla f(x,y) = (6x^2, 4y^3) = (0,0)$$
$$\Leftrightarrow (x,y) = (0,0). \text{ Note } f(0,0) = 0.$$

② $\mathcal{D}(k) = \{(x,y) : x^2 + y^2 = 1\}$.

For $(x,y) \in \mathcal{D}(k)$:

$$f(x,y) = 2x^3 + (1-x^2)^2 = \underbrace{x^4 + 2x^3 - 2x^2 + 1}_{g(x)}.$$

Consider $g(x)$ on $[-1, 1]$.

$$g'(x) = 4x^3 + 6x^2 - 4x$$
$$= 2x(2x-1)(x+2) = 0$$

$$\Leftrightarrow x = 0, \frac{1}{2}, -2 (\text{not in the domain}).$$

Note $g(0) = 1$, $g(\frac{1}{2}) = \frac{13}{16}$, $g(1) = 2$, $g(-1) = -2$.

So, absolute maxima $f(1,0) = 2$

absolute minima $f(-1,0) = -2$.

Week 8 Appendix

Lemma. Let $U \subseteq \mathbb{R}^n$ be open and let $f \in C^2(U)$. If $a \in U$ such that $D^2f(a)(h) > 0$ for all $0 \neq h \in \mathbb{R}^n$ then there exists $m > 0$ such that

$$D^2f(a)(x) \geq m\|x\|^2,$$

for all $x \in \mathbb{R}^n$.

Proof. Consider the compact set $K = \{x \in \mathbb{R}^n : \|x\| = 1\}$. Since $f \in C^2(U)$ we have that $D^2f(a)$ is continuous and positive on K . By the EVT, there exists $m > 0$ such that $m = \min\{D^2f(a)(x) : x \in K\}$. For $0 \neq x \in \mathbb{R}^n$ we then see that $\frac{x}{\|x\|} \in K$ and so

$$D^2f(a)\left(\frac{x}{\|x\|}\right) = \frac{1}{\|x\|^2}D^2f(a)(x) \geq m.$$

□

Lemma. Let $U \subseteq \mathbb{R}^n$ be open and let $f \in C^2(U)$. Suppose $a \in U$ such that $\nabla f(a) = 0$. Let $r > 0$ such that $B_r(a) \subseteq U$. There exists a function $\varepsilon : B_r(0) \rightarrow \mathbb{R}$ such that

$$\lim_{h \rightarrow 0} \varepsilon(h) = 0$$

and

$$f(a + h) - f(a) = \frac{1}{2}D^2f(a)(h) + \|h\|^2\varepsilon(h)$$

for $\|h\|$ sufficiently small.

Proof. Consider

$$\varepsilon(h) := \frac{f(a + h) - f(a) - \frac{1}{2}D^2f(a)(h)}{\|h\|^2}$$

for $0 \neq h \in B_r(0)$ and define $\varepsilon(0) = 0$. We are left to prove that $\varepsilon(h) \rightarrow 0$ as $h \rightarrow 0$. Let $h \in B_r(0)$. Since $\nabla(f)(a) = 0$ we have by Taylor's Theorem that

$$f(a + h) - f(a) = \frac{1}{2}D^2f(c)(h)$$

for some $c \in L(a, a + h)$. Then,

$$\begin{aligned} 0 &\leq |\varepsilon(h)|\|h\|^2 = \left| \frac{1}{2}D^2f(c)(h) - \frac{1}{2}D^2f(a)(h) \right| \\ &\leq \frac{1}{2} \sum_i \sum_j \left| \frac{\partial^2 f}{\partial x_i \partial x_j}(c) - \frac{\partial^2 f}{\partial x_i \partial x_j}(a) \right| |h_i h_j| \\ &\leq \frac{1}{2} \sum_i \sum_j \left| \frac{\partial^2 f}{\partial x_i \partial x_j}(c) - \frac{\partial^2 f}{\partial x_i \partial x_j}(a) \right| \|h\|^2 \end{aligned}$$

and

$$\frac{1}{2} \sum_i \sum_j \left(\frac{\partial^2 f}{\partial x_i \partial x_j}(c) - \frac{\partial^2 f}{\partial x_i \partial x_j}(a) \right) \rightarrow 0$$

as $h \rightarrow 0$ because $c \rightarrow a$ as $h \rightarrow 0$ and $f \in C^2(U)$. \square

Theorem. (Second Derivative Test) Let $U \subseteq \mathbb{R}^n$ be open and let $f \in C^2(U)$. Suppose $a \in U$ such that $\nabla f(a) = 0$.

1. If $D^2 f(a)(h) > 0$ for all $0 \neq h \in \mathbb{R}^n$ then $f(a)$ is a local minimum of f .
2. If $D^2 f(a)(h) < 0$ for all $0 \neq h \in \mathbb{R}^n$ then $f(a)$ is a local maximum of f .
3. If there exist $h, k \in \mathbb{R}^n$ such that $D^2 f(a)(h) > 0$ and $D^2 f(a)(k) < 0$ then a is a saddle point of f .

Proof. Let $r > 0$ such that $B_r(a) \subseteq U$. There exists a function $\varepsilon : B_r(0) \rightarrow \mathbb{R}$ such that

$$\lim_{h \rightarrow 0} \varepsilon(h) = 0$$

and

$$f(a + h) - f(a) = \frac{1}{2} D^2 f(a)(h) + \|h\|^2 \varepsilon(h)$$

for $\|h\|$ sufficiently small.

1. Suppose $D^2 f(a)(h) > 0$ for all $0 \neq h \in \mathbb{R}^n$. Let $m > 0$ such that

$$D^2 f(a)(x) \geq m\|x\|^2,$$

for all $x \in \mathbb{R}^n$. Then,

$$f(a + h) - f(a) = \frac{1}{2} D^2 f(a)(h) + \|h\|^2 \varepsilon(h) \geq \left(\frac{m}{2} + \varepsilon(h) \right) \|h\|^2 > 0$$

for all $\|h\|$ sufficiently small, since $m > 0$ and $\varepsilon(h) \rightarrow 0$ as $h \rightarrow 0$. Therefore $f(a + h) > f(a)$ for all $\|h\|$ sufficiently small, and so $f(a)$ is a local minimum of f .

2. Follows from (1) by replacing f with $-f$.

3. Let $h \in \mathbb{R}^n$. For small $t \in \mathbb{R}$,

$$\begin{aligned} f(a + th) - f(a) &= \frac{1}{2} D^2 f(a)(th) + \|th\|^2 \varepsilon(th) \\ &= t^2 \left(\frac{1}{2} D^2 f(a)(h) + \|h\|^2 \varepsilon(th) \right). \end{aligned}$$

Letting $t \rightarrow 0$, we have that $\varepsilon(th) \rightarrow 0$ and so $f(a + th) - f(a)$ takes on the same sign as $D^2 f(a)(h)$, which can be both positive and negative. Therefore a is a saddle point. \square

Week 9 Notes.

Module 1: Inverse Function Theorem.

Remark 9.1.1: Recall

$I = (a, b)$. If $f: I \rightarrow \mathbb{R}$ is continuous and injective, and $y \in f(I)$ is a point such that

- 1) f is differentiable at $x = f^{-1}(y) \in I$.
- 2) $f'(x) \neq 0$.

Then f^{-1} is differentiable at y and $(f^{-1})'(y) = \frac{1}{f'(x)}$.

Remark 9.1.2: Goal.

Develop a multivariable version of this theorem.

We need to generalize the idea of $\frac{1}{f'(x)} = (f'(x))^{-1}$ to something like $Df(x)^{-1}$ (matrix inverse).

Definition 9.1.3:

Let $U \subseteq \mathbb{R}^n$, $f: U \rightarrow \mathbb{R}^n$. We define the Jacobian of f at a by

$$Jf(a) := \det(Df(a)).$$

Theorem 9.1.4 [Inverse Function Theorem].

Let $U \subseteq \mathbb{R}^n$ be open, $f \in C^1(U, \mathbb{R}^n)$. If at a such that $Jf(a) \neq 0$, then there exists an open $a \in W \subseteq U$ such that

- 1) f is injective on W .
- 2) $f^{-1} \in C^1(f(W), \mathbb{R}^n)$.
- 3) For all $y \in f(W)$

$$D(f^{-1})(y) = [Df(x)]^{-1}$$

where $x = f^{-1}(y)$.

Proof: Appendix. This is quite technical.

Example 9.1.5. $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

$$f(x, y) = (x+y, \sin x + \cos y).$$

Note that

$$f_x(x, y) = (1, \cos x).$$

$$f_y(x, y) = (1, -\sin y).$$

So that $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$.

Question: Prove that f^{-1} exists and is differentiable on some open set containing $(0, 1)$ and compute $D(f^{-1})(0, 1)$.

Note: $f(x, y) = (0, 1)$. (Preimage).

$$\Leftrightarrow (x+y, \sin x + \cos y) = (0, 1)$$

$$\Leftrightarrow y = -x, \sin x + \cos(-x) = 1.$$

$$\Leftrightarrow y = -x, \sin x + \cos x = 1.$$

$$\Leftrightarrow (x, y) = (2k\pi, -2k\pi), k \in \mathbb{Z}.$$

$$\text{or } (x, y) = \left(\frac{\pi}{2} + 2k\pi, -\frac{\pi}{2} - 2k\pi\right), k \in \mathbb{Z}$$

Case 1: $a = (2k\pi, -2k\pi), k \in \mathbb{Z}$.

$$Jf(a) = \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} = -1 \neq 0.$$

Then by the inverse function theorem, we have the results we need to prove.

Moreover,

$$D(f^{-1})(0,1) = [Df(a)]^{-1} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}.$$

Case 2: $a = (\frac{\pi}{2} + 2k\pi, -\frac{\pi}{2} - 2k\pi)$.

$$Jf(a) = \begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} = 1 \neq 0.$$

Again, by the inverse function theorem, we have the results we need to prove. Moreover:

$$D(f^{-1})(0,1) = Df(a)^{-1} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}.$$

Remark 9.1.6:

The way we restrict f to make it injective depends on our choice of $f^{-1}(y)$.

Module 2: Implicit Function Theorem.

Remark 9.2.1: Idea.

When/where can $f(x, y, z) = 0$ be solved to express z as a function of x, y ?

That is when/where

$$\{ (x, y, z) \in \mathbb{R}^3 : f(x, y, z) = 0 \} = \{ (x, y, g(x, y)) : f(x, y, g(x, y)) = 0 \}$$

Example 9.2.2.

$$f(x, y, z) = x^2 + y^2 + z^2 - 1 = 0.$$

$U = \{ (x, y, z) \in \mathbb{R}^3 : z \geq 0 \}$ is open.

On U , we have

$$z = \sqrt{1 - x^2 - y^2} := g(x, y)$$

and $f(x, y, g(x, y)) = 0$.

Theorem 9.2.3 [Implicit Function Theorem].

Let $U \subseteq \mathbb{R}^{n+p}$ be open. $f = (f_1, \dots, f_n) \in C^1(U, \mathbb{R}^n)$

Let $x_0 \in \mathbb{R}^n, t_0 \in \mathbb{R}^p$ such that $f(x_0, t_0) = 0$.

If

$$\det \left[\frac{\partial f_i}{\partial x_j} (x_0, t_0) \right]_{n \times n} \neq 0 \xrightarrow{\substack{\text{ignore other} \\ \text{variables}}}$$

then there exists an open $t_0 \in V \subseteq \mathbb{R}^p$ and

a unique $g \in C^1(V, \mathbb{R}^n)$ such that.

① $g(t_0) = x_0$ and

② $f(g(t), t) = 0, \forall t \in V$.

Proof: Appendix.

Remark 9.2.4: Summary:

$t \in V \subseteq \mathbb{R}^P \rightarrow$ variables to keep.

$g(t) \in \mathbb{R}^n \rightarrow$ variables replaced by an implicit function of t .

Example 9.2.5: $xyz + \sin(x+y+z) = 0$.

Consider $f(x, y, z) = xyz + \sin(x+y+z)$

so that $f \in C^1(\mathbb{R}^3)$.

Note: $f(0, 0, 0) = 0$.

Now,

$$f_z(x, y, z) = xy + \cos(x+y+z)$$

$$\Rightarrow f_z(0, 0, 0) = 1 \neq 0.$$

$$\Rightarrow \det [1] = 1 \neq 0.$$

By the implicit function theorem, there exists open $V \subseteq \mathbb{R}^2$ with $(0, 0) \in V$ and $g(x, y)$ in $C^1(V)$ such that $g(0, 0) = 0$ and.

$$f(x, y, g(x, y)) = 0.$$

for all $(x, y) \in V$. i.e. $z = g(x, y)$ on V .

Example 9.2.6.

Prove there exists $U, V: \mathbb{R}^4 \rightarrow \mathbb{R}$.

and $(2, 1, -1, -2) \in U \subseteq \mathbb{R}^4$ open such that.

① $U, V \in C^1(U)$.

② $U(2, 1, -1, -2) = 4, V(2, 1, -1, -2) = 3$.

③ For all $(x, y, z, w) \in U$

$$U^2 + V^2 + W^2 = 29.$$

$$\frac{u^2}{x^2} + \frac{v^2}{y^2} + \frac{w^2}{z^2} = 17.$$

Solution:

$$f: \mathbb{R}^6 \rightarrow \mathbb{R}^2$$

$$f(u, v, x, y, z, w)$$

$$= (u^2 + v^2 + w^2 - 29, \frac{u^2}{x^2} + \frac{v^2}{y^2} + \frac{w^2}{z^2} - 17).$$

$$\Rightarrow f(4, 3, 2, 1, -1, -2) = 0.$$

and

$$\det \begin{bmatrix} \frac{\partial f_1}{\partial u} & \frac{\partial f_1}{\partial v} \\ \frac{\partial f_2}{\partial u} & \frac{\partial f_2}{\partial v} \end{bmatrix} = \begin{vmatrix} 2u & 2v \\ \frac{2u}{x^2} & \frac{2v}{y^2} \end{vmatrix} \\ = 4uv \left(\frac{1}{y^2} - \frac{1}{x^2} \right).$$

This is non-zero at $(4, 3, 2, 1, -1, -2)$.

By the Implicit function theorem, there exists open $U \ni (2, 1, -1, -2) \in \mathbb{R}^4$ and $g \in C^1(U, \mathbb{R}^2)$ such that $g(2, 1, -1, -2) = (4, 3)$ and $\forall (x, y, z, w) \in U$,

$$f(g(x, y, z, w), x, y, z, w) = 0.$$

Let $g = (g_1, g_2)$.

$$u(x, y, z, w) = g_1(x, y, z, w)$$

$$v(x, y, z, w) = g_2(x, y, z, w).$$

$$u, v \in C^1(U)$$

$$u(2, 1, -1, -2) = 4.$$

$$v(2, 1, -1, -2) = 3.$$

$$f(g(x, y, z, w), x, y, z, w) = 0.$$

$$\Rightarrow f(u(x, y, z, w), v(x, y, z, w), x, y, z, w) = 0.$$

$$\Rightarrow w^2 + v^2 + w^2 = 29.$$

$$\frac{w^2}{x^2} + \frac{v^2}{y^2} + \frac{w^2}{z^2} = 17.$$

Module 3: Lagrange Multipliers.

Remark 9.3.1: Goal.

Solve optimization problems with constraints.

Definition 9.3.2:

Let $U \subseteq \mathbb{R}^n$ be open, $f: U \rightarrow \mathbb{R}$.

① Let $a \in U$. We call $f(a)$ a local max of f .

Subject to the constraints

$g_i: U \rightarrow \mathbb{R}$.

$1 \leq i \leq m$.

If $g_i(a) = 0$ for all i and there exists $r > 0$ such that whenever $x \in B_r(a)$ and $g_i(x) = 0$ for all i , then $f(x) \leq f(a)$.

② Similar definition for local min.

Theorem 9.3.3.

Let $U \subseteq \mathbb{R}^n$ be open, $m < n$, $f, g_1, g_2, \dots, g_m \in C^1(U)$.

Suppose $a \in U$ such that

$$\det \left[\frac{\partial g_i}{\partial x_j}(a) \right]_{m \times m} \neq 0$$

If $f(a)$ is a local extremum of f subject to the constraints g_i , then $\exists x_1, x_2, \dots, x_m \in \mathbb{R}$ such that

$$\nabla f(a) + \sum_{i=1}^m x_i \nabla g_i(a) = 0.$$

Remark 9.3.4.

The proof for the above theorem is quite technical. We only consider the case $m=2, n=3$ here.

We want to show that there exists $\lambda, \mu \in \mathbb{R}$ such that

$$\lambda \frac{\partial g_1}{\partial x_j}(a) + \mu \frac{\partial g_2}{\partial x_j}(a) = -\frac{\partial f}{\partial x_j}(a)$$

for $j = 1, 2, 3$.

Let $A = \begin{bmatrix} \frac{\partial g_1}{\partial x_1}(a) & \frac{\partial g_1}{\partial x_2}(a) \\ \frac{\partial g_2}{\partial x_1}(a) & \frac{\partial g_2}{\partial x_2}(a) \end{bmatrix}$.

so that $\det A \neq 0$.

In particular

$$[\lambda, \mu] A = \left[-\frac{\partial f}{\partial x_1}(a), -\frac{\partial f}{\partial x_2}(a) \right]^{\text{partial}}$$

has a unique solution $[\lambda, \mu]$. Expand this we get the case for x_1, x_2 . We are left to show that


$$\lambda \frac{\partial g_1}{\partial x_3}(a) + \mu \frac{\partial g_2}{\partial x_3}(a) = -\frac{\partial f}{\partial x_3}(a).$$

The idea is then to use the implicit function theorem to replace x_3 with $h(x_1, x_2)$ and prove the above result with what we have with x_1, x_2 .

Example 9-3-5.

Maximize and minimize $f(x, y, z) = x + 2y$,
subject to the constraints

$$\begin{aligned} \textcircled{1} \quad & x + y + z = 1 \\ \textcircled{2} \quad & y^2 + z^2 = 4. \end{aligned}$$



Geometrically, such a max/min must exist!

Let. $f(x, y, z) = x + 2y$, $g(x, y, z) = x + y + z - 1$
 $h(x, y, z) = y^2 + z^2 - 4$.

Note:

$$\begin{vmatrix} g_x & g_y \\ h_x & h_y \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ 0 & 2y \end{vmatrix} = 2y \neq 0 \text{ for } y \neq 0.$$

Moreover, if $g(x, 0, z) = h(x, 0, z) = 0$
then $z = \pm\sqrt{2}$, $x = 1 \mp \sqrt{2}$. In which case.

$$f(1 \mp \sqrt{2}, 0, \pm\sqrt{2}) = 1 \mp \sqrt{2}.$$

Otherwise, such a max/min is of the form

$$\nabla f = \lambda \nabla g + \mu \nabla h.$$

$$\Rightarrow [1, 2, 0] = \lambda [1, 1, 1] + \mu [0, 2y, 2z].$$

$$\Rightarrow \lambda = 1, \lambda + 2\mu y = 2, \lambda + 2\mu z = 0.$$

$$\Rightarrow y = \frac{1}{2\mu}, z = \frac{-1}{2\mu}.$$

$$\text{But } y^2 + z^2 = 4 \Rightarrow \frac{3}{4M^2} = 4.$$

$$\Rightarrow M = \pm \frac{1}{2\sqrt{2}}.$$

$$\Rightarrow y = \pm \sqrt{2}, z = \mp \sqrt{2}.$$

$$\Rightarrow x = 1.$$

Checking:

$$f(1, \sqrt{2}, -\sqrt{2}) = 1 + 2\sqrt{2} : \text{max}$$

$$f(1, -\sqrt{2}, \sqrt{2}) = 1 - 2\sqrt{2} : \text{min}.$$

Week 9 Appendix

Lemma 1. Let $U \subseteq \mathbb{R}^n$ be open. Suppose $a \in U$ so that we may find $r > 0$ such that $\overline{B_r(a)} \subseteq U$. Let $f : U \rightarrow \mathbb{R}^n$ be continuous and injective when restricted to $\overline{B_r(a)}$ and assume its first order partials exist on $B_r(a)$. If $Jf \neq 0$ on $B_r(a)$ then there exists $\varepsilon > 0$ such that $B_\varepsilon(f(a)) \subseteq f(B_r(a))$.

Proof. Consider $g : \overline{B_r(a)} \rightarrow \mathbb{R}$ given by $g(x) = \|f(x) - f(a)\|$. Since f is continuous and injective on $\overline{B_r(a)}$ we have that g is continuous and $g(x) > 0$ for all $x \neq a$. By the EVT,

$$m = \inf\{g(x) : \|x - a\| = r\} > 0.$$

Take $\varepsilon = m/2$. We claim that $B_\varepsilon(f(a)) \subseteq f(B_r(a))$.

Let $y \in B_\varepsilon(f(a))$. Again by the EVT, there exists $b \in \overline{B_r(a)}$ such that

$$\|f(b) - y\| = \inf\{\|f(x) - y\| : x \in \overline{B_r(a)}\}.$$

For the sake of contradiction, suppose that $\|b - a\| = r$. Then,

$$\varepsilon > \|f(a) - y\| \geq \|f(b) - y\| \geq \|f(b) - f(a)\| - \|f(a) - y\| = g(b) - \|f(a) - y\| \geq m - \varepsilon = 2\varepsilon - \varepsilon = \varepsilon,$$

which is a contradiction. Therefore we have that $b \in B_r(a)$.

If we can show that $y = f(b)$ we are done. This is where the information about the partial derivatives and the Jacobian come into play. Consider the continuous function $h : \overline{B_r(a)} \rightarrow \mathbb{R}$ given by $h(x) = \|f(x) - y\|$. By construction, $h(b)$ is the minimum value of h . Moreover, $h^2(b)$ is also the minimum value of h^2 . Since $b \in B_r(a)$, which is open, we have that $\nabla h^2(b) = 0$ (note that in last week's proof we really just needed the first order partials to exist at a , not necessarily differentiability at a). However,

$$h^2(x) = \sum_{i=1}^n (f_i(x) - y_i)^2,$$

and so for every $1 \leq j \leq n$,

$$0 = \frac{\partial h^2}{\partial x_j}(b) = \sum_{i=1}^n 2(f_i(b) - y_i) \frac{\partial f_i}{\partial x_j}(b).$$

Thus, $Df(b)x = 0$, where $x = (2(f_1(b) - y_1), 2(f_2(b) - y_2), \dots, 2(f_n(b) - y_n))^T$. Since $Df(b)$ is invertible ($Jf(b) \neq 0$) we have that $x = 0$. Hence $f(b) = y$, as required. \square

Lemma 2. Let $U \subseteq \mathbb{R}^n$ be open and nonempty. If $f : U \rightarrow \mathbb{R}^n$ is continuous, injective, has all first-order partials existing on U , AND is such that $Jf \neq 0$ on U , then f^{-1} is continuous on $f(U)$.

Proof. To prove that $f^{-1} : f(U) \rightarrow \mathbb{R}^n$ is continuous it suffices to prove that $f(W)$ is open whenever W is open in \mathbb{R}^n and $W \subseteq U$ (Why? Piazza!). Well, let W be such a set and take $b \in f(W)$ so that $b = f(a)$ for some $a \in W$. Since W is open there exists $r > 0$ such that $\overline{B_r(a)} \subseteq W$. By the previous lemma, there then exists $\varepsilon > 0$ such that

$$B_\varepsilon(b) \subseteq f(B_r(a)).$$

Thus, $B_\varepsilon(b) \subseteq f(W)$, and so $f(W)$ is open. \square

Lemma 3. Let $U \subseteq \mathbb{R}^n$ be open and let $f \in C^1(U, \mathbb{R}^n)$. If $a \in U$ such that $Jf(a) \neq 0$ then there exists $r > 0$ such that $B_r(a) \subseteq U$, f is injective on $B_r(a)$, $Jf \neq 0$ on $B_r(a)$, and

$$\det \left(\frac{\partial f_i}{\partial x_j}(c_i) \right) \neq 0$$

for all $c_1, c_2, \dots, c_n \in B_r(a)$.

Proof. Let $W = U \times U \times \dots \times U$ (n-times). Consider $h : W \rightarrow \mathbb{R}$ defined by

$$h(x_1, x_2, \dots, x_n) = \det \left(\frac{\partial f_i}{\partial x_j}(x_i) \right)$$

Since $f \in C^1(U, \mathbb{R}^n)$ and a determinant is a polynomial in its entries, we have that h is continuous. Note that $h(a, a, \dots, a) = Jf(a) \neq 0$. Thus we may find an open interval $h(a, a, \dots, a) \in I \subseteq \mathbb{R}$ such that $0 \notin I$. Then, $h^{-1}(I)$ is open (note that W is open) and so there exists $R > 0$ such that $B_R(a, a, \dots, a) \subseteq h^{-1}(I)$. But then we may find $r > 0$ such that

$$B_r(a) \times \dots \times B_r(a) \subseteq B_R(a, a, \dots, a) \subseteq h^{-1}(I).$$

We then see that $Jf \neq 0$ on $B_r(a)$, and

$$\det \left(\frac{\partial f_i}{\partial x_j}(c_i) \right) \neq 0$$

for all $c_1, c_2, \dots, c_n \in B_r(a)$.

We are left to show that f injective on $B_r(a)$. For the sake of contradiction suppose there exists $x \neq y$ in $B_r(a)$ such that $f(x) = f(y)$. Since f is differentiable on $B_r(a)$, every f_i is differentiable on $B_r(a)$. Fix $1 \leq i \leq n$. By the MVT there exists $c_i \in L(x, y)$ such

that $0 = f_i(x) - f_i(y) = Df_i(c_i)(x - y)$. Letting $A = \left[\frac{\partial f_i}{\partial x_j}(c_i) \right]$ we see that $A(x - y) = 0$. Since $x - y \neq 0$, A is not invertible and so

$$\det \left(\frac{\partial f_i}{\partial x_j}(c_i) \right) = 0,$$

a contradiction. \square

Recall. (Cramer's Rule) Let A be a $n \times n$ invertible matrix and consider a system of equations $Ax = b$. This system has a unique solution $x = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$ given by

$$x_i = \frac{\det(A(i))}{\det A},$$

where $A(i)$ is the matrix obtained from A by replacing its i^{th} column by b .

Theorem. (Inverse Function Theorem) Let $U \subseteq \mathbb{R}^n$ be open and let $f \in C^1(U, \mathbb{R}^n)$. If $a \in U$ such that $Jf(a) \neq 0$ then there exists an open set $W \subseteq U$ such that

1. f is injective on W
2. $f^{-1} \in C^1(f(W), \mathbb{R}^n)$
3. For every $y \in f(W)$, if $x = f^{-1}(y)$ then

$$Df^{-1}(y) = [Df(x)]^{-1}.$$

Proof. Since this is a rather long and technical proof, we break it into digestible, enumerated pieces.

1. By Lemma 3 there exists $r > 0$ with $W := B_r(a) \subseteq U$ such that f is injective on W , $Jf \neq 0$ on W , and

$$\det \left(\frac{\partial f_i}{\partial x_j}(c_i) \right) \neq 0$$

for all $c_1, c_2, \dots, c_n \in W$. Moreover, by Lemma 2, f^{-1} is continuous on $f(W)$.

2. We claim that $f^{-1} \in C^1(f(W), \mathbb{R}^n)$. Fix $y_0 \in f(W)$ and $1 \leq i, j \leq n$. Choose $0 \neq t \in \mathbb{R}$ sufficiently small so that $y_0 + te_j \in f(W)$. We may then find $x_0, x_1 = x_1(t) \in W$ such that $f(x_0) = y_0$ and $f(x_1) = y_0 + te_j$. By the MVT, for every $1 \leq i \leq n$ there exists $c_i = c_i(t) \in L(x_0, x_1)$ such that

$$\nabla f_i(c_i)(x_1 - x_0) = f_i(x_1) - f_i(x_0) = \begin{cases} t & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}.$$

Therefore,

$$\nabla f_i(c_i)\left(\frac{x_1 - x_0}{t}\right) = \frac{1}{t}(f_i(x_1) - f_i(x_0)) = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}.$$

Now let A_j be the $n \times n$ matrix whose i^{th} row is $\nabla f_i(c_i)$. By assumption, $\det(A_j) \neq 0$. Moreover, $A_j\left(\frac{x_1 - x_0}{t}\right) = e_j$. For $1 \leq k \leq n$, we then see that

$$\frac{(f^{-1})_k(y_0 + te_j) - (f^{-1})_k(y_0)}{t} = \frac{x_{1,k} - x_{0,k}}{t},$$

where by Cramer's Rule, $Q_k(t) := \frac{x_{1,k} - x_{0,k}}{t}$ is a quotient of determinants of matrices whose entries are either 0, 1, or a first-order partial of f evaluated at a c_ℓ . As $t \rightarrow 0$ we clearly have that $y_0 + te_j \rightarrow y_0$. But then, by the continuity of f^{-1} , we have that $x_1 \rightarrow x_0$ and so $c_i \rightarrow x_0$. Since f is C^1 , we therefore that that $Q_k(t) \rightarrow Q_k$, where Q_k is a quotient of determinants of matrices whose entries are either 0, 1, or a first-order partial of f evaluated at a $x_0 = f^{-1}(y_0)$. Since $f \in C^1$ and f^{-1} is continuous at y_0 , it follows that Q_k is continuous at each $y_0 \in f(W)$. Moreover,

$$\lim_{t \rightarrow 0} \frac{(f^{-1})_k(y_0 + te_j) - (f^{-1})_k(y_0)}{t} = \lim_{t \rightarrow 0} \frac{x_{1,k} - x_{0,k}}{t} = Q_k.$$

Hence all of the partial derivatives of f^{-1} exist and are continuous at y_0 (ie. $f^{-1} \in C^1(f(W), \mathbb{R}^n)$).

3. Finally, we quickly run the chain rule and note that for $y \in f(W)$,

$$I = DI(y) = D(f \circ f^{-1})(y) = Df(f^{-1}(y))D(f^{-1})(y).$$

The result follows. □

Theorem. (Implicit Function Theorem) Suppose $U \subseteq \mathbb{R}^{n+p}$ is open and $f = (f_1, f_2, \dots, f_n) \in C^1(U, \mathbb{R}^n)$. Suppose $x_0 \in \mathbb{R}^n$ and $t_0 \in \mathbb{R}^p$ such that $f(x_0, t_0) = 0$. If

$$\det\left(\frac{\partial f_i}{\partial x_j}(x_0, t_0)\right)_{n \times n} \neq 0,$$

then there is an open set $t_0 \in V \subseteq \mathbb{R}^p$ and a unique function $g \in C^1(V, \mathbb{R}^n)$ such that $g(t_0) = x_0$ and $f(g(t), t) = 0$ for all $t \in V$.

Proof. For every $(x, t) \in U$ let

$$F(x, t) := (f(x, t), t) = (f_1(x, t), \dots, f_n(x, t), t_1, t_2, \dots, t_p).$$

Notice that $F(x_0, t_0) = (0, t_0)$. Then, F is a function from U to \mathbb{R}^{n+p} with

$$DF = \begin{bmatrix} \left(\frac{\partial f_i}{\partial x_j} \right)_{n \times n} & B \\ 0_{p \times n} & I_{p \times p} \end{bmatrix},$$

where $0_{p \times n}$ is the $p \times n$ zero matrix, $I_{p \times p}$ is the $p \times p$ identity matrix, and B is a matrix whose entries are first-order partials of the f_i 's with respect to the t_j 's. Taking the determinant of this crazy matrix evaluated at (x_0, t_0) , we have that

$$JF(x_0, t_0) = \det \left(\frac{\partial f_i}{\partial x_j}(x_0, t_0) \right)_{n \times n} \cdot \det I_{p \times p} \neq 0.$$

Therefore, by the Inverse Function Theorem there exists an open set $(x_0, t_0) \in W \subseteq U$ such that F is injective on W and $F^{-1} \in C^1(F(W), \mathbb{R}^{n+p})$.

To ease notation, let $G = F^{-1} = (G_1, G_2, \dots, G_n, G_{n+1}, \dots, G_{n+p})$. Consider $\varphi : F(W) \rightarrow \mathbb{R}^n$ given by

$$\varphi = (G_1, G_2, \dots, G_n).$$

By construction we have that

$$\varphi(F(x, t)) = x$$

for all $(x, t) \in W$ and

$$F(\varphi(x, t), t) = (x, t),$$

for all $(x, t) \in F(W)$.

Consider $V = \{t \in \mathbb{R}^p : (0, t) \in F(W)\}$ and the function $g : V \rightarrow \mathbb{R}^n$ given by $g(t) = \varphi(0, t)$. Since G is C^1 , it follows that φ is also C^1 . Hence, $g \in C^1(V, \mathbb{R}^n)$. Also note that V is open since $F(W)$ is open. Finally, we compute that

$$g(t_0) = \varphi(0, t_0) = \varphi(F(x_0, t_0)) = x_0,$$

and note that for all $(x, t) \in F(W)$,

$$f(\varphi(x, t), t) = x.$$

In particular,

$$0 = f(\varphi(0, t), t) = f(g(t), t) = 0$$

for all $t \in V$.

Uniqueness follows from the injectivity of F . (Please share the details on Piazza!)

□

Week 10 Notes.

Module 1: Riemann Integration

Remark 10.1.1: Recall $f: [a, b] \rightarrow \mathbb{R}$ bounded.

① A partition of $[a, b]$ is a set $P = \{x_0, x_1, \dots, x_n\}$ where $a = x_0 < x_1 < \dots < x_n = b$.

② For a partition $P = \{x_0, \dots, x_n\}$:

$$U(f, P) = \sum_{i=1}^n M_i (x_i - x_{i-1}).$$

$$L(f, P) = \sum_{i=1}^n m_i (x_i - x_{i-1}).$$

where

$$m_i = \inf \{f(x) : x \in [x_{i-1}, x_i]\}.$$

$$M_i = \sup \{f(x) : x \in [x_{i-1}, x_i]\}.$$

③ We say f is integrable if

$$\inf_P \{U(f, P)\} = \sup_P \{L(f, P)\}.$$

$$\begin{array}{ccc} \text{Upper Riemann} & \diagdown & \text{Lower Riemann Integral.} \\ \text{Integral} & \int_a^b f(x) dx & \int_a^b f(x) dx \end{array}$$

Definition 10.1.2:

A rectangle in \mathbb{R}^n is a set of the form

$$R = [a_1, b_1] \times [a_2, b_2] \times \dots \times [a_n, b_n].$$

where $a_i \leq b_i$.


Definition 10.1.3:

A partition of $R = [a_1, b_1] \times \cdots \times [a_n, b_n]$ is a grid of rectangles on R obtained by partitioning each $[a_i, b_i]$.

Example 10.1.4: In \mathbb{R}^2 :

R	
R_1	R_2
R_3	R_4
R_5	R_6

This is a partition



This is not a partition.

Definition 10.1.5

Let $R = [a_1, b_1] \times \cdots \times [a_n, b_n] \subseteq \mathbb{R}^n$ be a rectangle.

The volume of R is:

$$V(R) = (b_1 - a_1)(b_2 - a_2) \cdots (b_n - a_n).$$

Definition 10.1.6:

Let $R \subseteq \mathbb{R}^n$ be a rectangle and let $f: R \rightarrow \mathbb{R}$ be bounded. Let $P = \{R_1, \dots, R_n\}$ be a partition of R .

① Upper sum of f relative to P is:

$$U(f, P) = \sum_{i=1}^n M_i V(R_i).$$

where

$$M_i = \sup \{f(x) : x \in R_i\}.$$

② Lower sum of f relative to P is:

$$L(f, P) = \sum_{i=1}^n m_i V(R_i).$$


where

$$m_i = \inf \{f(x) : x \in R_i\}.$$

Definition 10.1.7:

Let P, Q be partitions of $R \subseteq \mathbb{R}^n$. We say P is a refinement of Q , written $P \leq Q$, if P is obtained from Q by partitioning the sides of R even further.

Example 10.1.8: $P \leq Q$:



Proposition 10.1.9:

If $P \leq Q$ on R and $f: R \rightarrow \mathbb{R}$ is bounded, then

$$U(f, P) \leq U(f, Q)$$

and

$$L(f, P) \geq L(f, Q).$$

Proof sketch:

Q. P

M	M_1	M_2
m	m_1	m_2
	M_3	M_4
	m_3	m_4

$$M_V(R) = M_V(R_1) + \dots + M_V(R_4)$$

$$\geq M_1 V(R_1) + \dots + M_4 V(R_4)$$

$$m_V(R) \leq m_1 V(R_1) + \dots + m_4 V(R_4).$$

Definition 10.1.10

Let $f: R \rightarrow \mathbb{R}$ be bounded, where $R \subseteq \mathbb{R}^n$ be a rectangle.

① The lower integral of f is:

$$\underline{\int}_R f = \sup \{ L(f, P) : P \}$$

② The Upper integral of f is:

$$\overline{\int}_R f = \inf \{ U(f, P) : P \}$$

③ We say f is (Riemann) integrable over R if.

$$\int_R f := \underline{\int}_R f = \overline{\int}_R f$$

Example 10.1.11: $f: [0, 1] \times [0, 1] \rightarrow \mathbb{R}$.

$$f(x, y) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & x \notin \mathbb{Q} \end{cases}$$

for all P , we have $U(f, P) = 1$, $L(f, P) = 0$

so, $\underline{\int}_R f = 0 \neq 1 = \overline{\int}_R f$, f is not integrable.

Module 2: Characterization Theorem.

Remark 10.2.1: Goal

Give an alternative definition of $f: R \rightarrow \mathbb{R}$ being integrable.

Lemma 10.2.2:

Let $f: R \rightarrow \mathbb{R}$ be bounded. If P, Q are partitions of R , then $L(f, P) \leq U(f, Q)$.

Proof sketch:

Find a common refinement S (eg overlap P and Q)

Then,

$$L(f, P) \leq L(f, S) \leq U(f, S) \leq U(f, Q).$$

Remark 10.2.3:

Let $f: R \rightarrow \mathbb{R}$ be bounded. For all Partitions P, Q

$$L(f, P) \leq U(f, Q).$$

$$\Rightarrow L(f, P) \leq \bar{S}_R f$$

$$\Rightarrow \underline{S}_R f \leq \bar{S}_R f.$$

Theorem 10.2.4:

Let $f: R \rightarrow \mathbb{R}$ be bounded. f is integrable iff for all $\varepsilon > 0$, there exists P such that.

$$U(f, P) - L(f, P) < \varepsilon$$

Proof

\Rightarrow : Assume f is integrable. So:

$$\underline{S}_R f = \bar{S}_R f.$$

Let $\varepsilon > 0$ be given. We may find partitions P, Q ,

such that.

$$\underline{\int}_R f - \frac{\epsilon}{2} < L(f, P)$$

$$U(f, Q) < \overline{\int}_R f + \frac{\epsilon}{2}$$

\Rightarrow

$$U(f, Q) < L(f, P) + \epsilon$$

Let S be a common refinement of P, Q . ($S \leq P, Q$).

$$U(f, S) \leq U(f, Q) < L(f, P) + \epsilon \leq L(f, S) + \epsilon$$

$$\Rightarrow U(f, S) - L(f, S) < \epsilon$$

\Leftarrow : Assume there exists partition P such that

$$U(f, P) - L(f, P) < \epsilon$$

for given $\epsilon > 0$.

So:

$$0 \leq \overline{\int}_R f - \underline{\int}_R f \leq U(f, P) - L(f, P) < \epsilon$$

$$\Rightarrow \overline{\int}_R f = \underline{\int}_R f.$$

□

Module 3: Jordan Content and Lebesgue Measure.

Remark 10.3.1: Goal

- ① Give a much stronger characterization of integrability
- ② To integrate over non-rectangles.

Definition 10.3.2:

- ① A set $A \subseteq \mathbb{R}^n$ has (Lebesgue) measure zero if for all $\epsilon > 0$, there exists rectangles R_i ($i \in \mathbb{N}$) such that

$$A \subseteq \bigcup_{i=1}^{\infty} R_i$$

and

$$\sum_{i=1}^{\infty} V(R_i) < \epsilon.$$

- ② A set $A \subseteq \mathbb{R}^n$ has (Jordan) content zero if for all $\epsilon > 0$, there exists rectangles R_1, R_2, \dots, R_m such that

$$A \subseteq \bigcup_{i=1}^m R_i$$

and:

$$\sum_{i=1}^m V(R_i) < \epsilon.$$

Proposition 10.3.3:

If $A \subseteq \mathbb{R}^n$ has content zero, then it has measure zero.

Proof:

Let $\epsilon > 0$ be given. Suppose A has content zero.

There exists rectangles R_1, \dots, R_m such that

$$A \subseteq R_1 \cup \dots \cup R_m$$

and

$$\sum_{i=1}^m V(R_i) < \varepsilon.$$

For $i \geq m$, let $R_i \subseteq \mathbb{R}^n$ be any rectangle with volume zero (e.g. take one side length to be $[a, a] = \{a\}$). So:

$$A \subseteq \bigcup_{i=1}^{\infty} R_i$$

and

$$\sum_{i=1}^{\infty} V(R_i) = \sum_{i=1}^m V(R_i) < \varepsilon.$$

□

Example 10.3.4 (measure zero but not content zero).

$$A = \mathbb{Q} \subseteq \mathbb{R}^1.$$

① Claim: \mathbb{Q} has measure zero.

$$\mathbb{Q} = \{q_1, q_2, q_3, \dots\}. \text{ (Since } \mathbb{Q} \text{ is countable).}$$

Let $\varepsilon > 0$ be given.

$$R_i = [q_i - \frac{\varepsilon}{2^{i+2}}, q_i + \frac{\varepsilon}{2^{i+2}}].$$

$$\text{Clearly } \mathbb{Q} \subseteq \bigcup_{i=1}^{\infty} R_i. \sum_{i=1}^{\infty} V(R_i) = \sum_{i=1}^{\infty} \frac{\varepsilon}{2^{i+1}} = \frac{\varepsilon}{2} < \varepsilon.$$

This shows \mathbb{Q} has measure zero.

② \mathbb{Q} does not have Jordan content zero.

Take $\varepsilon = 1$. To cover \mathbb{Q} by finitely many rectangles, \mathbb{Q} has to be bounded.

Note: this proof can be used to show any countable set has Lebesgue measure zero.

Proposition 10.3.5:

If $A_1, A_2, \dots, \subseteq \mathbb{R}^n$ have measure zero, then

$A = \bigcup_{i=1}^{\infty} A_i$ has measure zero.

Proof Sketch:

Let $\epsilon > 0$ be given. $A_i \subseteq \bigcup_{j=1}^{\infty} R_{i,j}$, $\sum_{j=1}^{\infty} V(R_{i,j}) < \frac{\epsilon}{2^i}$

$$A \subseteq \bigcup_{i,j} R_{i,j}. \sum_{i,j} V(R_{i,j}) = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} V(R_{i,j}) < \sum_{i=1}^{\infty} \frac{\epsilon}{2^i} = \epsilon.$$

Proposition 10.3.6:

If $A \subseteq \mathbb{R}^n$ is compact and has measure zero, then A has content zero.

Proof:

Let $\epsilon > 0$ be given. By assignment, there exists open rectangles R_i such that

$$A \subseteq \bigcup_{i=1}^{\infty} R_i$$

and

$$\sum_{i=1}^{\infty} V(R_i) < \epsilon$$

By compactness, $A \subseteq \bigcup_{i=1}^m R_i$ for some m .

Moreover, $\sum_{i=1}^m V(R_i) \leq \sum_{i=1}^{\infty} V(R_i) < \epsilon$.

Module 4: Integrability and Measure.

Remark 10.4.1: Goal

Describe the relationship between integrability and measure zero.

Definition 10.4.2:

Let $A \subseteq \mathbb{R}^n$, $f: A \rightarrow \mathbb{R}$ be bounded. For $a \in A$ and $\delta > 0$, we define:

$$M(a, f, \delta) = \sup \{f(x) : x \in A, \|x-a\| < \delta\}.$$

$$m(a, f, \delta) = \inf \{f(x) : x \in A, \|x-a\| < \delta\}.$$

$$\Theta(f, a) = \lim_{\delta \rightarrow 0} (M(a, f, \delta) - m(a, f, \delta)).$$

$\Theta(f, a)$ is called the oscillation of f at a .

Remark 10.4.3 [Piazza]

① $\Theta(f, a)$ always exists

② f is continuous at a iff $\Theta(f, a) = 0$.

Proposition 10.4.4.

Let $A \subseteq \mathbb{R}^n$ be closed. If $f: A \rightarrow \mathbb{R}$ is bounded, then $\forall \varepsilon > 0, \exists x \in A : \Theta(f, x) \geq \varepsilon$ is closed.

Proof:

Let $B = \{x \in A : \Theta(f, x) \geq \varepsilon\}$. Take $x \in A \setminus B$.

So, $\Theta(f, x) < \varepsilon$ and so there exists $\delta > 0$ with

$$M(x, f, \delta) - m(x, f, \delta) < \varepsilon$$

Consider $y \in B_{\delta/2}(x) \cap A$. Then, for $z \in A$ with $|y-z| < \frac{\delta}{2}$,

$$|z-x| \leq |z-y| + |y-x| < \delta$$

and so

$$\begin{aligned} m(x, f, \delta) &\leq f(z) \leq M(x, f, \delta) \\ \Rightarrow M(y, f, \delta/2) - m(y, f, \delta/2) &< \varepsilon \\ \Rightarrow \theta(f, y) &< \varepsilon \\ \Rightarrow B_{\delta/2}(x) \cap A &\subseteq A \setminus B \\ \Rightarrow A \setminus B &\text{ is relatively open in } A \\ \Rightarrow B &\text{ is relatively closed in } A \\ \Rightarrow B &\text{ is closed since } A \text{ is closed.} \end{aligned}$$

□

Proposition 10.4.5.

Let $R \subseteq \mathbb{R}^n$ be rectangle, $f: R \rightarrow \mathbb{R}$ be bounded.

Let $\varepsilon > 0$. If $\theta(f, x) < \varepsilon$ for all $x \in R$, then $\exists P$ such that $U(f, P) - L(f, P) < \varepsilon \cdot V(R)$.

Proof.

For all $x \in R$, we may find $\delta_x > 0$ such that

$$M(x, f, \delta_x) - m(x, f, \delta_x) < \varepsilon.$$

For all $x \in R$, let R_x be an open rectangle such that

$$x \in R_x \subseteq B_{\delta_x}(x).$$

Let $U_x = R_x \cap R$. We see that $R = \bigcup_{x \in R} U_x$ is a relatively open cover of R . Since R is compact,

there exists $x_1, \dots, x_m \in R$ such that $R = U_{x_1} \cup \dots \cup U_{x_m}$.
 Let P be a partition of R so fine that each
 subrectangle of P is contained in some $\overline{U_{x_i}}$. Note

$$\begin{aligned}\overline{U_{x_i}} &= \overline{R_{x_i} \cap R} \\ &\subseteq \overline{B_{\delta_{x_i}}(x_i)} \cap R \\ &\subseteq B_{\delta_{x_i}}(x_i) \cap R.\end{aligned}$$

So, for every $R_i \in P$

$$\begin{aligned}M_i - m_i &\leq \varepsilon \\ \Rightarrow U(f, P) - L(f, P) &= \sum_{R_i \in P} (M_i - m_i) V(R_i) \\ &< \sum_{R_i \in P} \varepsilon V(R_i) = \varepsilon V(R) \quad \square.\end{aligned}$$

Theorem 10.4.6.

Let $R \subseteq \mathbb{R}^n$ be rectangle, $f: R \rightarrow \mathbb{R}$ be bounded. Let $A = \{x \in R : f \text{ is not continuous at } x\}$. Then f is integrable iff A has measure zero.

Proof: Appendix.

Week 10 Appendix

In the following, $R \subseteq \mathbb{R}^n$ will denote a rectangle and $f : R \rightarrow \mathbb{R}$ will be a bounded function. Also, we let

$$A = \{x \in R : f \text{ is not continuous at } x\}.$$

Theorem. If A has measure zero then f is integrable.

Proof. Suppose A has measure zero and let $\varepsilon > 0$ be given. Let $B = \{x \in R : o(f, x) \geq \varepsilon\}$, so that B is compact (R is bounded) from our first proposition. Since $B \subseteq A$ (remember that the oscillation function is zero at points of continuity), we have that B also has measure zero. Since B is compact, B also has content zero. In particular, we may find finitely many rectangles U_1, \dots, U_m whose interiors cover B (I am using an assignment problem here), such that $\sum v(U_i) < \varepsilon$.

Let X denote the set of subrectangles of R which are contained in at least one U_i . Let Y denote the set of subrectangles of R which are contained in $R \setminus B$. Now, since the U_i 's cover B , we may find a partition $P = \{R_1, \dots, R_k\}$ fine enough so that the elements of P are from either X or Y .

Since f is bounded, there exists M such that $|f(x)| \leq M$ for all $x \in R$. In particular, for every $R_i \in P$, $M_i - m_i \leq 2M$ (as in the definition of upper and lower sum). By the definition of X ,

$$\sum_{R_i \in X} (M_i - m_i)v(R_i) \leq 2M \sum_{R_i \in X} v(R_i) \leq 2M \sum_{i=1}^m v(U_i) < 2M\varepsilon.$$

Now, if $R_i \in Y$ and $x \in R_i$, we have that $o(f, x) < \varepsilon$. By our second technical proposition (from class), we may find a partition $P_i = \{S_{i_1}, \dots, S_{i_{\alpha(i)}}\}$ of R_i such that

$$\sum_{j=1}^{\alpha(i)} (M_j - m_j)v(S_{i_j}) < \varepsilon v(R_i).$$

By replacing each $R_i \in Y$ with $S_{i_1}, S_{i_2}, \dots, S_{i_{\alpha(i)}}$ (and leaving the $R_i \in X$ alone), this creates a refinement $Q \leq P$. Finally,

$$\begin{aligned} U(f, Q) - L(f, Q) &= \sum_{R_i \in X} (M_i - m_i)v(R_i) + \sum_{R_i \in Y} \sum_{j=1}^{\alpha(i)} (M_j - m_j)v(S_{i_j}) \\ &< 2M\varepsilon + \sum_{R_i \in Y} \varepsilon v(R_i) \\ &\leq 2M\varepsilon + \varepsilon v(R). \end{aligned}$$

Since the final term can be made arbitrarily small, our function is integrable!

□

Theorem. If f is integrable then A has measure zero.

Proof. Suppose f is integrable and let $\varepsilon > 0$ be given. For every $n \in \mathbb{N}$, let

$$B_n = \{x \in R : o(f, x) \geq 1/n\}.$$

Since $A = B_1 \cup B_2 \cup \dots$, it suffices to show that each B_n has measure zero. Fix $n \in \mathbb{N}$.

Since f is integrable, we may find a partition P of R such that $U(f, P) - L(f, P) < \varepsilon/n$. Let X denote the collection of rectangles in P which intersect B_n . In particular, the elements of X cover B_n and are rectangles! Now, if $R_i \in X$, then $M_i - m_i \geq 1/n$ by definition of the oscillation function. Then,

$$\begin{aligned} \frac{1}{n} \sum_{R_i \in X} v(R_i) &\leq \sum_{R_i \in X} (M_i - m_i)v(R_i) \\ &\leq \sum_{R_i \in P} (M_i - m_i)v(R_i) \\ &= U(f, P) - L(f, P) < \frac{\varepsilon}{n}. \end{aligned}$$

In particular, $\sum_{R_i \in X} v(R_i) < \varepsilon$ and so B_n has measure (content) zero. □

Week 11 Notes.

Module 0: Equivalent Notations.

	Old Notation.	New notation.
Partition.	Grid on R .	Partition of R .
Volume.	$ R $.	$V(R)$.
Lower integral	$L \int_R f$.	$\underline{\int}_R f$.
Upper integral	$U \int_R f$.	$\overline{\int}_R f$.

Module 1: General Integrability.

Remark II.1.1. Goal.

To be able to integrate over regions other than rectangles.

Definition II.1.2:

Let $A \subseteq \mathbb{R}^n$ be bounded. Let R be a rectangle such that $A \subseteq R$. The characteristic function of A on R is:

$$\chi_A : R \rightarrow \mathbb{R}.$$

$$\chi_A(x) = \begin{cases} 1 & x \in A \\ 0 & x \notin A. \end{cases}$$

Definition II.1.3:

Let $A \subseteq \mathbb{R}^n$ be bounded and R be a rectangle with $A \subseteq R$. We extend $f: R \rightarrow \mathbb{R}$ by setting $f(x) = 0$ for all $x \in R \setminus A$. We say $f: A \rightarrow \mathbb{R}$ is integrable iff. $f \cdot \chi_A: R \rightarrow \mathbb{R}$ is integrable, in which we define.

$$\int_A f = \int_R f \cdot \chi_A.$$

Assignment: This definition is independent of the choice of R containing A .

Remark II.1.4:

When is $\chi_A: R \rightarrow \mathbb{R}$ integrable?

Theorem 11.1.5:

Let $A \subseteq \mathbb{R}^n$ be bounded, $A \subseteq R$ be a rectangle.

The function $\chi_A : R \rightarrow \mathbb{R}$ is integrable iff $\partial(A)$ has measure zero.

Proof: Let $a \in R$.

① $a \in \text{Int}(A)$

Then, there exists an open ball $B_\delta(a) \subseteq A$. Since

$\chi_A = 1$ on $B_\delta(a)$, χ_A is clearly continuous at a .

② $a \notin \overline{A}$

Then, $a \in \text{Int}(\mathbb{R}^n \setminus A)$ and so there exists an open ball $B_\delta(a) \subseteq \mathbb{R}^n \setminus A$. Since $\chi_A = 0$ on $B_\delta(a) \cap R$, χ_A is clearly continuous at a .

③ $a \in \overline{A} \setminus \text{Int}(A) = \partial(A)$.

Then, $a \in \overline{A}$ and $a \in \mathbb{R}^n \setminus \text{Int}(A) = \overline{\mathbb{R}^n \setminus A}$. In particular for all $\delta > 0$, there exists $x \in A$ and $y \in \mathbb{R} \setminus A$ such that $\|x - a\|, \|y - a\| < \delta$. Thus,

$$0 < \chi_A(a) \leq 1.$$

and so χ_A is not continuous at a . The result follows from our big Theorem. \square

Definition 11.1.6:

① We call A a Jordan region iff $\partial(A)$ has measure zero. (iff χ_A is integrable on $A \subseteq R$).

② If A is a Jordan region with $A \subseteq R$, we define the

volume of A by

$$\text{Vol}(A) = \int_R \chi_A = \int_A 1.$$

Proposition 11.1.7:

Let $A, B \subseteq \mathbb{R}^n$ be Jordan regions.

① $A \cup B$ is a Jordan region

② If $A \cap B = \emptyset$ and $f: A \cup B \rightarrow \mathbb{R}$ is integrable, then

$$\int_{A \cup B} f = \int_A f + \int_B f.$$

Proof sketch:

$$\begin{aligned} \text{① } \partial(A \cup B) &= \overline{A \cup B} \setminus \text{Int}(A \cup B) \\ &\subseteq (\overline{A} \cup \overline{B}) \setminus (\text{Int}(A) \cup \text{Int}(B)) \\ &\subseteq (\overline{A} \setminus \text{Int}(A)) \cup (\overline{B} \setminus \text{Int}(B)) \\ &= \partial(A) \cup \partial(B) \end{aligned}$$

② Let $A \cup B \subseteq R$.

$$\begin{aligned} \int_{A \cup B} f &= \int_R f \chi_{A \cup B} = \int_R f (\chi_A + \chi_B) \\ &= \int_R f \chi_A + \int_R f \chi_B \\ &= \int_A f + \int_B f. \end{aligned}$$

Module 2: Fubini's Theorem.

Remark 11.2.1:

Can we use multiple single-variable integrals to compute integrals of multivariable functions?

Would the order matter?

i.e. $\int_c^d \int_a^b f(x,y) dx dy = \int_a^b \int_c^d f(x,y) dy dx ? ?$

Remark 11.2.2:

Let $B \subseteq \mathbb{R}^2$ be Jordan region.

$f: B \rightarrow \mathbb{R}$ be integrable.

$$\int_B f(v) dv \equiv \iint_B f(x,y) dA.$$

$B \subseteq \mathbb{R}^3$ be Jordan region.

$f: B \rightarrow \mathbb{R}$ be integrable.

$$\int_B f(v) dv \equiv \iiint_B f(x,y,z) dV.$$

Lemma 11.2.3.

Let $R = [a,b] \times [c,d] \subseteq \mathbb{R}^2$, $f: R \rightarrow \mathbb{R}$ be bounded.

If $f(x, \cdot): [c, d] \rightarrow \mathbb{R}$ given by $f(x, \cdot)(y) = f(x, y)$ is integrable for all $x \in [a, b]$, then

$$L \iint_R f(x,y) dA \lesssim L \int_a^b \left(\int_c^d f(x,y) dy \right) dx.$$

$$\lesssim U \int_a^b \left(\int_c^d f(x,y) dy \right) dx$$

$$\lesssim U \iint_R f(x,y) dA.$$

Proof:

The middle inequality is trivial. We will prove the last inequality and leave the first for discussion.

Let $\Sigma \geq 0$ be given. Choose a partition G on R such that $U(f, G) - \Sigma \leq U \iint_R f(x, y) dA$.

Say

$$G = \{R_{ij} : 1 \leq i \leq k, 1 \leq j \leq l\}.$$

$$R_{ij} = [x_{i-1}, x_i] \times [y_{j-1}, y_j].$$

where $x_0 = a$, $x_k = b$, $y_0 = c$, $y_l = d$.

Set $M_{ij} = \sup \{f(v) : v \in R_{ij}\}$.

$$\begin{aligned} U \int_a^b \left(\int_c^d f(x, y) dy \right) dx \\ = \sum_{i=1}^k U \int_{x_{i-1}}^{x_i} \left(\sum_{j=1}^l \int_{y_{j-1}}^{y_j} f(x, y) dy \right) dx \\ \leq \sum_{i=1}^k \sum_{j=1}^l U \int_{x_{i-1}}^{x_i} \left(\int_{y_{j-1}}^{y_j} f(x, y) dy \right) dx \end{aligned}$$

(Here we use $U \int_a^b (f+g)(x) dx \leq U \int_a^b f(x) dx + U \int_a^b g(x) dx$)

$$\leq \sum_{i=1}^k \sum_{j=1}^l \int_{x_{i-1}}^{x_i} \left(\int_{y_{j-1}}^{y_j} M_{ij} dy \right) dx.$$

$$= \sum_{i=1}^k \sum_{j=1}^l M_{ij} (x_i - x_{i-1})(y_j - y_{j-1}).$$

$$= \sum_{R_{ij}} M_{ij} |R_{ij}|.$$

$$= U(f, G)$$

$$\leq U \iint_R f(x, y) dA + \Sigma$$

□.

Theorem 11.2.4 [Fubini's Theorem].

Let $R = [a, b] \times [c, d] \subseteq \mathbb{R}^2$.

$f: R \rightarrow \mathbb{R}$ be integrable.

If $f(x, \cdot)$ and $f(\cdot, y)$ are integrable over $[c, d]$ and $[a, b]$ respectively, for all $x \in [a, b]$ and $y \in [c, d]$, then.

$$\begin{aligned}\iint_R f(x, y) dA &= \int_a^b \int_c^d f(x, y) dy dx \\ &= \int_c^d \int_a^b f(x, y) dx dy.\end{aligned}$$

Proof:

Since f is integrable

$$L \iint_R f(x, y) dA = U \iint_R f(x, y) dA.$$

By the Lemma, this implies

$$L \int_a^b \int_c^d f(x, y) dy dx = U \int_a^b \int_c^d f(x, y) dy dx$$

So:

$$\iint_R f(x, y) dA = \int_a^b \left(\int_c^d f(x, y) dy \right) dx.$$

Rearrange the roles of x, y in the lemma, we proves the theorem. \square

Remark 11.2.5:

The conditions of Fubini's Theorem is met when f is continuous.

Remark 11.2.6:

We call

$$\int_{a_1}^{b_1} \int_{a_2}^{b_2} \cdots \int_{a_n}^{b_n} f(x_1, x_2, \dots, x_n) dx_1 dx_2 \cdots dx_n$$

iterated integrals.

Example 11.2.7:

$$R = [1, 2] \times [0, \pi].$$

$$\iint_R y \sin(xy) dA$$

[Note: $f, f(x, \cdot), f(\cdot, y)$ are all continuous on closed Jordan Regions \Rightarrow integrable].

$$\begin{aligned} & \text{Fubini's Theorem} \quad \int_0^\pi \int_1^2 y \sin(xy) dx dy \\ &= \int_0^\pi (-\cos(xy)) \Big|_{x=1}^{x=2} dy \\ &= \int_0^\pi -\cos 2y + \cos y dy. \\ &= \left[-\frac{1}{2} \sin 2y + \sin y \right]_0^\pi \\ &= 0. \end{aligned}$$

Module 3: Iterated Integrals.

Remark 11.3.1: Goal

Generalizing Fubini's Theorem.

Theorem 11.3.2:

$$R = [a_1, b_1] \times \cdots \times [a_n, b_n] \subseteq \mathbb{R}^n.$$

$$R_n = [a_1, b_1] \times \cdots \times [a_{n-1}, b_{n-1}]$$

If $f(x, \cdot)$ is integrable for all $x \in R_n$, then

$\int_{a_n}^{b_n} f(x, t) dt$ is integrable on R_n and

$$\int_R f(v) dv = \int_{R_n} \int_{a_n}^{b_n} f(x, t) dt dx.$$

Remark 11.3.3:

If $f: R \rightarrow \mathbb{R}$ is continuous, then

$$\int_R f(v) dv = \int_{a_1}^{b_1} \int_{a_2}^{b_2} \cdots \int_{a_n}^{b_n} f(x_1, \dots, x_n) dx_n \cdots dx_1$$

Remark 11.3.4:

We will use iterated integrals to integrate over "nice" regions which are not rectangles.

For simplicity, we shall work in \mathbb{R}^2 and \mathbb{R}^3 .

Definition 11.3.5:

① We say $A \subseteq \mathbb{R}^2$ is type 1 if


$$A = \{(x, y) : x \in [a, b], \varphi(x) \leq y \leq \psi(x)\}.$$

for some continuous $\varphi, \psi: [a, b] \rightarrow \mathbb{R}$.


② We say $A \subseteq \mathbb{R}^2$ is type 2 if

$A = \{(x, y) : y \in [a, b], \varphi(y) \leq x \leq \psi(y)\}$
 for some continuous $\varphi, \psi : [a, b] \rightarrow \mathbb{R}$.

Example 11.3.6: In \mathbb{R}^2 :



Type 2:



Definition 11.3.6: $A \subseteq \mathbb{R}^3$.

① Type 1.

$$A = \{(x, y, z) : (x, y) \in H, \varphi(x, y) \leq z \leq \psi(x, y)\}.$$

② Type 2:

$$A = \{(x, y, z) : (x, z) \in H, \varphi(x, z) \leq y \leq \psi(x, z)\}.$$

③ Type 3:

$$A = \{(x, y, z) : (y, z) \in H, \varphi(y, z) \leq x \leq \psi(y, z)\},$$

where $H \subseteq \mathbb{R}^2$ is a closed Jordan region and $\varphi, \psi : H \rightarrow \mathbb{R}$ are continuous.

Remark 11.3.7:

Regions of type 1, 2, or 3 are Jordan regions.

Theorem 11.3.8:

$A \subseteq \mathbb{R}^2$, $f: A \rightarrow \mathbb{R}$ is continuous.

① If A is type 1, so that

$$A = \{(x, y) : x \in [a, b], \varphi(x) \leq y \leq \psi(x)\},$$

for some continuous $\varphi, \psi: [a, b] \rightarrow \mathbb{R}$.

then $\int_A f(x, y) dV = \int_a^b \int_{\varphi(x)}^{\psi(x)} f(x, y) dy dx.$

② If A is type 2, so that

$$A = \{(x, y) : y \in [a, b], \varphi(y) \leq x \leq \psi(y)\},$$

then

$$\int_A f(x, y) dV = \int_a^b \int_{\varphi(y)}^{\psi(y)} f(x, y) dx dy.$$

Proof (of ①).

Let $R = [a, b] \times [c, d]$ be a rectangle containing A . Extend f to R by setting $f = 0$ on $R \setminus A$. By Fubini's Theorem,

$$\begin{aligned} \int_A f(x, y) dV &= \int_R f(x, y) dV \\ &= \int_a^b \int_c^d f(x, y) dy dx. \end{aligned}$$

However, $f(x, y) = 0$ if it is not the case that

$$\varphi(x) \leq y \leq \psi(x).$$

$\therefore \int_A f(x, y) dV = \int_a^b \int_{\varphi(x)}^{\psi(x)} f(x, y) dy dx \quad \square$

Theorem 11.3.9:

$A \subseteq \mathbb{R}^3$, $f: A \rightarrow \mathbb{R}$ is continuous

If A is type 1, then.

$$\int_A f(x, y, z) dV = \int_H \int_{\psi(x, y)}^{\psi(x, y)} f(x, y, z) dz dy$$


Type 2, Type 3 are similar.

Module 4: Examples.

Example 11.4.1:

Let $D \subseteq \mathbb{R}^2$ be the region bounded by $y=0$, $y=x^2$, $x=1$. Compute

$$\iint_D x \cos y \, dA.$$



$$\iint_D x \cos y \, dA = \int_0^1 \int_0^{x^2} x \cos y \, dy \, dx.$$

$$= \int_0^1 [x \sin y]_{y=0}^{y=x^2} \, dx.$$


$$= \int_0^1 x \sin x^2 \, dx.$$

$$= \left[-\frac{1}{2} \cos x^2 \right]_0^1.$$

$$= -\frac{1}{2} \cos(1) + \frac{1}{2}$$

Example 11.4.2:


$$\int_0^1 \int_{3y}^3 e^{x^2} \, dx \, dy$$




$$\begin{aligned}
& \int_0^1 \int_{3y}^3 e^{x^2} dx dy = \iint_D e^{x^2} dA. \\
& = \int_0^3 \int_0^{\frac{1}{3}x} e^{x^2} dy dx \\
& = \int_0^3 \left[e^{x^2} y \right]_{y=0}^{y=\frac{1}{3}x} dx \\
& = \int_0^3 \frac{1}{3}x e^{x^2} dx \\
& = \left[\frac{1}{6} e^{x^2} \right]_0^3 = \frac{1}{6} e^9 - \frac{1}{6}
\end{aligned}$$

Example 11.4.3 :

Find the volume of the tetrahedron T enclosed by $x=0$, $y=0$, $z=0$ and $2x+y+z=4$.



The base :



$$T = \{(x, y, z) : 0 \leq x \leq 2, 0 \leq y \leq -2x + 4, 0 \leq z \leq 4 - 2x - y\}$$

$$H = \{(x, y) : 0 \leq x \leq 2, 0 \leq y \leq -2x + 4\}.$$

↪ Type 1.

$$T = \{(x, y, z) : (x, y) \in H, 0 \leq z \leq 4 - 2x - y\}.$$

$$\iiint 1 \cdot dV$$

$$= \int_H^T 1 \cdot dz \, dA.$$

continuous

$$= \int_0^2 \int_0^{-2x+4} \int_0^{4-2x-y} 1 \cdot dz \, dy \, dx.$$

$$= \int_0^2 \int_0^{-2x+4} \int_{4-2x-y}^0 1 \cdot dy \, dx$$

$$= \int_0^2 \left[(4-2x)y - \frac{1}{2}y^2 \right]_{y=0}^{y=4-2x} dx$$

$$= \int_0^2 \frac{1}{2} (4-2x)^2 dx = \frac{16}{3}$$

Module 5: Change of Variables.

Remark 11.5.1:

Recall $f: A \rightarrow \mathbb{R}^m$, $A \subseteq \mathbb{R}^n$, $a \in A$. The Jacobian of f at a is:

$$Jf(a) = \det(Df(a)).$$

Theorem 11.5.2:

Let $U \subseteq \mathbb{R}^n$ be open, $A \subseteq U$ be closed Jordan region. Let $f: A \rightarrow \mathbb{R}$ be continuous and let $\psi \in C^1(U, \mathbb{R}^n)$. Suppose $\exists B \subseteq A$.

- ① $\text{Vol}(CB) \geq 0$.
- ② ψ is injective on $A \setminus B$.
- ③ $J\psi(a) \neq 0$, $\forall a \in A \setminus B$.

and suppose $f: \psi(A) \rightarrow \mathbb{R}$ is continuous. Then $\psi(A)$ is a Jordan region, f is injective on $\psi(A)$ and

$$\int_{\psi(A)} f(x) dx = \int_A f(\psi(x)) |J\psi(x)| dx.$$

Theorem 11.5.3: Polar Coordinates.

$$\begin{aligned}
 & (x, y) \quad r = \sqrt{x^2 + y^2} \\
 & \theta \quad x = r \cos \theta \\
 & \quad y = r \sin \theta \\
 & \quad 0 \leq \theta \leq 2\pi.
 \end{aligned}$$

We call (r, θ) the polar coordinates of $(x, y) \in \mathbb{R}^2$. Consider $\psi \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ given by

$$\varphi(r, \theta) = (r \cos \theta, r \sin \theta).$$

Note: φ is injective on $\mathbb{R}^2 \setminus \{(0, 0) : 0 \leq \theta \leq 2\pi\}$.

Volume zero.

$$|\mathcal{J}(\varphi(r, \theta))| = \left| \det \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix} \right| = |r| = r.$$

$$\iint_{\varphi(D)} f(x, y) dA = \iint_D f(r \cos \theta, r \sin \theta) r dA$$

Example 11.5-4:

$$\iint_D \cos(r^2) dA.$$

D is the region bounded by $x^2 + y^2 = 9$ and above the x -axis.

$$D = \{(r, \theta) : 0 \leq r \leq 3, 0 \leq \theta \leq \pi\}.$$


$$\iint_D \cos(r^2) dA = \iint_D \cos(r^2) r dA.$$

$$= \int_0^\pi \int_0^3 \cos(r^2) r dr d\theta$$

$$= \int_0^\pi \left[\frac{1}{2} \sin(r^2) \right]_0^3 dr$$

$$= \int_0^\pi \frac{1}{2} \sin 9 \theta = \frac{\pi}{2} \sin 9.$$

Theorem 11.5-5. Cylindrical Coordinates.



We call (r, θ, z) the cylindrical coordinates of (x, y, z) .

$$\varphi(r, \theta, z) = (r \cos \theta, r \sin \theta, z).$$

$$|\mathcal{J} \varphi(r, \theta, z)| = \left| \det \begin{bmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \right| \geq r.$$

$$\underset{\Psi(CA)}{\iiint f(x, y, z) dV} = \iiint_A f(r \cos \theta, r \sin \theta, z) r dV$$

Example 11.5-6.


$$\iiint_A e^z dV$$

where A is enclosed by

① The paraboloid $z = 1 + x^2 + y^2$.

② The cylinder $x^2 + y^2 = 5$.

③ The xy -plane.




$$A = \{(r, \theta, z) : 0 \leq r \leq \sqrt{5}, 0 \leq \theta \leq 2\pi, 0 \leq z \leq 1 + r^2\}$$

$$\begin{aligned} \iiint_A e^z dV &= \int_0^{\sqrt{5}} \int_0^{2\pi} \int_0^{1+r^2} e^z r dz d\theta dr \\ &= \int_0^{\sqrt{5}} \int_0^{2\pi} r e^{1+r^2} - r d\theta dr. \end{aligned}$$

$$\begin{aligned}
 &= 2\pi \int_0^5 r e^{1tr^2} - r dr \\
 &= 2\pi \left[\frac{1}{2} e^{1tr^2} - \frac{1}{2} r^2 \right]_0^5 \\
 &= \pi (e^6 - 5 - 1) .
 \end{aligned}$$

Theorem 11.5.7: Spherical Coordinates.



$$x = r \cos \theta$$

$$r = \rho \sin \phi$$

$$y = r \sin \theta$$

$$z = \rho \cos \phi .$$

$$x = \rho \sin \phi \cos \theta, y = \sin \theta \sin \phi .$$

$$z = \rho \cos \phi, x^2 + y^2 + z^2 = \rho^2 .$$

Consider $\Psi(\rho, \theta, \phi) = (\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi)$

$$\begin{aligned}
 |\nabla \Psi(\rho, \theta, \phi)| &= \left| \det \begin{bmatrix} \sin \phi \cos \theta & \rho \sin \phi \sin \theta & \rho \cos \phi \cos \theta \\ \sin \phi \sin \theta & \rho \sin \phi \cos \theta & \rho \cos \phi \sin \theta \\ \cos \phi & 0 & -\rho \sin \phi \end{bmatrix} \right| \\
 &= \rho^2 \sin \phi .
 \end{aligned}$$

$$\iiint_A f(x, y, z) dV = \iiint_A f(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \rho^2 \sin \phi dV$$

Example 11.5.8.

Find the volume of the sphere $x^2+y^2+z^2=a^2$.

$S = \{(r, \theta, \phi) : 0 \leq r \leq a, 0 \leq \theta \leq 2\pi, 0 \leq \phi \leq \pi\}$.

$$\text{Vol}(S) = \int_S 1 \, dV$$

$$= \iiint_S 1 \, dx \, dy \, dz$$

$$= \int_0^a \int_0^{2\pi} \int_0^\pi r^2 \sin \phi \, d\phi \, d\theta \, dr.$$

$$= \int_0^a \int_0^{2\pi} [-r^2 \cos \phi]_0^\pi \, d\theta \, dr$$

$$= \int_0^a \int_0^{2\pi} 2r^2 \, d\theta \, dr$$

$$= 2\pi \int_0^a 2r^2 \, dr$$


$$= 2\pi \cdot \frac{2}{3}r^3 \Big|_0^a = \frac{4\pi}{3}a^3.$$

Example 11.5.9.

Find the volume of the solid which

① lies above the cone $z = \sqrt{x^2+y^2}$.

② below the sphere $x^2+y^2+z^2=z$.



$$x^2+y^2+z^2=z \Leftrightarrow x^2+y^2+(z-\frac{1}{2})^2=\frac{1}{4}$$

$$\text{Cone: } \rho \cos \phi = \sqrt{\rho^2 \sin^2 \phi \cos^2 \theta + \rho^2 \sin^2 \phi \sin^2 \theta}.$$

$$= \rho \sin \phi.$$

$$C = \{(\rho, \theta, \phi) : \rho = 0 \text{ or } \phi = \frac{\pi}{4}\}.$$

$$\text{Sphere: } \rho^2 = \rho \cos \phi$$

$$S = \{(\rho, \theta, \phi) : \rho = 0 \text{ or } \rho = \cos \phi\}.$$

Let D be the above solid.

$$\iiint_D 1 \, dV = \iiint_D \rho^2 \sin \phi \, d\rho \, d\phi \, d\theta.$$

$$= \int_0^{2\pi} \int_0^{\pi/4} \int_0^{\cos \phi} \rho^2 \sin \phi \, d\rho \, d\phi \, d\theta.$$

$$= \frac{\pi}{8}.$$