
Mach number

DS2V | Argon, $M_\infty = 10$, $T_\infty = 200$ K, $T_w = 500$ K | Matched late averages near $tU_\infty/D = 18$–20




$q =$ minor/major semiaxis. $D = 0.3048$ m; common nose at $x = 0$. Each paired geometry has the same two semiaxes.

White marker: axis shock from $\rho/\rho_\infty = 2.5$. Lower half mirrored. Rightmost panels repeat the same circular reference.

Density ratio ρ/ρ_∞ (square-root color spacing)

DS2V | Argon, $M_\infty = 10$, $T_\infty = 200$ K, $T_w = 500$ K | Matched late averages near $tU_\infty/D = 18$–20




$q =$ minor/major semiaxis. $D = 0.3048$ m; common nose at $x = 0$. Each paired geometry has the same two semiaxes.

White marker: axis shock from $\rho/\rho_\infty = 2.5$. Lower half mirrored. Rightmost panels repeat the same circular reference.

Translational temperature (K)

DS2V | Argon, $M_\infty = 10$, $T_\infty = 200$ K, $T_w = 500$ K | Matched late averages near $tU_\infty/D = 18$–20




$q =$ minor/major semiaxis. $D = 0.3048$ m; common nose at $x = 0$. Each paired geometry has the same two semiaxes.

White marker: axis shock from $\rho/\rho_\infty = 2.5$. Lower half mirrored. Rightmost panels repeat the same circular reference.

Pressure (Pa)

DS2V | Argon, $M_\infty = 10$, $T_\infty = 200$ K, $T_w = 500$ K | Matched late averages near $tU_\infty/D = 18$–20



$q =$ minor/major semiaxis. $D = 0.3048$ m; common nose at $x = 0$. Each paired geometry has the same two semiaxes.

White marker: axis shock from $\rho/\rho_\infty = 2.5$. Lower half mirrored. Rightmost panels repeat the same circular reference.