


Causal Data Science

Lecture 5:2 Backdoor adjustment

Lecturer: Sara Magliacane

UvA - Spring 2023



Identification strategies for causal effects

- Given a causal graph G , an **identification strategy** is a formula to estimate an interventional distribution from a combination of observational ones


- **Backdoor criterion (this class), Adjustment criterion (next class)**

$$p(x_j | \text{do}(x_i)) = \int_{x_Z} p(x_j | x_i, x_Z) p(x_Z) dx_Z$$

- **Frontdoor criterion (next class)**

$$p(x_j | \text{do}(x'_i)) = \int_{x_M} p(x_M | x'_i) \int_{x_i} p(x_j | x_M, x_i) p(x_i) dx_i$$

- **Instrumental variables (next class)**




Valid adjustment sets

- Given a causal Bayesian network (G, p) with DAG $G = (\mathbf{V}, \mathbf{E})$
- We call **(valid) adjustment sets** for the total causal effect of the **treatment** X_i on the **outcome** X_j with $i \neq j$, the sets $\mathbf{Z} \subseteq \mathbf{V} \setminus \{i, j\}$ such that:

$$p(x_j | \text{do}(x_i)) = \int_{x_{\mathbf{Z}}} p(x_j | x_i, x_{\mathbf{Z}}) p(x_{\mathbf{Z}}) dx_{\mathbf{Z}}$$

$$P(X_j | \text{do}(X_i = x_i)) = \sum_{x_{\mathbf{Z}}} P(X_j | X_i = x_i, X_{\mathbf{Z}} = x_{\mathbf{Z}}) P(X_{\mathbf{Z}} = x_{\mathbf{Z}})$$


(ADJUSTMENT FORMULA)



Additional assumptions for adjustment: Positivity

- Adjustment formula for estimating the causal effect of X_i on X_j with adjustment sets $\mathbf{Z} \subseteq \mathbf{V} \setminus \{i, j\}$:

$$p(x_j | \text{do}(x_i)) = \int_{x_{\mathbf{Z}}} p(x_j | x_i, x_{\mathbf{Z}}) p(x_{\mathbf{Z}}) dx_{\mathbf{Z}}$$




Additional assumptions for adjustment: Positivity

- Adjustment formula for estimating the causal effect of X_i on X_j with adjustment sets $\mathbf{Z} \subseteq \mathbf{V} \setminus \{i, j\}$:

$$p(x_j | \text{do}(x_i)) = \int_{x_{\mathbf{Z}}} p(x_j | x_i, x_{\mathbf{Z}}) p(x_{\mathbf{Z}}) dx_{\mathbf{Z}}$$

- Values in interventional regime **should be also possible** in the observational regime, specifically: $p(x_i, x_{\mathbf{Z}}) > 0$ for $x_i \in \mathcal{X}_i, x_{\mathbf{Z}} \in \mathcal{X}_{\mathbf{Z}}$
- If all women are treated and men are untreated, we cannot adjust for gender




Valid adjustment sets

- Given a causal Bayesian network (G, p) with DAG $G = (\mathbf{V}, \mathbf{E})$
- We call **(valid) adjustment sets** for the total causal effect of the **treatment** X_i on the **outcome** X_j with $i \neq j$, the sets $\mathbf{Z} \subseteq \mathbf{V} \setminus \{i, j\}$ such that:

$$p(x_j | \text{do}(x_i)) = \int_{x_{\mathbf{Z}}} p(x_j | x_i, x_{\mathbf{Z}}) p(x_{\mathbf{Z}}) dx_{\mathbf{Z}}$$

We will see two graphical criteria for identifying these sets directly from the graph: backdoor and adjustment criterion.



Sometimes less is more

- Adjusting for all possible covariates doesn't always help:

$\text{lm}(X_4 \sim X_1 + X_2)$


```
linear_regressorX12 = LinearRegression()  
linear_regressorX12.fit(X21, Y)  
linear_regressorX12.coef_[0,1]
```

```
array([29.87150906])
```

$\text{lm}(X_4 \sim X_1 + X_2 + X_3)$

```
linear_regressorX123 = LinearRegression()  
linear_regressorX123.fit(X, Y)  
linear_regressorX123.coef_[0,1]
```

```
array([0.15806091])
```




$$E[X_4 | do(X_1 = 1)] - E[X_4 | do(X_1 = 0)] = 30$$




Special case: no causal parents for treatment

- If treatment X_i has no causal parents, then the interventional distribution is the same as the observational distribution conditioned on $X_i = x_i$:



$$\begin{aligned} P(X_{\bar{v}} | \text{do}(X_i = x_i)) &= \prod_{j \in \bar{v}, j \neq i} P(x_j | X_{\text{pa}(j)}) \cdot P(x_i | \phi) \\ &= \prod_{j \in \bar{v}} P(x_j | X_{\text{pa}(j)}) \Big|_{X_i = x_i} \quad \text{eval. in } X_i = x_i \end{aligned}$$




Special case: no causal parents for treatment

- If treatment X_i has no causal parents, then the interventional distribution is the same as the observational distribution conditioned on $X_i = x_i$:


$$P(X_{\bar{v}} \mid \text{do}(X_i = x_i)) = \prod_{j \in v} P(X_j \mid X_{\text{pa}(j)}) \Big|_{X_i = x_i} = \frac{P(X_{\bar{v}})}{P(X_i \mid X_{\text{pa}(i)})} \Big|_{X_i = x_i}$$


$$= P(X_{\bar{v}} \mid X_i = x_i)$$



Special case: no causal parents for treatment

- If treatment X_i has no causal parents, then the interventional distribution is the same as the observational distribution conditioned on $X_i = x_i$:






Special case: using the parents of the treatment

- **The set of causal parents** of the treatment allow us to estimate the interventional distribution:

$$p(X_{\mathbf{V}} \mid \text{do}(X_i = x_i)) = \prod_{j \in \mathbf{V} \setminus \{i\}} p(X_j \mid X_{\text{Pa}(j)}) \mid_{X_i = x_i}$$




Special case: using the parents of the treatment

- **The set of causal parents** of the treatment allow us to estimate the interventional distribution:

$$p(X_{\mathbf{V}} \mid \text{do}(X_i = x_i)) = \prod_{j \in \mathbf{V} \setminus \{i\}} p(X_j \mid X_{\text{Pa}(j)}) \mid_{X_i = x_i}$$

$$= \frac{p(X_{\bar{v}})}{p(X_i \mid X_{\text{Pa}(i)})} \Big|_{X_i = x_i}$$




Special case: using the parents of the treatment

- **The set of causal parents** of the treatment allow us to estimate the interventional distribution:

$$p(X_{\mathbf{V}} \mid \text{do}(X_i = x_i)) = \prod_{j \in \mathbf{V} \setminus \{i\}} p(X_j \mid X_{\text{Pa}(j)}) \mid_{X_i = x_i}$$

$$= \frac{p(X_{\bar{v}})}{p(X_i \mid X_{\text{Pa}(i)})} \Big|_{X_i = x_i} = \frac{p(X_{\bar{v}}) \cdot p(X_{\text{Pa}(i)})}{p(X_i, X_{\text{Pa}(i)})} \Big|_{X_i = x_i}$$




Special case: using the parents of the treatment

- **The set of causal parents** of the treatment allow us to estimate the interventional distribution:

$$p(X_{\mathbf{V}} \mid \text{do}(X_i = x_i)) = \prod_{j \in \mathbf{V} \setminus \{i\}} p(X_j \mid X_{\text{Pa}(j)}) \mid_{X_i = x_i}$$

$$= \frac{p(X_{\bar{v}})}{p(X_i \mid X_{\text{Pa}(i)})} \Big|_{X_i = x_i} = \frac{p(X_{\bar{v}}) \cdot p(X_{\text{Pa}(i)})}{p(X_i, X_{\text{Pa}(i)})} \Big|_{X_i = x_i}$$

$$= p(X_{\bar{v}, \{i, \text{Pa}(i)\}} \mid X_i, X_{\text{Pa}(i)}) \cdot p(X_{\text{Pa}(i)}) \Big|_{X_i = x_i}$$




Special case: using the parents of the treatment

- **The set of causal parents** of the treatment allow us to estimate the interventional distribution:

$$p(X_{\mathbf{V}} \mid \text{do}(X_i = x_i)) = \prod_{j \in \mathbf{V} \setminus \{i\}} p(X_j \mid X_{\text{Pa}(j)}) \mid_{X_i = x_i}$$

$$= P(X_{\mathbf{V} \setminus \{i, \text{Pa}(i)\}} \mid X_i, X_{\text{Pa}(i)}) \cdot P(X_{\text{Pa}(i)}) \Big|_{X_i = x_i}$$

$$= p(X_{\mathbf{V} \setminus \{i, \text{Pa}(i)\}} \mid X_i = x_i, X_{\text{Pa}(i)}) p(X_{\text{Pa}(i)})$$



Special case: using the parents of the treatment

- **The set of causal parents** of the treatment allow us to estimate the interventional distribution:


$$p(X_V | \text{do}(X_i = x_i)) = p(X_{V \setminus \{i, \text{Pa}(i)\}} | X_i = x_i, X_{\text{Pa}(i)})p(X_{\text{Pa}(i)})$$

- We can marginalise out the other variables which are not X_i , X_j or $X_{\text{Pa}(i)}$

$$p(X_j | \text{do}(X_i = x_i)) = \int_{x_{\text{Pa}(i)}} p(X_j | X_i = x_i, X_{\text{Pa}(i)})p(X_{\text{Pa}(i)})dx_{\text{Pa}(i)}$$



$$p(x_j | \text{do}(x_i)) = \int_{x_Z} p(x_j | x_i, x_Z)p(x_Z)dx_Z$$


**The parents of the treatment
are a valid adjustment set**



Special case: using the parents of the treatment

- The causal parents of the treatment are a valid adjustment set
- Sometimes some of the parents might not be observed
- We want to list **all valid adjustment sets**







Identifiability of an interventional distribution

- We say an interventional distribution $p(X_i | \text{do}(X_j = x_j))$ is **identifiable**, if we can rewrite it using observational distributions
 - For example if a **valid adjustment set** exists
 - But also, front door criterion, instrumental variables


$$p(X_4 | \text{do}(X_2)) = ?$$






Backdoor criterion intuition

- Directed path from i to j are **causal (both directly and indirectly)**
- **Backdoor paths** (all paths that start with an arrow into $i \leftarrow \dots j$)
induce **spurious associations**
- If we block the spurious associations, we only get the causal effect




Backdoor criterion intuition

- Directed path from i to j are **causal (both directly and indirectly)**
- **Backdoor paths** (all paths that start with an arrow into $i \leftarrow \dots j$)
induce **spurious associations**
- If we block the spurious associations, we only get the causal effect
- Including **descendants of i that are also ancestors of j (*mediators*)**
would **block causal paths**
- Including **descendants of i that are also descendants of j** would
create **collider bias**




Backdoor criterion [Pearl 2009]

- Given a CBN (G, p) with $G = (\mathbf{V}, \mathbf{E})$, a set $\mathbf{Z} \subseteq \mathbf{V} \setminus \{i, j\}$ satisfies the **backdoor criterion** for estimating the causal effect of X_i on X_j with $i \neq j$:
- \mathbf{Z} does **not contain any descendant of i** , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, **and**



Backdoor criterion [Pearl 2009]


- Given a CBN (G, p) with $G = (\mathbf{V}, \mathbf{E})$, a set $\mathbf{Z} \subseteq \mathbf{V} \setminus \{i, j\}$ satisfies the **backdoor criterion** for estimating the causal effect of X_i on X_j with $i \neq j$:
 - \mathbf{Z} does **not contain any descendant of i** , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, **and**
 - \mathbf{Z} blocks all **backdoor paths** from i to j (all paths that start with an arrow into $i \leftarrow \dots j$)



Backdoor criterion [Pearl 2009]

- Given a CBN (G, p) with $G = (\mathbf{V}, \mathbf{E})$, a set $\mathbf{Z} \subseteq \mathbf{V} \setminus \{i, j\}$ satisfies the **backdoor criterion** for estimating the causal effect of X_i on X_j with $i \neq j$:
 - \mathbf{Z} does **not contain any descendant of i** , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, **and**
 - \mathbf{Z} blocks all **backdoor paths** from i to j (all paths that start with an arrow into $i \leftarrow \dots j$)

The backdoor criterion finds **some (not necessarily all)** valid adjustment sets




Backdoor criterion [Pearl 2009]

- Given a CBN (G, p) with $G = (\mathbf{V}, \mathbf{E})$, a set $\mathbf{Z} \subseteq \mathbf{V} \setminus \{i, j\}$ satisfies the **backdoor criterion** for estimating the causal effect of X_i on X_j with $i \neq j$:
 - \mathbf{Z} does **not contain any descendant of i** , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, **and**
 - \mathbf{Z} blocks all **backdoor paths** from i to j (all paths that start with an arrow into $i \leftarrow \dots j$)


The adjustment criterion is an extension of backdoor that finds all of them - next class

The backdoor criterion finds **some (not necessarily all)** valid adjustment sets



Backdoor criterion example:


- \mathbf{Z} does ~~not contain any descendant of i~~ , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, ~~and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$




- X_2 doesn't have descendants except X_4

Backdoor criterion example:

- ~~\mathbf{Z} does not contain any descendant of i , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$



$$\bar{Z} = \{X_3\}$$

- A **path** between **i** and **j** is **blocked by** $\mathbf{A} \subseteq \mathbf{V} \setminus \{i, j\}$ **if** at least one condition holds:
 - ✓ There is a **non-collider** on the path that is in \mathbf{A} , **or**
 - There is a **collider** k on the path, but $k \notin \mathbf{A}$ and $\text{Desc}(k) \cap \mathbf{A} = \emptyset$




Backdoor criterion example:

- ~~\mathbf{Z} does not contain any descendant of i , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$




$$\bar{\mathbf{Z}} = \{X_3\}$$

- A **path** between i and j is **blocked by** $\mathbf{A} \subseteq \mathbf{V} \setminus \{i, j\}$ **if** at least one condition holds:
 - ✓ There is a **non-collider** on the path that is in \mathbf{A} , **or**
 - There is a **collider** k on the path, but $k \notin \mathbf{A}$ and $\text{Desc}(k) \cap \mathbf{A} = \emptyset$




Backdoor criterion example:

- ~~\mathbf{Z} does not contain any descendant of i , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$




$$\bar{\mathbf{Z}} = \{X_3\}$$

- A **path** between **i and j** is **blocked by $\mathbf{A} \subseteq \mathbf{V} \setminus \{i, j\}$ if** at least one condition holds:
 - ✓ There is a **non-collider** on the path that is in \mathbf{A} , or
 - There is a **collider** k on the path, but $k \notin \mathbf{A}$ and $\text{Desc}(k) \cap \mathbf{A} = \emptyset$




Backdoor criterion example:

- ~~\mathbf{Z} does not contain any descendant of i , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$




$$\bar{\mathbf{Z}} = \{X_3\}$$

- A **path** between i and j is **blocked by** $\mathbf{A} \subseteq \mathbf{V} \setminus \{i, j\}$ **if** at least one condition holds:
 - There is a **non-collider** on the path that is in \mathbf{A} , **or**
 - ✗ There is a **collider** k on the path, but $k \notin \mathbf{A}$ and $\text{Desc}(k) \cap \mathbf{A} = \emptyset$




Backdoor criterion example:

- ~~\mathbf{Z} does not contain any descendant of i , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$




- A **path** between i and j is **blocked by** $\mathbf{A} \subseteq \mathbf{V} \setminus \{i, j\}$ **if** at least one condition holds:
 - ✓ There is a **non-collider** on the path that is in \mathbf{A} , **or**
 - There is a **collider** k on the path, but $k \notin \mathbf{A}$ and $\text{Desc}(k) \cap \mathbf{A} = \emptyset$

$\bar{Z} = \{X_3, X_1\}$ or
 $\bar{Z} = \{X_3, X_5\}$




Backdoor criterion example:

- ~~Z does not contain any descendant of i , $\text{Desc}(i) \cap Z = \emptyset$, and~~
- Z blocks all backdoor paths from i to j , i.e. $i \leftarrow \dots j$




- A path between i and j is blocked by $\mathbf{A} \subseteq V \setminus \{i, j\}$ if at least one condition holds:
 - There is a non-collider on the path that is in \mathbf{A} , or
 - There is a collider k on the path, but $k \notin \mathbf{A}$ and $\text{Desc}(k) \cap \mathbf{A} = \emptyset$

Is $Z = \{X_1, X_3\}$ an adjustment set that satisfies the backdoor criterion?



Backdoor criterion example:


- \mathbf{Z} does ~~not contain any descendant of i~~ , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, ~~and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$



$$\bar{Z} = \{X_3, X_1\}$$


$$\bar{Z} = \{X_3, X_5\}$$

$$\bar{Z} = \{X_3, X_5, X_1\}$$




Backdoor criterion example 2:

- ~~\mathbf{Z} does not contain any descendant of i , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$




Does $Z = \emptyset$ satisfy the backdoor criterion for $P(X_4 | do(X_2))$?




Backdoor criterion example 2:

- ~~\mathbf{Z} does not contain any descendant of i , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and~~
- \mathbf{Z} blocks all backdoor paths from i to j , i.e. $i \leftarrow \dots j$




$$P(X_4 | \text{do}(X_2 = x_2)) = \sum_{x_Z} P(X_4 | X_2 = x_2, X_Z = x_Z) P(X_Z = x_Z) = P(X_4 | X_2 = x_2)$$




Backdoor criterion example 2:

- ~~\mathbf{Z} does not contain any descendant of i , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and~~
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$



$$X_2 \perp\!\!\!\perp X_4$$

$$P(X_4 | \text{do}(X_2 = x_2)) = P(X_4 | X_2 = x_2) = P(X_4)$$



Backdoor criterion example 4:

- **Z** does **not contain any descendant of i** , $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, **and**
- **Z** blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$

$\text{lm}(X_4 \sim X_1 + X_2)$

```
linear_regressorX12 = LinearRegression()  
linear_regressorX12.fit(X21, Y)  
linear_regressorX12.coef_[0,1]
```



array([29.87150906])



$\text{lm}(X_4 \sim X_1 + X_2 + X_3)$


```
linear_regressorX123 = LinearRegression()  
linear_regressorX123.fit(X, Y)  
linear_regressorX123.coef_[0,1]
```

array([0.15806091])



$X_3 \notin \bar{Z}$

$\bar{Z} = \{X_2\}$




Linear SCMs - adjustment can be done with regression

- One can prove that in linear SCMs (not just Gaussian) the formula:

$$p(x_j | \text{do}(x_i)) = \int_{x_Z} p(x_j | x_i, x_Z) p(x_Z) dx_Z$$

is equivalent to the **coefficient of X_i for the linear regression of X_j on X_i, X_Z**



Linear SCMs - adjustment can be done with regression

- One can prove that in linear SCMs (not just Gaussian) the formula:


$$p(x_j | \text{do}(x_i)) = \int_{x_Z} p(x_j | x_i, x_Z) p(x_Z) dx_Z$$


is equivalent to the coefficient of X_i for the linear regression of X_j on X_i, X_Z

lm($X_4 \sim X_1 + X_2$)

```
linear_regressorX12 = LinearRegression()  
linear_regressorX12.fit(X21, Y)  
linear_regressorX12.coef_[0,1]
```


```
array([29.87150906])
```






Example - effect of smoking on adult asthma?


- Pretreatment variables are not always ok to adjust for
- Childhood asthma unblocks backdoor path
 - M-bias
 - Cannot adjust only on that





Backdoor criterion exercise in Canvas

- \mathbf{Z} does **not contain any descendant of i** , i.e. $\text{Desc}(i) \cap \mathbf{Z} = \emptyset$, and
- \mathbf{Z} blocks all **backdoor paths** from i to j , i.e. $i \leftarrow \dots j$



$$P(X_5 | \text{do}(X_1)) = ?$$



Identification strategies for causal effects

- Given a causal graph G , an **identification strategy** is a formula to estimate an interventional distribution from a combination of observational ones


- **Backdoor criterion (this class), Adjustment criterion (next class)**

$$p(x_j | \text{do}(x_i)) = \int_{x_Z} p(x_j | x_i, x_Z) p(x_Z) dx_Z$$


- **Frontdoor criterion (next class)**

$$p(x_j | \text{do}(x'_i)) = \int_{x_M} p(x_M | x'_i) \int_{x_i} p(x_j | x_M, x'_i) p(x_i) dx_i$$

- **Instrumental variables (next class)**



Questions?



<https://xkcd.com/2048/>