

Coursework I

CID number: A mysterious number series

MATH40007: Introduction to Applied Mathematics, 2023

**Imperial College
London**

July 30, 2023

Problem 1

Coursework background: Let $N > 1$ be a positive integer. Given an N -node graph viewed as a resistive electric circuit with all edges having unit conductance, the resistance matrix \mathbf{R} is defined as the matrix with components R_{ij} given by the effective resistance between node i (set to unit voltage) and node j (grounded) with all diagonal elements taken to vanish, i.e. $R_{ii} = 0$. Recall that the effective resistance is just the reciprocal of the effective conductance C_{eff} introduced in lectures.

The total effective resistance of the graph is defined in terms of this resistance matrix as

$$R^{(total)} = \sum_{i < j} R_{ij}.$$

Let the orthonormal (i.e., unit length and mutually orthogonal) eigenvectors and corresponding eigenvalues of the N - by N graph Laplacian \mathbf{K} be denoted by

$$\{\mathbf{e}_k \mid k = 1, \dots, N\}, \quad \{\lambda_k \mid k = 1, \dots, N\},$$

where we take

$$\mathbf{e}_1 = \frac{1}{\sqrt{N}} \begin{pmatrix} 1 \\ 1 \\ \cdot \\ \cdot \\ 1 \end{pmatrix}, \quad \lambda_1 = 0$$

and where we denote the elements of \mathbf{e}_k for $k = 2, \dots, N$ as

$$\mathbf{e}_k = \begin{pmatrix} e_{k1} \\ e_{k2} \\ \cdot \\ \cdot \\ e_{kN} \end{pmatrix}, \quad k = 2, \dots, N$$

The following two formulas can be established relating the elements of the resistance matrix and the total effective resistance to the eigenvectors/eigenvalues of \mathbf{K} :

$$R_{ij} = \sum_{k=2}^N \frac{1}{\lambda_k} (e_{ki} - e_{kj})^2$$

and

$$R^{(total)} = N \sum_{k=2}^N \frac{1}{\lambda_k}.$$

Coursework exercise: Consider the 4-node graph shown in Figure 1:




Figure 1: A 4-node graph viewed as an electric circuit.

- (a) Using the node labelling given in the figure, find the resistance matrix \mathbf{R} for this graph.
 - (b) Hence, using your results from part (a), compute R^{total} for this graph using formula (1).
 - (c) Find the eigenvalues $\{\lambda_k \mid k = 1, \dots, 4\}$ and orthonormal eigenvectors $\{\mathbf{e}_k \mid k = 1, \dots, 4\}$ of the graph Laplacian \mathbf{K} .
 - (d) Use the results of parts (a) and (c) to verify formula (2).
 - (e) Use the results of parts (b) and (c) to verify formula (3).
 - (f) Prove formula (2). Hint: consider using the eigenvectors as a basis.
- [Note: you are not required to prove formula (3)].

Solution.

- (a) As each node in the graph is connected to three edges, the degree matrix \mathbf{D} of the graph is:

$$\mathbf{D} = \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix}$$

Since all the nodes in the graph are connected with each other, the adjacent matrix \mathbf{W} of the graph is:

$$\mathbf{W} = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

since all the nodes in the graph are connected with each other.

Therefore, the Laplacian matrix \mathbf{K} of the graph is:

$$\mathbf{K} = \mathbf{D} - \mathbf{W} = \begin{pmatrix} 3 & -1 & -1 & -1 \\ -1 & 3 & -1 & -1 \\ -1 & -1 & 3 & -1 \\ -1 & -1 & -1 & 3 \end{pmatrix}$$

Since the graph is complete, it follows that the effective conductances between every pair of two nodes are the same. So, we could only calculate the conductance of one pair of the 4 nodes. We can assume that the voltage of node ① is 1 and the node ④ is grounded. Then, set the effective conductance between node ① and node ④ is \mathbf{C}_{eff} . We have this

$$\mathbf{K}\mathbf{x} = \mathbf{f}$$

We suppose the potentials at ② and ③ are x_1 and x_2 respectively. As KCL holds at node ② and ③, it follows that

$$\begin{pmatrix} 3 & -1 & -1 & -1 \\ -1 & 3 & -1 & -1 \\ -1 & -1 & 3 & -1 \\ -1 & -1 & -1 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ x_2 \\ x_3 \\ 0 \end{pmatrix} = \begin{pmatrix} \mathbf{C}_{\text{eff}} \\ 0 \\ 0 \\ -\mathbf{C}_{\text{eff}} \end{pmatrix}$$

Because the nodes ④ is grounded, we can cancel the fourth row and the fourth column of the Laplacian matrix. Then, the grounded Laplacian matrix, with node ④ grounded, gives the linear system

$$\hat{\mathbf{K}}\hat{\mathbf{x}} = \begin{pmatrix} 3 & -1 & -1 \\ -1 & 3 & -1 \\ -1 & -1 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \mathbf{C}_{\text{eff}} \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{cases} 3 - x_2 - x_3 = \mathbf{C}_{\text{eff}} \\ -1 + 3x_2 - x_3 = 0 \\ -1 - x_2 + 3x_3 = 0 \end{cases} \Rightarrow \begin{cases} \mathbf{C}_{\text{eff}} = 2 \\ x_2 = \frac{1}{2} \\ x_3 = \frac{1}{2} \end{cases}$$

Therefore, the effective conductance between each pair of the nodes is 2. As the effective resistance is the reciprocal of the effective conductance, it follows that

$$\mathbf{R}_{\text{eff}} = \frac{1}{\mathbf{C}_{\text{eff}}} = \frac{1}{2}.$$

The graph is complete, so the effective resistances between every pair of two nodes are the same. Hence, by the definition of the resistance matrix \mathbf{R} for this graph is

$$\mathbf{R} = \begin{pmatrix} 0 & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 0 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 0 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix}$$

(b)

$$R^{(total)} = \sum_{i < j} R_{ij} = \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} = \frac{1}{2} \times 6 = 3$$

(c) The graph Laplacian matrix \mathbf{K} is
$$\begin{pmatrix} 3 & -1 & -1 & -1 \\ -1 & 3 & -1 & -1 \\ -1 & -1 & 3 & -1 \\ -1 & -1 & -1 & 3 \end{pmatrix}$$

so, the characteristic polynomial of \mathbf{K} is

$$\det(\lambda I_n - \mathbf{K}) = \begin{vmatrix} \lambda - 3 & 1 & 1 & 1 \\ 1 & \lambda - 3 & 1 & 1 \\ 1 & 1 & \lambda - 3 & 1 \\ 1 & 1 & 1 & \lambda - 3 \end{vmatrix} = \lambda(\lambda - 4)^3.$$

$$\lambda(\lambda - 4)^3 = 0 \implies \lambda = 0 \text{ or } 4.$$

Hence, we have $\lambda_1 = 0$, $\lambda_2 = 4$, $\lambda_3 = 4$, $\lambda_4 = 4$.

$$\{\lambda_k | k = 1, \dots, 4\} = \{\lambda_1 = 0, \lambda_2 = 4, \lambda_3 = 4, \lambda_4 = 4\}$$

When $\lambda_1 = 0$, the corresponding eigenvector is

$$e_1 = \frac{1}{\sqrt{4}} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

When $\lambda_2 = \lambda_3 = \lambda_4 = 4$, we have

$$\mathbf{K}\mathbf{x} = 4\mathbf{x}$$

$$\mathbf{K}\mathbf{x} - 4\mathbf{x} = 0$$

$$(\mathbf{K} - 4\mathbf{I})\mathbf{x} = 0$$

$$\begin{pmatrix} -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = 0$$

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = 0$$

$$x_1 + x_2 + x_3 + x_4 = 0$$

Therefore, the eigenspace E_4 of \mathbf{K} is the right null space of the matrix

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$
 It is easy to know that the rank of this matrix is 1 by elementary row operations and this is a 4×4 matrix. Hence, by the rank nullity theorem, the dimension of the eigenspace of the matrix is 3. Therefore,

$$E_4 = \text{span} \left\{ \begin{pmatrix} -1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ 0 \\ 1 \end{pmatrix} \right\}$$

By some calculation, we can find one collection of the orthogonal eigenvectors are:

$$e_1 = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

$$e_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \\ 0 \\ 0 \end{pmatrix}$$

$$e_3 = \begin{pmatrix} -\frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{6}} \\ \frac{\sqrt{6}}{3} \\ 0 \end{pmatrix}$$

$$e_4 = \begin{pmatrix} -\frac{1}{2\sqrt{3}} \\ \frac{1}{2\sqrt{3}} \\ -\frac{1}{2\sqrt{3}} \\ \frac{\sqrt{3}}{2} \end{pmatrix}$$

It is easy to check their length is unit and they are mutually orthogonal. Hence,

$$\{e_k | k = 1, \dots, 4\} = \{e_1, e_2, e_3, e_4\} = \left\{ \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -\frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{6}} \\ \frac{\sqrt{6}}{3} \\ 0 \end{pmatrix}, \begin{pmatrix} -\frac{1}{2\sqrt{3}} \\ \frac{1}{2\sqrt{3}} \\ -\frac{1}{2\sqrt{3}} \\ \frac{\sqrt{3}}{2} \end{pmatrix} \right\}$$

(d) The formula (2) is

$$R_{ij} = \sum_{k=2}^N \frac{1}{\lambda_k} (e_{ki} - e_{kj})^2$$

$$\text{As } \lambda_2 = \lambda_3 = \lambda_4 = 4 \text{ and } e_1 = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} e_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \\ 0 \\ 0 \end{pmatrix} e_3 = \begin{pmatrix} -\frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{6}} \\ \frac{\sqrt{6}}{3} \\ 0 \end{pmatrix} e_4 = \begin{pmatrix} -\frac{1}{2\sqrt{3}} \\ -\frac{1}{2\sqrt{3}} \\ -\frac{1}{2\sqrt{3}} \\ \frac{\sqrt{3}}{2} \end{pmatrix}$$

$$\begin{aligned} R &= \begin{pmatrix} 0 & \frac{1}{4}(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}})^2 & \frac{1}{4}[\frac{1}{2} + (\frac{\sqrt{6}}{3} + \frac{1}{\sqrt{6}})^2] & \frac{1}{4}[\frac{1}{2} + \frac{1}{6} + (\frac{\sqrt{3}}{2} + \frac{1}{2\sqrt{3}})^2] \\ \frac{1}{4}(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}})^2 & 0 & \frac{1}{4}[\frac{1}{2} + (\frac{\sqrt{6}}{3} + \frac{1}{\sqrt{6}})^2] & \frac{1}{4}[\frac{1}{2} + \frac{1}{6} + (\frac{\sqrt{3}}{2} + \frac{1}{2\sqrt{3}})^2] \\ \frac{1}{4}[\frac{1}{2} + (\frac{\sqrt{6}}{3} + \frac{1}{\sqrt{6}})^2] & \frac{1}{4}[\frac{1}{2} + (\frac{\sqrt{6}}{3} + \frac{1}{\sqrt{6}})^2] & 0 & \frac{1}{4}[\frac{1}{2} + \frac{1}{6} + (\frac{\sqrt{3}}{2} + \frac{1}{2\sqrt{3}})^2] \\ \frac{1}{4}[\frac{1}{2} + \frac{1}{6} + (\frac{\sqrt{3}}{2} + \frac{1}{2\sqrt{3}})^2] & \frac{1}{4}[\frac{1}{2} + \frac{1}{6} + (\frac{\sqrt{3}}{2} + \frac{1}{2\sqrt{3}})^2] & \frac{1}{4}[\frac{1}{2} + \frac{1}{6} + (\frac{\sqrt{3}}{2} + \frac{1}{2\sqrt{3}})^2] & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 0 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 0 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix} \end{aligned}$$

which is the same as the result in part (a). Therefore, the formula (2) is correct.

(e) The formula (3) is

$$R^{(total)} = N \sum_{k=2}^N \frac{1}{\lambda_k}$$

Since $N = 4$ and $\lambda_2 = \lambda_3 = \lambda_4 = 4$,

$$R^{(total)} = N \sum_{k=2}^N \frac{1}{\lambda_k} = 4 \times (\frac{1}{4} + \frac{1}{4} + \frac{1}{4}) = 3.$$

which is the same as the result in part (b). Therefore, the formula (3) is correct.

(f) *Proof.* We use the eigenvectors $E = \{e_1, e_2, e_3, e_4\}$ as a basis of \mathbb{R}^4 . Regard \mathbf{K} as a linear transformation,

$$\mathbf{K} : \mathbb{R}^4 \rightarrow \mathbb{R}^4$$

Set the standard basis of \mathbb{R}^4 is B . From the Linear Algebra and Groups course, we have basis change formula, where

$${}_B[id]_E = \begin{pmatrix} e_{11} & e_{21} & e_{31} & e_{41} \\ e_{12} & e_{22} & e_{32} & e_{42} \\ e_{13} & e_{23} & e_{33} & e_{43} \\ e_{14} & e_{24} & e_{34} & e_{44} \end{pmatrix}$$

We set ${}_B[id]_E$ as P . As the matrix ${}_B[id]_E$ is consist of four orthogonal unit vectors, the inverse of it is

$${}_E[id]_B = ({}_B[id]_E)^T = \begin{pmatrix} e_{11} & e_{12} & e_{13} & e_{14} \\ e_{21} & e_{22} & e_{23} & e_{24} \\ e_{31} & e_{32} & e_{33} & e_{34} \\ e_{41} & e_{42} & e_{43} & e_{44} \end{pmatrix}$$

By the basis change formula, we have

$$[\mathbf{K}]_E = P^{-1}\mathbf{K}P$$

As the column of P is consist of eigenvectors of \mathbf{K} , $[\mathbf{K}]_E$ is a diagonal matrix whose element on the diagonal is the eigenvalues of \mathbf{K} , we set it as D , where

$$D = \begin{pmatrix} \lambda_1 & 0 & 0 & 0 \\ 0 & \lambda_2 & 0 & 0 \\ 0 & 0 & \lambda_3 & 0 \\ 0 & 0 & 0 & \lambda_4 \end{pmatrix}$$

Hence,

$$D = P^{-1}\mathbf{K}P$$

$$\mathbf{K} = PDP^{-1}$$

$$\begin{aligned} \mathbf{K} &= \begin{pmatrix} e_{11} & e_{21} & e_{31} & e_{41} \\ e_{12} & e_{22} & e_{32} & e_{42} \\ e_{13} & e_{23} & e_{33} & e_{43} \\ e_{14} & e_{24} & e_{34} & e_{44} \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 & 0 & 0 \\ 0 & \lambda_2 & 0 & 0 \\ 0 & 0 & \lambda_3 & 0 \\ 0 & 0 & 0 & \lambda_4 \end{pmatrix} \begin{pmatrix} e_{11} & e_{12} & e_{13} & e_{14} \\ e_{21} & e_{22} & e_{23} & e_{24} \\ e_{31} & e_{32} & e_{33} & e_{34} \\ e_{41} & e_{42} & e_{43} & e_{44} \end{pmatrix} \\ &= \begin{pmatrix} e_1 & e_2 & e_3 & e_4 \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 & 0 & 0 \\ 0 & \lambda_2 & 0 & 0 \\ 0 & 0 & \lambda_3 & 0 \\ 0 & 0 & 0 & \lambda_4 \end{pmatrix} \begin{pmatrix} e_1^T \\ e_2^T \\ e_3^T \\ e_4^T \end{pmatrix} \end{aligned}$$

Use the result of part (c),

$$\mathbf{K} = \begin{pmatrix} 1 & e_{21} & e_{31} & e_{41} \\ 1 & e_{22} & e_{32} & e_{42} \\ 1 & e_{23} & e_{33} & e_{43} \\ 1 & e_{24} & e_{34} & e_{44} \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 4 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 & 1 \\ e_{21} & e_{22} & e_{23} & e_{24} \\ e_{31} & e_{32} & e_{33} & e_{34} \\ e_{41} & e_{42} & e_{43} & e_{44} \end{pmatrix} \quad (*)$$

Then, use the way in part (a) that we calculate R_{ij} and substitue \mathbf{K} with $(*)$, we can immediately get

$$R_{ij} = \sum_{k=2}^N \frac{1}{\lambda_k} (e_{ki} - e_{kj})^2.$$

□