

MATH50003

Numerical Analysis

II.4 Interval Arithmetic

Dr Sheehan Olver

Part II

Representing Numbers

1. **Integers** via modular arithmetic
2. **Reals** via floating point
3. **Floating point arithmetic** and bounding errors
4. **Interval arithmetic** for rigorous computations

II.4 Interval Arithmetic

Use set operations with rounding to prove rigorous bounds

For sets $X, Y \subseteq \mathbb{R}$ consider the set operations

$$X + Y := \{x + y : x \in X, y \in Y\},$$

$$XY := \{xy : x \in X, y \in Y\},$$

$$X/Y := \{x/y : x \in X, y \in Y\}$$

We will use floating point arithmetic to define operations so that

$$X + Y \subseteq X \oplus Y,$$

$$XY \subseteq X \otimes Y,$$

$$X/Y \subseteq X \oslash Y$$

Proposition 3 (interval bounds). *For intervals $X = [a, b]$ and $Y = [c, d]$ satisfying $0 < a \leq b$ and $0 < c \leq d$, and $n > 0$, we have:*

$$X + Y = [a + c, b + d]$$

$$X/n = [a/n, b/n]$$

$$XY = [ac, bd]$$

Definition 14 (floating point interval arithmetic). For intervals $A = [a, b]$ and $B = [c, d]$ satisfying $0 < a \leq b$ and $0 < c \leq d$, and $n > 0$, define:

$$[a, b] \oplus [c, d] := [\mathbb{fl}^{\text{down}}(a + c), \mathbb{fl}^{\text{up}}(b + d)]$$

$$[a, b] \ominus [c, d] := [\mathbb{fl}^{\text{down}}(a - d), \mathbb{fl}^{\text{up}}(b - c)]$$

$$[a, b] \oslash n := [\mathbb{fl}^{\text{down}}(a/n), \mathbb{fl}^{\text{up}}(b/n)]$$

$$[a, b] \otimes [c, d] := [\mathbb{fl}^{\text{down}}(ac), \mathbb{fl}^{\text{up}}(bd)]$$

Example 18 (small sum).

Example 19 (exponential with intervals).

$$\exp(x) = \sum_{k=0}^n \frac{x^k}{k!} + \underbrace{\exp(t) \frac{x^{n+1}}{(n+1)!}}_{\delta_{x,n}}$$

$$\exp(X) \subseteq \left(\bigoplus_{k=0}^n X \mathbin{\bigwedge} k \mathbin{\bigcirc} k!\right) \oplus \left[\mathfrak{fl}^{\mathrm{down}}\left(-\frac{3}{(n+1)!}\right), \mathfrak{fl}^{\mathrm{up}}\left(\frac{3}{(n+1)!}\right)\right]$$

**Let's implement Interval
arithmetic in Lab 4.**