

## Sec 1 G2 Numbers — Additional Practice (Sections 2.1–2.5)

**Instructions.** Answer all questions. Use a calculator only where stated.

### 2.1 Concept of negative numbers and the number line

#### What negative numbers mean

- **Positive numbers** are above 0 (often used for *above, gain, profit*).
- **Negative numbers** are below 0 (often used for *below, loss, debt*).

#### Examples

- Temperature:  $-3^{\circ}\text{C}$  means 3 degrees below  $0^{\circ}\text{C}$ .
- Money:  $-\$20$  can mean owing \$20 or losing \$20.
- Height:  $-50$  m (below sea level),  $+50$  m (above sea level).

#### Reading a number line

- Numbers increase as you move **to the right**.
- Numbers decrease as you move **to the left**.
- So  $-7$  is **less than**  $-2$  because  $-7$  is further left.

#### Key comparisons

- Any negative number  $< 0$ .
- Any positive number  $> 0$ .
- Among negatives: the one with the **larger** magnitude is actually **smaller**.
  - Example:  $-9 < -2$ .

#### “Which is closer to 0?”

Think “distance from 0”.

- Example:  $-1$  is closer to 0 than  $-5$ , so  $-1$  is closer.

#### Practice Questions

**Q1.** A scuba diver is 6 m below sea level.

- Represent his position as an integer.
- He rises 4 m. Represent the change as an integer.

- (c) Find his new position as an integer.
- Q2.** Consider  $-3.5$ ,  $-1$ ,  $0.6$ ,  $2.4$ ,  $\frac{5}{2}$ .
- Arrange in ascending order.
  - Fill in:  $-1$  \_\_\_  $-3.5$  ( $>$ ,  $<$ ,  $=$ ).
  - Which number is closest to 0?
- Q3.** Points  $A = -4$ ,  $B = 1.5$ ,  $C = -0.5$  on a number line.
- Which point is leftmost?
  - Fill in:  $-0.5$  \_\_\_  $1.5$  ( $>$ ,  $<$ ,  $=$ ).
  - Give one integer between  $C$  and  $B$ .
- Q4.** Insert  $<$ ,  $>$ , or  $=$ .
- $-7$  \_\_\_  $-2$
  - $-3$  \_\_\_  $0$
  - $4$  \_\_\_  $-4$
  - $-1.2$  \_\_\_  $-1.20$
- Q5.** Average monthly temperature is  $28^{\circ}\text{C}$ . Differences (from 28) for months A, B, C are  $-3$ ,  $+1$ ,  $-5$  respectively.
- What does  $-5^{\circ}\text{C}$  mean?
  - Find each actual temperature.
  - Order months from coldest to hottest.
- Q6.** A shop's profit/loss over 4 weeks is  $+120$ ,  $-45$ ,  $+80$ ,  $-150$  dollars.
- Which week has the greatest profit?
  - What does  $-45$  mean?
  - Find the total over 4 weeks.
- Q7.** A hot-air balloon is 300 m above ground. Changes in height:  $+50$ ,  $-120$ ,  $+30$ ,  $-80$  (in order).
- Height after each change.
  - Greatest height reached?
  - Is it ever below ground?
- Q8.** List all integers  $n$  with  $-12 \leq n \leq 6$  that are:
- negative multiples of 4
  - positive multiples of 3
- Q9.** List all **positive and negative composite integers**  $n$  such that  $-15 < n < 15$ .
- Q10.** A game score starts at 0. Events:  $+7$ ,  $-3$ ,  $+4$ ,  $-10$ .
- Score after each event.
  - Final score.
  - When does the score become negative for the first time?

## 2.2 Addition and subtraction of integers

### Adding integers

- If the signs are the **same**, add the sizes and keep the sign.
  - Example:  $(-6) + (-4) = -10$ .
- If the signs are **different**, subtract the smaller size from the larger size, and keep the sign of the larger size.
  - Example:  $9 + (-14) = -(14 - 9) = -5$ .

### Subtracting integers

A very important rule:

- **Subtracting a negative becomes adding.**
- **Subtracting a positive stays subtracting.**

### Examples

- $12 - (-7) = 12 + 7 = 19$
- $-3 - 5 = -8$
- $-3 - (-5) = -3 + 5 = 2$

### Using context (profit/loss, rise/fall)

Treat each change as a signed number, then add them.

- Example: Start at  $-250$  m (below sea level), rise  $+310$  m  $\Rightarrow -250 + 310 = 60$  m (now above sea level).

### Practice Questions

**Q1.** Evaluate **without a calculator**.

- (a)  $-18 + 25$
- (b)  $14 - (-9)$
- (c)  $-7 - 13$
- (d)  $-30 - (-12)$

**Q2.** Evaluate **without a calculator**.

- (a)  $8 + (-3) + (-11)$
- (b)  $-5 + 14 - 9$
- (c)  $-12 - (-7) + (-10)$
- (d)  $25 - (-8) - 17$

**Q3.** Find the missing integer.

- (a)  $x + (-7) = -15$
- (b)  $12 - y = -3$

- (c)  $-4 - z = 9$
- (d)  $a - (-8) = -2$

**Q4.** A submarine is 250 m below sea level. It dives 120 m, then rises 310 m.

- (a) Write the starting position and each change using integers.
- (b) Final position (integer).
- (c) How far from sea level is it finally?

**Q5.** Morning temperature is  $-4^{\circ}\text{C}$ . It increases by  $9^{\circ}\text{C}$ , then decreases by  $12^{\circ}\text{C}$ .

- (a) Final temperature.
- (b) How many degrees below  $0^{\circ}\text{C}$  is it?

**Q6.** Heights relative to sea level:

- Place A: 430 m below sea level
- Place B: 4095 m above sea level
- Place C: 1800 m above sea level

- (a) Represent each height as an integer.
- (b) Difference in height between B and A.
- (c) Difference in height between C and A.

**Q7.** A stall's monthly profit/loss: Jan +350, Feb  $-120$ , Mar +80, Apr  $-400$ .

- (a) Which month has the greatest profit?
- (b) What does  $-400$  mean?
- (c) Total profit/loss for the 4 months.

**Q8.** City A is GMT +8, City B is GMT  $-5$ .

- (a) How many hours ahead is A compared to B?
- (b) If it is 3:00 pm in B, what time is it in A?
- (c) If it is 9:30 am in A, what time is it in B?

**Q9.** Find the sum and average of  $-3, 7, -10, 2$ .

**Q10.** Points  $P = -6$ ,  $R = 5$  on a number line. Point  $Q = x$  is exactly halfway between them. Find  $x$ .

## 2.3 Multiplication, division and combined operations of integers

### Sign rules for multiplication/division

- Same signs  $\Rightarrow$  **positive**
  - Example:  $(-6) \times (-2) = +12$
- Different signs  $\Rightarrow$  **negative**
  - Example:  $(-6) \times 2 = -12$

Same rules for division:

- $(-48) \div (-6) = +8$
- $45 \div (-9) = -5$

### **Powers (indices) with negatives**

Be careful with brackets.

- $(-3)^2 = 9$  (because the negative is inside the square)
- $-3^2 = -(3^2) = -9$  (square happens first, then the minus)

**Useful idea:** brackets tell you what is being powered.

### **Roots**

- $\sqrt{81} = 9$
- $\sqrt[3]{-125} = -5$  (cube root keeps the sign)

### **Order of operations (BODMAS)**

This is the order you calculate in:

1. **B**rackets
2. **O**rders (powers, roots)
3. **D**ivision and **M**ultiplication (left to right)
4. **A**ddition and **S**ubtraction (left to right)

#### **Example**

$$3 + 5 \times (-2) - 18 \div (-3)$$

Do  $\times$  and  $\div$  first:

- $5 \times (-2) = -10$
- $18 \div (-3) = -6$

So:

$$3 + (-10) - (-6) = 3 - 10 + 6 = -1$$

### **Brackets can change the value**

- $5 + 4 \times 3 = 5 + 12 = 17$
- $(5 + 4) \times 3 = 9 \times 3 = 27$

### **Practice Questions**

**Q1.** Evaluate **without a calculator**.

- (a)  $(-6) \times 7$
- (b)  $(-48) \div (-6)$
- (c)  $45 \div (-9)$

(d)  $(-3) \times (-5) \times 2$

**Q2.** Evaluate **without a calculator**.

(a)  $-3^2$

(b)  $(-3)^2$

(c)  $\sqrt[3]{-125}$

(d)  $-\sqrt{81}$

(e)  $(-2)^3 \times (-2)^2$

**Q3.** Evaluate **without a calculator**:  $3 + 5 \times (-2) - 18 \div (-3)$ .

**Q4.** Evaluate **without a calculator**:

$$[(-4)^2 - \sqrt{64}] \div (-2) + 3.$$

**Q5.** A drone descends at 40 cm per second.

(a) Represent the change in height in 2 minutes by a negative number.

(b) If its height was 5000 cm, find its height after 2 minutes.

**Q6.** Water leaks from a tank at 250 ml per minute.

(a) Represent the amount leaked in 12 minutes by a negative number (in ml).

(b) The tank started with 5 L. How much remains after 12 minutes?

**Q7.** In a 12-question test: +5 marks per correct answer, -2 per wrong answer, -1 per unanswered.

(a) Lowest possible score.

(b) Score if a student gets 7 correct, 3 wrong, 2 unanswered.

**Q8.** Insert **one pair of brackets** to make each statement correct.

(a)  $5 + 4 \times 3 - 2 = 25$

(b)  $16 - 4 \div 2^2 = 3$

**Q9.** A student writes:  $(-3)^2 - 3^2 = -9 - 9 = -18$ .

(a) State the mistake.

(b) Give the correct answer.

**Q10.** Let  $x = -\frac{7}{3}$ .

(a) Write the reciprocal of  $x$ .

(b) Evaluate  $x \times$  (its reciprocal).

(c) Evaluate  $x \div \frac{14}{9}$ .

## 2.4 Rational numbers, irrational numbers and real numbers

**Rational numbers (numbers you can write as a fraction)**

A **rational number** can be written as  $\frac{a}{b}$  where  $a, b$  are integers and  $b \neq 0$ .

These include:

- Integers:  $-4, 0, 7$
- Fractions:  $\frac{2}{5}$
- Terminating decimals:  $0.72$
- Recurring decimals:  $0.\overline{3}$

### Examples

- $0.72 = \frac{72}{100} = \frac{18}{25}$
- $0.\overline{3} = \frac{1}{3}$

### Terminating vs recurring decimals

- **Terminating:** ends (like  $0.125$ )
- **Recurring:** repeats a pattern forever (like  $0.41\overline{6}$ )

### Irrational numbers (cannot be written as a fraction)

These cannot be written exactly as a fraction, and their decimals go on forever **without repeating**.

Common examples:

- $\pi$
- $\sqrt{2}, \sqrt{5}$  (when the number under the square root is *not* a perfect square)

### Examples

- $\sqrt{50} = \sqrt{25 \times 2} = 5\sqrt{2}$  is irrational because  $\sqrt{2}$  is irrational.
- $\sqrt{16} = 4$  is rational (perfect square).

### Real numbers

**Real numbers** include *all* rational and irrational numbers.

So:

- rational numbers  $\subset$  real numbers
- irrational numbers  $\subset$  real numbers

### Comparing surds (like $\sqrt{5}$ )

You can compare by knowing nearby squares:

- $2^2 = 4, 3^2 = 9$  so  $\sqrt{5}$  is between 2 and 3.
- Roughly,  $\sqrt{5} \approx 2.24$  (good enough for ordering).

### Practice Questions

**Q1.** Express as a fraction in simplest form (or mixed number where stated).

- $0.72$
- $-2.5$

- (c) 3.04  
 (d)  $5\frac{3}{8}$  as an improper fraction

**Q2.** Using a calculator, evaluate and state whether the decimal is **terminating** or **recurring**.

- (a)  $\frac{7}{16}$   
 (b)  $\frac{5}{12}$   
 (c)  $-\frac{11}{40}$   
 (d)  $\frac{13}{6}$

**Q3.** From  $\sqrt{50}$ , 3.1416,  $0.\overline{12}$ ,  $\frac{\pi}{3}$ ,  $-\sqrt{81}$ ,  $\frac{2}{7}$ ,  $\sqrt{\frac{1}{4}}$ :

- (a) List the rational numbers.  
 (b) List the irrational numbers.

**Q4.** Arrange in ascending order:  $-\sqrt{5}$ ,  $-2.1$ ,  $-2$ ,  $1.7$ ,  $\sqrt{3}$ .

**Q5.** Which of the following are **integers**?

$$-4, 0, \frac{5}{2}, \sqrt{16}, -\sqrt{7}, 3.0, -\frac{8}{4}$$

**Q6.** Express each as a fraction in simplest form.

- (a)  $0.\overline{3}$   
 (b)  $0.\overline{27}$   
 (c)  $1.\overline{2}$

**Q7.** (a) Express  $0.\overline{18}$  as a fraction in simplest form.

(b) Hence, find  $0.\overline{18} + 0.\overline{09}$  as a fraction in simplest form.

**Q8.** State whether each number is **rational** or **irrational**.

- (a)  $\frac{2\pi}{\pi}$   
 (b)  $\frac{\sqrt{12}}{\sqrt{3}}$   
 (c)  $\sqrt{2} + \sqrt{8}$   
 (d)  $\frac{\sqrt{50}}{5}$

**Q9.** (a) Between which two consecutive integers does  $\sqrt{2}$  lie? Give  $\sqrt{2}$  correct to 2 d.p.

(b) Between which two consecutive integers does  $\sqrt{7}$  lie? Give  $\sqrt{7}$  correct to 2 d.p.

**Q10.** Two students claim:

- A: "Every irrational number is a real number."
- B: "Every real number is rational."

Who is correct? Give a brief reason.



## 2.5 Operations on real numbers

This section is mainly about doing calculations correctly with:

- fractions
- mixed numbers
- decimals
- negatives
- and sometimes roots

### Converting between mixed and improper fractions

- Mixed number  $\rightarrow$  improper:

$$2\frac{5}{6} = \frac{2 \times 6 + 5}{6} = \frac{17}{6}$$

- Improper  $\rightarrow$  mixed: divide top by bottom.

$$\frac{17}{12} = 1\frac{5}{12}$$

### Simplifying fractions

Divide top and bottom by the same number.

- $\frac{18}{24}$  divide by 6  $\Rightarrow \frac{3}{4}$

### Adding/subtracting fractions

You need a **common denominator**.

- $\frac{3}{4} + \frac{2}{5} = \frac{15}{20} + \frac{8}{20} = \frac{23}{20}$

### Multiplying fractions

Multiply tops, multiply bottoms, then simplify.

- $\frac{3}{8} \times \frac{5}{6} = \frac{15}{48} = \frac{5}{16}$

### Dividing fractions

Flip the second fraction (take the reciprocal), then multiply.

- $\frac{3}{8} \div \left(\frac{1}{4}\right) = \frac{3}{8} \times 4 = \frac{3}{2}$

### Decimals with negatives

Treat them like signed numbers and combine carefully.

- $0.008 - (-1.02) = 0.008 + 1.02 = 1.028$

### “Halfway between two numbers”

Halfway means “average of the two numbers”:

$$\text{halfway} = \frac{\text{first} + \text{second}}{2}$$

Example:

$$\text{halfway between } -6 \text{ and } 5 = \frac{-6 + 5}{2} = -\frac{1}{2}$$

### Practice Questions

**Q1.** (a) Simplify  $\frac{18}{24}$ .

(b) Convert  $2\frac{5}{6}$  to an improper fraction.

(c) Convert  $\frac{17}{12}$  to a mixed number.

**Q2.** Evaluate **without a calculator**.

(a)  $\frac{3}{4} + \frac{2}{5}$

(b)  $\frac{5}{6} - \frac{1}{4}$

(c)  $1\frac{1}{3} + 2\frac{1}{6}$

(d)  $-\frac{3}{5} + \frac{7}{10}$

**Q3.** Evaluate **without a calculator**.

(a)  $\frac{3}{8} \div \left(-\frac{1}{4}\right)$

(b)  $-2\frac{1}{2} \times \frac{3}{5}$

(c)  $\frac{5}{12} \div \frac{8}{9}$

**Q4.** Evaluate **without a calculator**:

$$\frac{3}{5} - \left[ \frac{1}{2} \div \left( -\frac{3}{4} \right) \right].$$

**Q5.** Evaluate **without a calculator**.

(a)  $1.25 + 0.81 - 5.4$

(b)  $0.008 - (-1.02)$

(c)  $(-3.18) - 5.4 + 0.02$

(d)  $6.7 + (-10.3) + 4$

**Q6.** Find the number exactly halfway between  $-0.625$  and  $-\frac{1}{3}$ . Express your answer as a fraction in simplest form.

**Q7.** In a class,  $\frac{1}{3}$  walk to school,  $\frac{2}{5}$  take the bus, and the rest take MRT. Find the fraction who take MRT.

**Q8.** Joanne spent  $\frac{1}{4}$  of her money on food,  $\frac{3}{10}$  on transport, and donated  $\frac{2}{5}$  of the remainder. She saved the rest.

(a) Find the fraction of her money that was donated.

(b) If she saved \$216, find her total money and the amount spent on food.

**Q9.** An empty file has mass 250 g. When filled with papers, the mass is 0.73 kg. If it can hold 240 sheets, find the mass of 1 sheet (in grams).

**Q10.** Using a calculator, evaluate (give answers correct to **5 significant figures**).

(a)  $\sqrt[3]{19} \times \pi^2$

(b)  $\sqrt[5]{3.75 + 0.006}$

# Solutions

## 2.1 Concept of negative numbers and the number line

- Q1.** (a)  $-6$  (b)  $+4$  (c)  $-2$  (below is negative; rising is positive)
- Q2.** (a)  $-3.5 < -1 < 0.6 < 2.4 < \frac{5}{2}$  (b)  $>$  (c)  $0.6$
- Q3.** (a)  $A$  (b)  $<$  (c)  $0$  or  $1$
- Q4.** (a)  $<$  (b)  $<$  (c)  $>$  (d)  $=$
- Q5.** (a)  $5^\circ\text{C}$  below average (b) A  $25^\circ\text{C}$ , B  $29^\circ\text{C}$ , C  $23^\circ\text{C}$  (c)  $C, A, B$
- Q6.** (a)  $+120$  (b) loss  $\$45$  (c)  $120 - 45 + 80 - 150 = +5$
- Q7.** (a)  $350, 230, 260, 180$  m (b)  $350$  m (c) No
- Q8.** (a)  $-12, -8, -4$  (b)  $3, 6$
- Q9.**  $\pm 4, \pm 6, \pm 8, \pm 9, \pm 10, \pm 12, \pm 14$
- Q10.** (a)  $7, 4, 8, -2$  (b)  $-2$  (c) after the  $-10$  event

## 2.2 Addition and subtraction of integers

- Q1.** (a)  $7$  (b)  $23$  (c)  $-20$  (d)  $-18$
- Q2.** (a)  $-6$  (b)  $0$  (c)  $-15$  (d)  $16$
- Q3.** (a)  $x = -8$  (b)  $y = 15$  (c)  $z = -13$  (d)  $a = -10$
- Q4.** (a) start  $-250$ , dive  $-120$ , rise  $+310$  (b)  $-60$  (c)  $60$  m below sea level
- Q5.** (a)  $-4 + 9 - 12 = -7^\circ\text{C}$  (b)  $7^\circ\text{C}$  below  $0$
- Q6.** (a) A  $-430$ , B  $+4095$ , C  $+1800$  (b)  $4525$  m (c)  $2230$  m
- Q7.** (a) Jan (b) loss  $\$400$  (c)  $350 - 120 + 80 - 400 = -90$  (overall loss  $\$90$ )
- Q8.** (a)  $8 - (-5) = 13$  h (b)  $15:00 + 13 = 04:00$  (next day) (c)  $09:30 - 13 = 20:30$  (previous day)
- Q9.** Sum  $= -4$ , average  $= -1$
- Q10.**  $x = \frac{-6 + 5}{2} = -\frac{1}{2}$

## 2.3 Multiplication, division and combined operations of integers

- Q1.** (a)  $-42$  (b)  $8$  (c)  $-5$  (d)  $30$
- Q2.** (a)  $-9$  (b)  $9$  (c)  $-5$  (d)  $-9$  (e)  $-32$
- Q3.**  $3 + 5(-2) - 18/(-3) = 3 - 10 + 6 = -1$
- Q4.**  $(16 - 8)/(-2) + 3 = -4 + 3 = -1$

**Q5.** (a)  $-40 \times 120 = -4800$  cm (b)  $5000 - 4800 = 200$  cm

**Q6.** (a)  $-250 \times 12 = -3000$  ml (b)  $5000 - 3000 = 2000$  ml  $= 2$  L

**Q7.** (a)  $12(-2) = -24$  (b)  $7(5) - 3(2) - 2(1) = 27$

**Q8.** (a)  $(5 + 4) \times 3 - 2 = 25$  (b)  $(16 - 4) \div 2^2 = 3$

**Q9.** (a)  $(-3)^2$  is  $+9$ , not  $-9$  (b)  $9 - 9 = 0$

**Q10.** (a)  $-\frac{3}{7}$  (b)  $1$  (c)  $-\frac{7}{3} \times \frac{9}{14} = -\frac{3}{2}$

## 2.4 Rational numbers, irrational numbers and real numbers

- Q1.** (a)  $\frac{18}{25}$  (b)  $-\frac{5}{2}$  (c)  $\frac{76}{25}$  (d)  $\frac{43}{8}$
- Q2.** (a)  $0.4375$  terminating (b)  $0.41\overline{6}$  recurring (c)  $-0.275$  terminating (d)  $2.1\overline{6}$  recurring
- Q3.** (a)  $3.1416, 0.\overline{12}, -9, \frac{2}{7}, \frac{1}{2}$  (b)  $\sqrt{50}, \frac{\pi}{3}$
- Q4.**  $-\sqrt{5} < -2.1 < -2 < 1.7 < \sqrt{3}$  ( $\sqrt{5} \approx 2.236, \sqrt{3} \approx 1.732$ )
- Q5.** Integers:  $-4, 0, 4, 3, -2$
- Q6.** (a)  $\frac{1}{3}$  (b)  $\frac{3}{11}$  (c)  $\frac{11}{9}$
- Q7.** (a)  $\frac{2}{11}$  (b)  $\frac{2}{11} + \frac{1}{11} = \frac{3}{11}$
- Q8.** (a) rational ( $=2$ ) (b) rational ( $=2$ ) (c) irrational ( $=3\sqrt{2}$ ) (d) irrational ( $=\sqrt{2}$ )
- Q9.** (a) between  $1$  and  $2$ ;  $\sqrt{2} \approx 1.41$  (b) between  $2$  and  $3$ ;  $\sqrt{7} \approx 2.65$
- Q10.** A correct; B false (e.g.  $\sqrt{2}$  is real but irrational)

## 2.5 Operations on real numbers

- Q1.** (a)  $\frac{3}{4}$  (b)  $\frac{17}{6}$  (c)  $1\frac{5}{12}$
- Q2.** (a)  $\frac{23}{20} = 1\frac{3}{20}$  (b)  $\frac{7}{12}$  (c)  $\frac{7}{2} = 3\frac{1}{2}$  (d)  $\frac{1}{10}$
- Q3.** (a)  $-\frac{3}{2}$  (b)  $-\frac{3}{2}$  (c)  $\frac{15}{32}$
- Q4.**  $\frac{1}{2} \div (-\frac{3}{4}) = -\frac{2}{3}$  so  $\frac{3}{5} - (-\frac{2}{3}) = \frac{19}{15} = 1\frac{4}{15}$
- Q5.** (a)  $-3.34$  (b)  $1.028$  (c)  $-8.56$  (d)  $0.4$
- Q6.**  $-\frac{23}{48}$
- Q7.**  $1 - (\frac{1}{3} + \frac{2}{5}) = \frac{4}{15}$
- Q8.** (a) remainder  $= \frac{9}{20}$ , donated  $= \frac{2}{5} \cdot \frac{9}{20} = \frac{9}{100}$  (b) saved fraction  $= \frac{27}{100}$  so total  $= 216 \div \frac{27}{100} = 800$ , food  $= \frac{1}{4} \cdot 800 = 200$
- Q9.**  $0.73$  kg  $= 730$  g, papers  $= 730 - 250 = 480$  g, per sheet  $= 480/240 = 2$  g
- Q10.** (a)  $\approx 26.336$  (b)  $\approx 1.3030$