IMA Journal of Complex Networks (2026) Page 1 of 19
doi:10.1093/comnet/xxx000

A Directed-Hypergraph Representation of Higher-Order Information Flows

YUNJU LEE*
Independent Researcher, Seoul, Republic of Korea
*Corresponding author: eka7268de@aol.com

[Received on 11 November 2026]

A statistical approach to higher-order interactions requires an understanding of higher-order correlations among components. The information dynamics of these correlations can be encoded using multiple transfer entropy and multiple reversed transfer entropy. This paper investigates the relationship between the topology of higher-order information flows and the entropy dynamics of individual components. Because a multiple transfer entropy has a source set and a target set, its structure can be naturally represented by a hyperarc. By distinguishing statistically direct and indirect relationships, we introduce an information flow hypergraph and analyze its BF-hypergraph substructure in terms of BF-sinks, BF-sources, and weak BF-hypercycles. Under the strong blocking flows condition and $\alpha_1 = 0$, periodic or recurrent entropy dynamics with a fixed nonempty flow topology requires the presence of a weak BF-hypercycle. We further show that stationarity imposes an additional local balance condition between B-inflows and F-outflows. The corresponding result for cycles in classical directed graph theory is recovered as a special case.

Keywords: mutual information; transfer entropy; information dynamics; higher-order correlation; complex network; directed hypergraphs

1. Introduction

Synergistic effects are key characteristics of higher-order dynamics [1–7] and distinguish many-body interactions from purely pairwise interactions. For a statistical characterization of such dynamics, several studies [8–11] have adopted information-theoretic approaches. In particular, partial information decomposition (PID) [11, 12] provides nonnegative measures of synergistic and redundant information, although their definitions are not unique. In this paper, we instead employ multiple mutual information, also referred to as interaction information or co-information [8, 13–19]. Although it is a signed quantity, multiple mutual information is uniquely defined consistently for arbitrary orders. It therefore captures higher-order correlation structures that cannot be reduced to ordinary pairwise mutual information [8, 9].

The temporal evolution of such higher-order correlations can be described using multiple transfer entropy and multiple reversed transfer entropy [20]. These quantities extend the decomposition of mutual-information dynamics to higher orders, capturing directional higher-order information flows. However, their topological organization has not yet been systematically characterized.

This paper introduces an information flow hypergraph to represent the topology of direct higher-order information flows. A multiple transfer entropy $T_{A\to B}$ contains a source set and a target set, which naturally correspond to the tail and head of a hyperarc [21–23]. Since nonzero transfer entropy may

© The author 2026. Published by Oxford University Press on behalf of the Institute of Mathematics and its Applications. All rights reserved.

also reflect indirect statistical dependencies, direct information flows are identified using dynamical neighbors. The resulting representation consists of hyperarcs corresponding to direct multiple transfer entropies and multiple reversed transfer entropies.

The entropy dynamics of an individual vertex is governed by a BF-hypergraph substructure composed of B-arcs and F-arcs [21, 22, 24]. Under the strong blocking flows condition and deterministic reversible dynamics, this structure constrains the possible evolution of the vertex entropy. In particular, a finite oriented BF-hypergraph without weak BF-hypercycles possesses a BF-sink and a BF-source. At a BF-sink, the entropy increases strictly, whereas at a BF-source it decreases strictly. This result generalizes the pairwise argument developed in Ref. [25].

A weak BF-hypercycle is therefore necessary for periodic or recurrent entropy dynamics when the nonempty flow topology is fixed and the standing assumptions hold at every step. However, unlike a cycle in classical directed graph theory [26, 27], a weak BF-hypercycle does not necessarily admit a stationary state. This paper derives an additional necessary condition for stationarity: every non-isolated vertex must have both a B-inflow and an F-outflow. Their aggregate contributions must also balance numerically.

The remainder of this paper is organized as follows. Section 2 reviews the definitions of higher-order information dynamics and directed hypergraphs. Section 3 introduces the information flow hypergraph and the blocking flows conditions. Section 4 analyzes BF-hypergraphs without weak BF-hypercycles. Section 5 derives a necessary condition for stationary information dynamics on a weak BF-hypercycle.

2. Definitions

2.1 *Higher-Order Information Dynamics*

Consider an N-component system labeled with a finite set V. The state of $v \in V$ at time t is denoted as $\Gamma_v(t)$ taking values in a discrete space. An ensemble description or stochastic dynamics of the entire system induces a joint probability. Information-theoretic quantities can then be used to characterize statistical correlations and directional information flows among the components. All entropies used below are finite, and logarithms are natural. The symbol $\subset$ allows equality, and an empty subset sum is zero.

DEFINITION 2.1 For $v \in V$, the **Shannon entropy** H_v [28, 29] is defined as:

$$H_v := -\sum_{\gamma_v \in \mathscr{M}} \mathbb{P}(\Gamma_v = \gamma_v) \log \mathbb{P}(\Gamma_v = \gamma_v). \tag{2.1}$$

And we also define Shannon entropy of $A := \{a_1, \ldots, a_k\} \subset V$ as:

$$H_A := -\sum_{\gamma_{a_1}, \ldots, \gamma_{a_k} \in \mathscr{M}} \mathbb{P}(\Gamma_{a_1} = \gamma_{a_1}, \ldots, \Gamma_{a_k} = \gamma_{a_k}) \log \mathbb{P}(\Gamma_{a_1} = \gamma_{a_1}, \ldots, \Gamma_{a_k} = \gamma_{a_k}). \tag{2.2}$$

Then, the **conditional entropy** is defined as: for $v, u \in V$,

$$H_u|_v := H_{\{v,u\}} - H_v, \tag{2.3}$$

and, for $A, B \subset V$,

$$H_B|_A := H_{A\cup B} - H_A. \tag{2.4}$$

Here, Γ_v, Γ_u, and $\Gamma_{a_1}, \ldots, \Gamma_{a_k}$ are discrete random variables in a state space $\mathscr{M}$.

DEFINITION 2.2 For $v, u \in V$, the **mutual information** [29] between Γ_v and Γ_u quantifies the reduction in the uncertainty of Γ_u obtained by conditioning on Γ_v:

$$I_{v:u} := H_u - H_u|_v = H_v + H_u - H_{\{v,u\}}. \tag{2.5}$$

The **conditional mutual information** is defined as

$$I_{v:u}|_w := H_v|_w + H_u|_w - H_{\{v,u\}}|_w, \tag{2.6}$$

with $w \in V$.

We also use the notation $I_{(A):v'}|_v := H_{\{v'\}}|_{\{v\}} - H_{\{v'\}}|_{\{v\}\cup A}$ for $v \in V, A \subset V$.

For discrete random variables, the Shannon entropy, conditional entropy, and mutual information are nonnegative.

DEFINITION 2.3 For $A \subset V$, the **multiple mutual information (co-information)** [13, 14, 16] is a signed generalization of mutual information that characterizes higher-order statistical dependence.

$$I_A^{\mathrm{co}} := \sum_{r=1}^{|A|} (-1)^{r+1} \sum_{S\subset A, |S|=r} H_S. \tag{2.7}$$

And the **conditional multiple mutual information** is defined as,

$$I_A^{\mathrm{co}}|_B := \sum_{r=1}^{|A|} (-1)^{r+1} \sum_{S\subset A, |S|=r} H_S|_B, \tag{2.8}$$

We note that the cases where $|A| = 2$ corresponds to the mutual information, and $|A| = 1$ for Shannon entropy.

The temporal evolution of correlations can be related to directional information flows [20]. In an information-theoretic framework, such directional dependencies are characterized using transfer entropy [30].

DEFINITION 2.4 In the one-step formulation adopted here, the **transfer entropy** [31, 32] from $u \in V$ to $v \in V$ measures a directional predictive dependence of v on u:

$$T_{u\to v} := I_{v':u}|_v, \tag{2.9}$$

where unprimed and primed variables are evaluated at times t and $t + \Delta t$, respectively, with $\Delta t > 0$.

DEFINITION 2.5 **Reversed transfer entropy** [33–35], also referred to as backward transfer entropy [33], is defined by reversing the temporal indexing in the transfer-entropy expression:

$$T^{\dagger}_{u\to v} := I_{v:u'}|_{v'}. \tag{2.10}$$

DEFINITION 2.6 **Multiple transfer entropy** and **multiple reversed transfer entropy** [20] are higher-order generalizations of transfer entropy and reversed transfer entropy:

$$\begin{aligned} T_{B\to A} &:= I^{\mathrm{co}}_{A'\cup B}|_{A} \\ T^{\dagger}_{B\to A} &:= I^{\mathrm{co}}_{A\cup B'}|_{A'}, \end{aligned} \tag{2.11}$$

for nonempty disjoint $A, B \subset V$.

We also note that multiple transfer entropy and multiple reversed transfer entropy become the transfer entropy and reversed transfer entropy when $|A| = |B| = 1$.

Using these quantities, Ref. [20] derived exact decompositions of the temporal changes in entropy with transfer entropy and reversed transfer entropy:

$$\Delta H_v = \sum_{\eta\in K_v} [T_{\eta\to v} - T^{\dagger}_{\eta\to v}] - E_{K_v\to v} + \alpha_1^{K_v\to v}, \tag{2.12}$$

where,

$$\begin{aligned} E_{K_v\to v} &:= \sum_{u=2}^{|K_v|} (-1)^u \sum_{S\subset K_v, |S|=u} [T_{S\to\{v\}} - T^{\dagger}_{S\to\{v\}}] \\ \alpha_1^{K_v\to v} &:= H_{\{v'\}}|_{\{v\}\cup K_v} - H_{\{v\}}|_{\{v'\}\cup K'_v}. \end{aligned} \tag{2.13}$$

In this identity, K_v may be any subset of $V\setminus\{v\}$. The term α_1 represents residual contributions that are not expressed as transfer entropy or reversed transfer entropy. Its roles under deterministic reversible dynamics will be discussed in Section 3.2.

2.2 *Directed Hypergraphs*

We summarize the definitions from directed-hypergraph theory that are required for the subsequent analysis [21–23, 36].

DEFINITION 2.7 A **directed hypergraph** $(V, \overrightarrow{E})$ consists of a finite set V of vertices and a nonempty set $\overrightarrow{E} \subset \{(e^+, e^-) | e^+, e^- \subset V \text{ and } e^+ \neq \emptyset, e^- \neq \emptyset, e^+ \cap e^- = \emptyset\}$ of hyperarcs. The set of reversed hyperarcs is defined as $\overleftarrow{E} := \{(e^-, e^+) | (e^+, e^-) \in \overrightarrow{E}\}$. The directed hypergraph $(V, \overleftarrow{E})$ is called the **symmetric image** of $(V, \overrightarrow{E})$. For a hyperarc $\overrightarrow{e} = (e^+, e^-) \in \overrightarrow{E}$, the sets e^+ and e^- are called its tail and head, respectively, and are denoted by $\mathscr{T}(\overrightarrow{e})$ and $\mathscr{H}(\overrightarrow{e})$.

We assume that a directed hypergraph $(V, \overrightarrow{E})$ has no isolated vertices, namely,

$$V = \bigcup_{\overrightarrow{e}\in\overrightarrow{E}} [\mathscr{T}(\overrightarrow{e}) \cup \mathscr{H}(\overrightarrow{e})]. \tag{2.14}$$

DEFINITION 2.8 An **oriented hypergraph** $(V, \overrightarrow{E})$ is a directed hypergraph $(V, \overrightarrow{E})$ satisfying $\overrightarrow{E} \cap \overleftarrow{E} = \emptyset$.

DEFINITION 2.9 A **BF-hypergraph** $(V, \overrightarrow{E})$ is a directed hypergraph $(V, \overrightarrow{E})$ such that $|e^+| = 1$ or $|e^-| = 1$ for every $(e^+, e^-) \in \overrightarrow{E}$, and $V = \bigcup_{\overrightarrow{e} \in \overrightarrow{E}} [\mathscr{T}(\overrightarrow{e}) \cup \mathscr{H}(\overrightarrow{e})]$. Specifically, the hyperarc $\overrightarrow{e_B} = (e^+, e^-) \in \overrightarrow{E}$ with $|e^-| = 1$ is called the B-arc, while the hyperarc $\overrightarrow{e_F} = (e^+, e^-) \in \overrightarrow{E}$ with $|e^+| = 1$ is called the F-arc. An ordinary directed edge in a directed graph satisfies both conditions and can therefore be regarded as both a B-arc and an F-arc.

DEFINITION 2.10 For a vertex $v \in V$ in a BF-hypergraph $(V, \overrightarrow{E})$, the **B-inflow** set of v, denoted by $\mathrm{In}_B(v)$, is the set of B-arcs directed toward v: $\mathrm{In}_B(v) := \{(e^+, e^-) \in \overrightarrow{E} \,|\, e^- = \{v\}\}$. Similarly, the **B-outflow** set of v is defined as $\mathrm{Out}_B(v) := \{(e^+, e^-) \in \overrightarrow{E} \,|\, v \in e^+ \text{ and } |e^-| = 1\}$. The definitions for **F-inflow** and **F-outflow** are $\mathrm{In}_F(v) := \{(e^+, e^-) \in \overrightarrow{E} \,|\, v \in e^- \text{ and } |e^+| = 1\}$ and $\mathrm{Out}_F(v) := \{(e^+, e^-) \in \overrightarrow{E} \,|\, e^+ = \{v\}\}$.

DEFINITION 2.11 A **BF-sink** is a vertex $v \in V$ such that $\mathrm{Out}_B(v) = \emptyset$, $\mathrm{Out}_F(v) = \emptyset$, and $\mathrm{In}_B(v) \cup \mathrm{In}_F(v) \neq \emptyset$. A **BF-source** is a vertex $v \in V$ such that $\mathrm{In}_B(v) = \emptyset$, $\mathrm{In}_F(v) = \emptyset$, and $\mathrm{Out}_B(v) \cup \mathrm{Out}_F(v) \neq \emptyset$.

DEFINITION 2.12 In a directed hypergraph $(V, \overrightarrow{E})$, a **weak hyperpath** from v_1 to v_{n+1} is defined as a $(2n+1)$-tuple $(v_1, \overrightarrow{e_1}, v_2, \overrightarrow{e_2}, \ldots, \overrightarrow{e_n}, v_{n+1})$ such that $v_i \in \mathscr{T}(\overrightarrow{e_i})$ and $v_{i+1} \in \mathscr{H}(\overrightarrow{e_i})$ with $\overrightarrow{e_i} \in \overrightarrow{E}$ for $\forall i \in \{1, 2, \ldots, n\}$. A vertex $s \in V$ is **connected** to $u \in V$ if a weak hyperpath from u to s exists. Furthermore, a nonempty subset $S \subset V$ is **connected** to the nonempty subset $U \subset V$ if there are $u \in U$ and $s \in S$ such that s is connected to u.

DEFINITION 2.13 The **transitive closure** of a directed hypergraph $\mathscr{G} = (V, \overrightarrow{E})$ is a directed hypergraph $\mathscr{G}^* = (V, \overrightarrow{E^*})$ such that, for any nonempty disjoint subsets $U, S \subset V$, $(U, S) \in \overrightarrow{E^*}$ if and only if S is connected to U in $\mathscr{G}$.

DEFINITION 2.14 A **weak BF-hypercycle** of length n in a BF-hypergraph $(V, \overrightarrow{E}_{BF})$ is a weak hyperpath $(v_1, \overrightarrow{e_1}, v_2, \overrightarrow{e_2}, \ldots, \overrightarrow{e_n}, v_{n+1})$ in $(V, \overrightarrow{E}_{BF})$ such that $v_1 = v_{n+1}$.

In the definition of weak BF-hypercycle, no distinctness condition is imposed on the intermediate vertices or hyperarcs.

3. Identification of Higher-Order Information Flows

3.1 *Information Flow Hypergraph*

Consider an N-component system with a joint probability distribution. Let $V = \{v_1, \ldots, v_N\}$ be the set of components. The set of all nonzero multiple transfer entropies is defined as,

$$\overrightarrow{E}_{total} := \{(A, B) | A, B \subset V \text{ are nonempty and disjoint, and } T_{A \to B} \neq 0\}. \tag{3.1}$$

However, $\overrightarrow{E}_{total}$ may contain indirect statistical relationships. For example, $T_{p\to q}$ can be nonzero even when the influence of p reaches q through intermediate components. Therefore, we effectively consider direct causal relationships using dynamical neighbors.

DEFINITION 3.1 A set of **dynamical neighbors** $K_A \subset V \setminus A$ of a nonempty $A \subset V$ is an inclusion-minimal set satisfying,

$$H_{A'}|_{A\cup K_A} = 0, \tag{3.2}$$

where the primed variables are evaluated at the next time step. Similarly, a set of **backward dynamical neighbors** $Q_A \subset V \setminus A$ is an inclusion-minimal set satisfying,

$$H_A|_{A'\cup Q'_A} = 0. \tag{3.3}$$

Here, Q'_A denotes the future variables corresponding to the vertex labels in Q_A.

We denote $K_v := K_{\{v\}}$ and $Q_v := Q_{\{v\}}$ for the singleton case.

To ensure that the dynamical neighbors are well defined, we assume the dynamics of N components is described as: for $F : \mathscr{M}^N \times \mathbb{R} \to \mathscr{M}^N$,

$$(\Gamma_{v_1}(t+\Delta t),\ldots,\Gamma_{v_N}(t+\Delta t)) = F(\Gamma_{v_1}(t),\ldots,\Gamma_{v_N}(t),t), \tag{3.4}$$

where Γ_v is the state of component $v \in V$. For the subsequent entropy results, we also assume that the full state can be reconstructed from its future state:

$$(\Gamma_{v_1}(t),\ldots,\Gamma_{v_N}(t)) = \widetilde{F}(\Gamma_{v_1}(t+\Delta t),\ldots,\Gamma_{v_N}(t+\Delta t),t+\Delta t). \tag{3.5}$$

Thus, $V \setminus A$ satisfies each reconstruction condition. Since V is finite, inclusion-minimal sets K_A and Q_A exist. Stochastic input examples in Appendix C are used only to illustrate local flow quantities.
However, the sets of dynamical neighbors are not necessarily unique. We therefore define,

$$\begin{aligned} \overline{\mathscr{K}}_A &:= \{K \subset V \setminus A | K \text{ is a set of dynamical neighbors of } A\}, \\ \overline{\mathscr{Q}}_A &:= \{Q \subset V \setminus A | Q \text{ is a set of backward dynamical neighbors of } A\}. \end{aligned} \tag{3.6}$$

Then, the set of direct multiple transfer entropies is defined as $\overrightarrow{E}_T := \{(A,B) \in \overrightarrow{E}_{total} | \exists K_B \in \overline{\mathscr{K}}_B \text{ such that } A \subset K_B\}$. The set $\overrightarrow{E}_T$ contains the forward terms used in the entropy expansion in Sec. 3.2. However, Proposition 3.2 shows that the connectivity of $(V, \overrightarrow{E}_{total})$ is not generally preserved in $(V, \overrightarrow{E}_T)$. Thus, some indirect statistical relationships cannot be represented as chains of direct information flows.

PROPOSITION 3.2 In general, $\mathscr{G}^*_T \neq \mathscr{G}^*_{total}$, where $\mathscr{G}_T = (V, \overrightarrow{E}_T)$, $\mathscr{G}_{total} = (V, \overrightarrow{E}_{total})$.

(proof) Appendix A.

However, if one modifies the definition of $\overrightarrow{E}_T$ as $\overrightarrow{E}_{\bar{T}} := \{(A,B) \in \overrightarrow{E}_{total} | \exists K_B \in \overline{\mathscr{K}}_B \text{ such that } A \cap K_B \neq \emptyset\}$, the specific example for Proposition 3.2, addressed in Appendix A, loses its meaning: X and Y

are connected in $\mathscr{G}_{\bar{T}} = (V, \overrightarrow{E}_{\bar{T}})$, as the redundant information flow exists. Specifically, $T_{\{Y,C\}\to\{X\}} = \ln 2 - h(\varepsilon) > 0$, indicating $(\{Y,C\},\{X\}) \in \overrightarrow{E}_{\bar{T}}$. Moreover, Proposition 3.3 shows that $\mathscr{G}^*_{\bar{T}} = \mathscr{G}^*_{total}$.

PROPOSITION 3.3 $\mathscr{G}^*_{\bar{T}} = \mathscr{G}^*_{total}$, where $\mathscr{G}_{\bar{T}} = (V, \overrightarrow{E}_{\bar{T}})$, $\mathscr{G}_{total} = (V, \overrightarrow{E}_{total})$.

(proof) Appendix B.

Although $\overrightarrow{E}_{\bar{T}}$ preserves the connectivity of $\overrightarrow{E}_{total}$, the subsequent analysis uses $\overrightarrow{E}_{T}$, which more strictly identifies direct information flows whose source sets are contained in a set of dynamical neighbors.

DEFINITION 3.4 A **transfer-entropy hypergraph** $\mathscr{G}_T$ of an N-component system is a directed hypergraph $(V, \overrightarrow{E}_T)$, where V is a set of N components and $\overrightarrow{E}_T$ is the set of direct multiple transfer entropies,

$$\overrightarrow{E}_T := \{(A,B) \in \overrightarrow{E}_{total} | \exists K_B \in \overline{\mathscr{K}}_B \text{ such that } A \subset K_B\}. \tag{3.7}$$

Here, a **reversed-transfer-entropy hypergraph** $\mathscr{G}_{T^\dagger}$ is a directed hypergraph $(V, \overrightarrow{E}_{T^\dagger})$, where $\overrightarrow{E}_{T^\dagger}$ is defined as,

$$\begin{aligned}\overrightarrow{E}_{T^\dagger} := \{(A,B) \mid A,B \subset V \text{ are nonempty and disjoint, } T^\dagger_{A\to B} \neq 0,\\ \exists Q_B \in \overline{\mathscr{Q}}_B \text{ such that } A \subset Q_B\}.\end{aligned} \tag{3.8}$$

We note that $\mathscr{G}_T$ and $\mathscr{G}_{T^\dagger}$ use the dynamical neighbors for their respective temporal directions. This ensures that the forward and backward terms in the entropy expansion below are represented. It does not assert completeness for the dynamics of all multiple mutual informations.

DEFINITION 3.5 The **information flow hypergraph** of an N-component system is the ordered pair $\mathscr{G}_{\Delta I} := (\mathscr{G}_T, \mathscr{G}_{T^\dagger})$, where $\mathscr{G}_T$ and $\mathscr{G}_{T^\dagger}$ are the transfer-entropy and reversed-transfer-entropy hypergraphs, respectively.

3.2 *Condition for* $\alpha_1 = 0$

The subsequent analysis uses the combined set $K_v^{\text{both}} := K_v \cup Q_v$, where K_v and Q_v are sets of dynamical neighbors and backward dynamical neighbors, respectively. By their definitions,

$$H_{\{v'\}}|_{\{v\}\cup K_v^{\text{both}}} = 0, \qquad H_{\{v\}}|_{\{v'\}\cup (K_v^{\text{both}})'} = 0. \tag{3.9}$$

Therefore,

$$\alpha_1^{K_v^{\text{both}}\to v} = 0. \tag{3.10}$$

The combined set need not be inclusion-minimal, and the smaller set K_v need not satisfy $\alpha_1^{K_v\to v} = 0$. To retain the direct-flow definitions, we expand the two temporal directions separately:

$$\begin{aligned}\Delta H_v &= I_{(K_v):v'}|_v - I_{(Q'_v):v}|_{v'} \\ &= \sum_{\emptyset\neq S\subset K_v} (-1)^{|S|+1} T_{S\to\{v\}} - \sum_{\emptyset\neq R\subset Q_v} (-1)^{|R|+1} T^\dagger_{R\to\{v\}}.\end{aligned} \tag{3.11}$$

Here, $I_{(K_v):v'}|_v = H_{v'}|_v$ and $I_{(Q'_v):v}|_{v'} = H_v|_{v'}$ are nonnegative. Every nonzero term in each sum is represented in the corresponding flow hypergraph.

3.3 *Blocking Flows Conditions*

To exclude a trivial cycle formed by simultaneously allowing a hyperarc and its reverse, $v_1 \to (e^+, e^-) \to v_2 \to (e^-, e^+) \to v_1$, we impose the blocking flows condition introduced in Ref. [20]. This condition assigns a fixed direction to each information-flow channel.

DEFINITION 3.6 An information flow hypergraph with **the blocking flows condition** is an information flow hypergraph $\mathscr{G}_{\Delta I} = (\mathscr{G}_T, \mathscr{G}_{T^\dagger})$ satisfying $\overrightarrow{E}_T \cap \overleftarrow{E}_T = \emptyset$, $\overrightarrow{E}_{T^\dagger} \cap \overleftarrow{E}_{T^\dagger} = \emptyset$, and $\overrightarrow{E}_T \cap \overrightarrow{E}_{T^\dagger} = \emptyset$, where $\mathscr{G}_T = (V, \overrightarrow{E}_T)$ and $\mathscr{G}_{T^\dagger} = (V, \overrightarrow{E}_{T^\dagger})$.

$\mathscr{G}_T$ and $\mathscr{G}_{T^\dagger}$ in $\mathscr{G}_{\Delta I}$ with the blocking flows condition are oriented hypergraphs. Here, the blocking flows condition is a restriction on the chosen dynamics and ensemble. Introducing separate intermediate channels can be a modeling choice, but does not automatically preserve the original one-step dynamics.

DEFINITION 3.7 An information flow hypergraph with **the strong blocking flows condition** is an information flow hypergraph $\mathscr{G}_{\Delta I} = (\mathscr{G}_T, \mathscr{G}_{T^\dagger})$ with the blocking flows condition satisfying $\overrightarrow{E}_{T^\dagger} = \overleftarrow{E}_T$, where $\mathscr{G}_T = (V, \overrightarrow{E}_T)$ and $\mathscr{G}_{T^\dagger} = (V, \overrightarrow{E}_{T^\dagger})$.

Then, under the strong blocking flows condition, the hypergraph $\mathscr{G}_{T^\dagger}$ becomes the symmetric image of $\mathscr{G}_T$. This is an additional assumption matching the presence or absence of hyperarcs, rather than their numerical values. It does not follow from stationarity or an aggregate entropy balance. The backward-neighbor definition does not require mutual forward causal parenthood.

Here, we illustrate how reversed transfer entropy in the inverse direction occurs. Consider three vertices v, u, w with independent fair binary states and the closed copying dynamics,

$$\Gamma_v(t+\Delta t) = \Gamma_u(t), \qquad \Gamma_u(t+\Delta t) = \Gamma_w(t), \qquad \Gamma_w(t+\Delta t) = \Gamma_v(t). \tag{3.12}$$

This is a deterministic reversible dynamics with an invariant independent uniform distribution. Then, $K_v = \{u\}$, $Q_v = \{w\}$, and the other neighbor sets follow cyclically. Its only nonzero direct transfer entropies are $T_{u\to v} = T_{v\to w} = T_{w\to u} = \ln 2$, and the reversed transfer entropies have the opposite directions and the same values. Thus, the strong blocking flows condition and $\alpha_1^{K_x^{\mathrm{both}}\to x} = 0$ hold for every vertex x. However, $\alpha_1^{K_v\to v} = -\ln 2$, showing why the backward neighbors are needed.

The strong blocking flows condition restricts the applicable class of dynamics. For example, let u, v be independent fair bits with $\Gamma_u(t+\Delta t) = \Gamma_u(t)$ and $\Gamma_v(t+\Delta t) = \Gamma_v(t) \oplus \Gamma_u(t)$. This dynamics is reversible, but $T_{u\to v} = T^\dagger_{u\to v} = \ln 2$, violating the blocking flows condition. More generally, a copying channel $v' = u$ has $T^\dagger_{v\to u} = H_u|_{u'}$; this vanishes if u' alone determines u. Appendix C provides local stochastic illustrations without the $\alpha_1 = 0$ condition.

4. Higher-Order Information Dynamics without Weak BF-hypercycles

As described in Eq. (3.11), the dynamics of individual entropy H_v is determined by higher-order information flows, $T_{S\to\{v\}}$ for $S\subset K_v$ and $T^{\dagger}_{R\to\{v\}}$ for $R\subset Q_v$. We fix $K_v\in\overline{\mathscr{K}}_v$ and $Q_v\in\overline{\mathscr{Q}}_v$ in Eq. (3.11). From the strong blocking flows condition, a nonzero $T^{\dagger}_{R\to\{v\}}$ implies the nonzero $T_{\{v\}\to R}$. We therefore focus on the hyperarcs associated with nonzero $T_{S\to\{v\}}$ and $T_{\{v\}\to R}$ in $\mathscr{G}_T=(V,\overrightarrow{E}_T)$ for an information flow hypergraph $\mathscr{G}_{\Delta I}=(\mathscr{G}_T,\mathscr{G}_{T^\dagger})$. All B- and F-arcs in $\mathscr{G}_T$ form a BF-hypergraph substructure $\mathscr{G}_{BF}=(V_{BF},\overrightarrow{E}_{BF})$ of $\mathscr{G}_T$. Here, V_{BF} is the set of vertices incident to these arcs; Proposition 4.2 below gives $V_{BF}=V$ under the standing assumptions. To clarify the role of cyclic structure, we first examine BF-hypergraphs without weak BF-hypercycles. In classical graph theory [26, 27], a directed graph without cycles has a sink and a source. Regarding a weak BF-hypercycle, a similar argument is suggested [23].

PROPOSITION 4.1 A finite BF-hypergraph $\mathscr{G}_{BF}$ without weak BF-hypercycles possesses a BF-sink and a BF-source.

(proof)
Assume that a BF-sink does not exist in $\mathscr{G}_{BF}=(V,\overrightarrow{E})$. Then, every vertex in V has at least one B-outflow or F-outflow. For a fixed vertex $v_1\in V$, we consider a weak hyperpath Y_n and a vertex set $V_n\subset V$ such that:

$$\begin{aligned} Y_n &:= (v_1,\overrightarrow{e_1},v_2,\overrightarrow{e_2},\ldots,\overrightarrow{e_n},v_{n+1}) \\ V_n &:= \bigcup_{i=1}^{n}\{v_i\}. \end{aligned} \tag{4.1}$$

The hyperpath Y_1 exists and $|V_1|=1$. If Y_n exists, then Y_{n+1} also exists, since v_{n+1} has at least one B-outflow or F-outflow. Meanwhile, if $|V_n|=n$, then $|V_{n+1}|=n+1$ from the fact, $v_{n+1}\notin V_n$, which is required to avoid the formation of a weak BF-hypercycle. However, this contradicts the assumption $|V|<\infty$. The existence of a BF-source follows symmetrically by applying the same argument to the symmetric image of the hypergraph. ■

Therefore, a BF-hypergraph without weak BF-hypercycles in $\mathscr{G}_T$ has a BF-sink and BF-source. While Proposition 4.1 is a general result, Proposition 4.2 demonstrates the structural requirements imposed on $\mathscr{G}_T$ by the standing assumptions, using Lemma 4.1.

LEMMA 4.1 $T_{S\to\{v\}}=0$ for every nonempty subset $S\subset K_v$, if and only if $I_{(K_v):v'}|_v := H_{\{v'\}}|_{\{v\}} - H_{\{v'\}}|_{\{v\}\cup K_v}=0$.

(proof)

($\Longrightarrow$) Since,

$$I_{(K_v):v'}|_v = \sum_{l=1}^{|K_v|}(-1)^{l+1}\sum_{S\subset K_v,|S|=l} T_{S\to\{v\}}, \tag{4.2}$$

the result follows.
($\Longleftarrow$) If $I_{(K_v):v'}|_v = 0$, $T_{u\to v} = 0$ for $\forall S := \{u\} \subset K_v$, such that $|S| = 1$, since,

$$I_{(K_v):v'}|_v = T_{u\to v} + I_{(K_v\setminus S):v'}|_{v,S}, \tag{4.3}$$

and $T_{u\to v} \geqslant 0$ and $I_{(K_v\setminus S):v'}|_{v,S} \geqslant 0$.
Now, suppose that, for $n \in \mathbb{N}$, $T_{S\to\{v\}} = 0$ for $\forall S \subset K_v$, such that $|S| \leqslant n$. Regarding an arbitrary set $P \subset K_v$ with $|P| = n+1$, one obtains $I_{(P):v'}|_v = 0$, as,

$$I_{(K_v):v'}|_v = I_{(P):v'}|_v + I_{(K_v\setminus P):v'}|_{v,P} = 0, \tag{4.4}$$

and $I_{(P):v'}|_v \geqslant 0$ and $I_{(K_v\setminus P):v'}|_{v,P} \geqslant 0$.
Then,

$$I_{(P):v'}|_v = \sum_{l=1}^{n}(-1)^{l+1}\sum_{S\subset P,|S|=l} T_{S\to\{v\}} + (-1)^{n+2}T_{P\to\{v\}} = 0. \tag{4.5}$$

From the assumption, $T_{P\to\{v\}} = 0$.
The induction leads to the desired result. The same argument with reversed temporal indices applies to $T^{\dagger}_{R\to\{v\}}$ and $I_{(Q'_v):v}|_{v'}$. ■

PROPOSITION 4.2 Under deterministic reversible dynamics and the strong blocking flows condition,

$$\begin{aligned} \mathrm{In}_B(v) \neq \emptyset \quad &\Longleftrightarrow \quad H_{v'}|_v > 0, \\ \mathrm{Out}_F(v) \neq \emptyset \quad &\Longleftrightarrow \quad H_v|_{v'} > 0. \end{aligned} \tag{4.6}$$

An F-inflow requires a B-inflow, and a B-outflow requires an F-outflow. Every non-isolated vertex has a B-inflow or an F-outflow.

(proof)
Suppose that $\mathrm{In}_B(v) = \emptyset$. Then, $T_{S\to\{v\}} = 0$ for every nonempty $S \subset K_v$. Lemma 4.1 and the definition of K_v give $H_{v'}|_v = 0$. Conversely, if $H_{v'}|_v = 0$, every multiple transfer entropy whose target contains v vanishes, since v' is determined by a conditioning variable. This proves the first equivalence and excludes an F-inflow without a B-inflow.
The strong blocking flows condition identifies an F-outflow in $\mathscr{G}_T$ with a B-inflow in $\mathscr{G}_{T^\dagger}$. Applying the reversed form of Lemma 4.1 to Q_v proves the second equivalence. If $H_v|_{v'} = 0$, every reversed transfer entropy whose target contains v vanishes. Strong blocking then excludes every transfer-entropy hyperarc whose source contains v, including a B-outflow.
If both conditional entropies vanish, all incoming and outgoing transfer-entropy hyperarcs vanish, including those with multiple vertices in both tail and head. Thus, v is isolated. This proves the remaining assertion. ■

The proposition does not require every non-isolated vertex to have a B-inflow. For example, let v be a fair bit and $u = 0$, with the reversible swap $v' = u$, $u' = v$. The only direct transfer-entropy hyperarc

is $v \to u$, and the reversed hyperarc is $u \to v$, both with value $\ln 2$. Strong blocking holds, but v is a BF-source and u is a BF-sink.

PROPOSITION 4.3 Consider an information flow hypergraph $\mathscr{G}_{\Delta I} = (\mathscr{G}_T, \mathscr{G}_{T^\dagger})$ satisfying the strong blocking flows condition under deterministic reversible dynamics. Let v be a BF-sink in $\mathscr{G}_T$. Then, $\Delta H_v > 0$. For a BF-source, $\Delta H_v < 0$.

(proof)
Consider a BF-sink v with its dynamical neighbors K_v and backward dynamical neighbors Q_v in $\mathscr{G}_T$. From Eq. (3.11), the dynamics of v's entropy H_v is:

$$\Delta H_v = I_{(K_v):v'}|_v - I_{(Q'_v):v}|_{v'}. \tag{4.7}$$

If a B-inflow $(e^+, \{v\})$ in $\mathscr{G}_{T^\dagger}$ is nonzero, the corresponding F-outflow $(\{v\}, e^+)$ in $\mathscr{G}_T$ is also nonzero from the strong blocking flows condition. However, it is not the case, as v is a BF-sink in $\mathscr{G}_T$. Hence, all B-inflows to v in $\mathscr{G}_{T^\dagger}$ are zero. The reversed form of Lemma 4.1 gives $I_{(Q'_v):v}|_{v'} = 0$, resulting in:

$$\Delta H_v = I_{(K_v):v'}|_v \geqslant 0. \tag{4.8}$$

From Lemma 4.1, if $I_{(K_v):v'}|_v = 0$, all B-inflows in $\mathscr{G}_T$ vanish. As a BF-sink has a B-inflow or an F-inflow, and an F-inflow requires a B-inflow by Proposition 4.2, it is not the case, resulting in $\Delta H_v > 0$. For a BF-source, $I_{(K_v):v'}|_v = 0$ because it has no B-inflow. Its B-outflow or F-outflow requires an F-outflow by Proposition 4.2. Therefore, $\Delta H_v = -I_{(Q'_v):v}|_{v'} < 0$. ■

In conclusion, periodic or recurrent entropy dynamics requires a weak BF-hypercycle when the nonempty flow topology is fixed and the standing assumptions hold at every step. Otherwise, Proposition 4.1 gives a fixed BF-sink and BF-source whose entropies change strictly in opposite directions, precluding a return of the entropy vector to, or arbitrarily close to, its initial value. This does not exclude asymptotic relaxation. A changing flow topology requires separate analysis: repeating the swap above yields periodic dynamics while reversing its single edge at each step.

5. Higher-Order Information Dynamics on a Single Weak BF-hypercycle

The previous section demonstrates that periodic or recurrent entropy dynamics requires a weak BF-hypercycle under a fixed nonempty topology and the standing assumptions. This does not imply that every vertex lies on a cycle. However, unlike a cycle in a directed graph, a weak BF-hypercycle can have structural monotonicity in entropy dynamics as in the case of a BF-sink, depending on the types of hyperarcs. Lemma 5.1 shows how the entropy balance is determined by the types of inflow and outflow.

LEMMA 5.1 Consider an information flow hypergraph $\mathscr{G}_{\Delta I} = (\mathscr{G}_T, \mathscr{G}_{T^\dagger})$ satisfying the strong blocking flows condition, under deterministic reversible dynamics. For a vertex v in $\mathscr{G}_{\Delta I}$:
(a) $\Delta H_v > 0$ if v has a B-inflow but no F-outflow in $\mathscr{G}_T$.
(b) ΔH_v may be positive, negative, or zero if v has both a B-inflow and an F-outflow in $\mathscr{G}_T$.
(c) $\Delta H_v < 0$ if v has an F-outflow but no B-inflow in $\mathscr{G}_T$.

(proof)
(a) Let v have a B-inflow $(P,\{v\})$ but no F-outflow in $\mathscr{G}_T$. The strong blocking flows condition requires that F-outflow $(\{v\},P)$ exists in $\mathscr{G}_{T^\dagger}$, but no B-inflow to v exists there. Then, from Lemma 4.1, $I_{(K_v):v'}|_v \neq 0$ and $I_{(Q'_v):v}|_{v'} = 0$. Hence,

$$\begin{aligned} \Delta H_v &= I_{(K_v):v'}|_v - I_{(Q'_v):v}|_{v'} \\ &= I_{(K_v):v'}|_v > 0. \end{aligned} \tag{5.1}$$

(b) Let v have a B-inflow $(P,\{v\})$ and an F-outflow $(\{v\},Q)$ in $\mathscr{G}_T$. The strong blocking flows condition requires that F-outflow $(\{v\},P)$ and B-inflow $(Q,\{v\})$ exist in $\mathscr{G}_{T^\dagger}$. Then, from Lemma 4.1, $I_{(K_v):v'}|_v \neq 0$ and $I_{(Q'_v):v}|_{v'} \neq 0$, where K_v and Q_v are sets of dynamical neighbors and backward dynamical neighbors, respectively. Hence,

$$\Delta H_v = I_{(K_v):v'}|_v - I_{(Q'_v):v}|_{v'}. \tag{5.2}$$

The presence of the two flow types does not determine the sign of this difference.
(c) If v has an F-outflow but no B-inflow, then $I_{(K_v):v'}|_v = 0$ and $I_{(Q'_v):v}|_{v'} > 0$. Thus, $\Delta H_v < 0$. ■

PROPOSITION 5.1 Consider an information flow hypergraph $\mathscr{G}_{\Delta I} = (\mathscr{G}_T, \mathscr{G}_{T^\dagger})$ satisfying the strong blocking flows condition, under deterministic reversible dynamics. A necessary condition for stationarity of entropy dynamics $(H_v)_{v\in V}$ is that every non-isolated vertex v has both a B-inflow and an F-outflow in $\mathscr{G}_T$.

(proof)
Stationarity requires $\Delta H_v = 0$ for every vertex v. Therefore, the necessary condition is obtained by excluding the monotonic cases. Proposition 4.2 ensures that a non-isolated vertex has a B-inflow or an F-outflow. Lemma 5.1 excludes the presence of only one of these types. Hence, both are required. Their aggregate contributions must also satisfy

$$I_{(K_v):v'}|_v = I_{(Q'_v):v}|_{v'}. \tag{5.3}$$

The presence of both flow types alone does not impose this numerical equality or imply stationarity of the joint distribution. ■

A cycle in graph theory satisfies the necessary condition of Proposition 5.1, as the directed edge of the cycle is both B-arc and F-arc.

6. Conclusion

Higher-order interactions can generate higher-order correlations that are not reducible to pairwise relationships. To characterize the topology of their temporal evolution, this paper introduced an information flow hypergraph by associating direct multiple transfer entropy and multiple reversed transfer entropy with hyperarcs. The BF-hypergraph substructure determines the entropy balance of individual vertices,

under the strong blocking flows condition and deterministic reversible dynamics, using K_v and Q_v for the two temporal directions. A finite BF-hypergraph without weak BF-hypercycles possesses a BF-sink and a BF-source. These assumptions require a non-isolated vertex to have a B-inflow or an F-outflow in a direct transfer entropy hypergraph $\mathscr{G}_T$. At a BF-sink of $\mathscr{G}_T$, the entropy increases strictly; at a BF-source, it decreases strictly. Consequently, periodic or recurrent entropy dynamics requires a weak BF-hypercycle when the nonempty flow topology is fixed and these assumptions hold at every step. However, a weak BF-hypercycle alone does not guarantee stationarity. A further necessary condition is that each non-isolated vertex has both a B-inflow and an F-outflow in $\mathscr{G}_T$, with numerically balanced aggregate contributions. This condition also clarifies the role of the residual term. A stationary state with a B-inflow but no compensating F-outflow is incompatible with the present assumptions. If backward reconstruction fails, the residual term in an entropy expansion need not vanish.

The present analysis focuses on sinks, sources, and weak BF-hypercycles. Directed hypergraphs possess additional structural properties, including spectral properties of hypergraph Laplacians [37], that may provide further insight into information dynamics. Another natural extension is to study the relationship between an explosive behavior of higher-order dynamics [2] and information-dynamic stationarity discussed in this paper.

A. Proof of Proposition 3.2

A.1 *Main Proof*

One can establish a non-trivial example involving dependence in the initial distribution. Define $V = \{X, Y, C, \zeta\}$ with binary states and the deterministic reversible dynamics:

$$\begin{aligned} \Gamma_X(t+\Delta t) &= \Gamma_C(t), \\ \Gamma_C(t+\Delta t) &= \Gamma_X(t), \\ \Gamma_Y(t+\Delta t) &= \Gamma_Y(t) \oplus \Gamma_\zeta(t), \\ \Gamma_\zeta(t+\Delta t) &= \Gamma_\zeta(t). \end{aligned} \tag{A.1}$$

At the step under consideration, let $\Gamma_X = 0$, let Γ_C, Γ_ζ be independent fair bits, and set $\Gamma_Y = \Gamma_C \oplus N$, where $N \sim B(\varepsilon)$ is independent of them and $0 < \varepsilon < 1/2$. The variable N specifies the initial distribution. Define $h(\varepsilon) := -\varepsilon \ln \varepsilon - (1-\varepsilon)\ln(1-\varepsilon)$.
Then, $K_X = \{C\}$, $K_Y = \{\zeta\}$, and $K_C = K_\zeta = \emptyset$. The only nonzero direct transfer-entropy hyperarcs are $C \to X$ and $\zeta \to Y$, both with value $\ln 2$. Indeed, any target containing C or ζ contains a future variable determined by the conditioning variables. For the remaining joint target $\{X, Y\}$, C and ζ are conditionally independent given Y, so its multiple transfer entropies from subsets of $K_{\{X,Y\}} = \{C, \zeta\}$ vanish. The imperfect correlation between C and Y does not provide another sufficient neighbor set for X.
While $T_{Y\to X} = \ln 2 - h(\varepsilon) > 0$, X is not connected to Y in $\mathscr{G}_T$. Hence, $\mathscr{G}_T^* \neq \mathscr{G}_{total}^*$ in this example. ■

The subsequent sections explain how initial dependence and common causes can yield indirect statistical relationships that are not represented by chains in $\mathscr{G}_T$.

A.2 *Total Neighbors*

For a vertex set $S \subset V$, let $R_S^1 \subset V$ be the union of possible dynamical neighbors of S: $R_S^1 := \bigcup_{K_S \in \overline{\mathscr{K}}_S} K_S$. Then, define $R_S^{i+1} \subset V$ as the union of possible dynamical neighbors of $R_S^i \subset V$ for $i \in \mathbb{N}$: $R_S^{i+1} := \bigcup_{K_{R_S^i} \in \overline{\mathscr{K}}_{R_S^i}} K_{R_S^i}$. Since V is finite, there is $N \in \mathbb{N}$ such that,

$$\bigcup_{i=1}^{N} R_S^i = \bigcup_{i=1}^{\infty} R_S^i. \tag{A.2}$$

An empty layer remains empty. Hence, the set of total neighbors of S is defined as:

$$K_S^{total} := \bigcup_{i=1}^{N} R_S^i. \tag{A.3}$$

LEMMA A.1 If $u \in K_S^{total}$, then $\exists s \in S$ such that s is connected to u in $\mathscr{G}_T$.

(proof)
It is enough to show that, for $u \in K_S$ for $K_S \in \overline{\mathscr{K}}_S$, $\exists (e^+, e^-) \in \overrightarrow{E}_T$ such that $u \in e^+$ and $e^- \cap S \neq \emptyset$. Let $K_S = \{u\} \cup W$, and $S = \{s_1, \ldots, s_m\}$. By the definition of K_S,

$$H_{S'}|_{S \cup W} > 0. \tag{A.4}$$

Then, there is $s \in S$ such that $H_{s'}|_{S \cup W} > 0$, as,

$$\begin{aligned} H_{S'}|_{S \cup W} &= H_{s'_1}|_{S \cup W} + \sum_{i=2}^{m} H_{s'_i}|_{\{s'_1, \ldots, s'_{i-1}\} \cup S \cup W} \\ &\leqslant \sum_{i=1}^{m} H_{s'_i}|_{S \cup W}. \end{aligned} \tag{A.5}$$

Let $L_s := [S \setminus \{s\}] \cup [K_S \setminus \{u\}]$. As $H_{s'}|_{\{s,u\} \cup L_s} \leqslant H_{S'}|_{\{s,u\} \cup L_s} = 0$, $H_{s'}|_{\{s,u\} \cup L_s} = 0$. Thus, one can obtain a minimal subset $K_s \subset L_s \cup \{u\}$ such that $H_{s'}|_{\{s\} \cup K_s} = 0$, by reducing the set $L_s \cup \{u\}$. Since $H_{s'}|_{\{s\} \cup L_s} > 0$, $u \in K_s$. Then,

$$\begin{aligned} T_{u \to s}|_{K_s \setminus \{u\}} &:= H_{s'}|_{\{s\} \cup [K_s \setminus \{u\}]} - H_{s'}|_{\{s\} \cup K_s} \\ &= H_{s'}|_{\{s\} \cup [K_s \setminus \{u\}]} > 0. \end{aligned} \tag{A.6}$$

Since,

$$T_{u \to s}|_{K_s \setminus \{u\}} = T_{u \to s} + \sum_{i=1}^{|K_s|-1} (-1)^i \sum_{W \subset K_s \setminus \{u\},\ |W|=i} T_{\{u\} \cup W \to \{s\}}, \tag{A.7}$$

there is a nonzero term $T_{e^+\to\{s\}}$ among $T_{u\to s}$, and $T_{\{u\}\cup W\to\{s\}}$ with $k := |e^+|$. One can confirm $e^+ \subset K_s$. Hence, the hyperarc $(e^+, \{s\})$ is an element of $\overrightarrow{E}_T$. Given that $u \in e^+$, s is connected to u in $\mathscr{G}_T$. Applying this argument successively to the layers R_S^j, $1 \leqslant j \leqslant N$, gives a weak hyperpath from u to S. ■

For the multi-step statements below, suppose that the neighbor relations used in these layers remain valid throughout the time interval. Then, the state dynamics of $S^{total} := S \cup K_S^{total}$ is autonomous: $\Gamma_{S^{total}}(t+\Delta t) = F(\Gamma_{S^{total}}(t), t)$, where $\Gamma_{S^{total}}(t) := (\Gamma_{w_1}(t), \ldots, \Gamma_{w_h}(t))$ with $S^{total} = \{w_1, \ldots, w_h\}$.

A.3 *Two Independent Autonomous Systems*

Consider two total neighbors S^{total} and U^{total} such that $S^{total} \cap U^{total} = \emptyset$.

LEMMA A.2 If $I^{co}_{S\cup U}(t_0) = 0$ for all nonempty $S \subset S^{total}$ and $U \subset U^{total}$, then $I^{co}_{S\cup U}(t) = 0$ and $T_{S\to U}(t) = 0$ for $t \geqslant t_0$.

(proof)
Since S^{total} and U^{total} are autonomous, one obtains,

$$\begin{aligned} \Gamma_{S^{total}}(t) &= F_t(\Gamma_{S^{total}}(t_0)) \\ \Gamma_{U^{total}}(t') &= G_{t'}(\Gamma_{U^{total}}(t_0)), \end{aligned} \tag{A.8}$$

for $t, t' \geqslant t_0$.
On the other hand, if $I^{co}_{S\cup U}(t_0) = 0$ for all nonempty $S \subset S^{total}$ and $U \subset U^{total}$, then

$$H_{S^{total}}(t_0) - H_{S^{total}|_{U^{total}}}(t_0) = 0.$$

This indicates the factorization of joint probability.
As $S^{total} \cap U^{total} = \emptyset$ and statistically independent at $t = t_0$, $\Gamma_{S^{total}}(t)$ and $\Gamma_{U^{total}}(t')$ are statistically independent for $\forall t, t' \geqslant t_0$. ■

A.4 *Classification of Indirect Relationships*

We analyze case by case whether S is connected to U in $\mathscr{G}_T$ for $\forall (U,S) \in \overrightarrow{E}_{total}$. Fix an arbitrary hyperarc $(U,S) \in \overrightarrow{E}_{total}$. If $U \cap K_S^{total} \neq \emptyset$, Lemma A.1 requires that S is connected to U in $\mathscr{G}_T$. Let $U \cap K_S^{total} = \emptyset$.
If $U^{total} \cap S^{total} = \emptyset$, Lemma A.2 implies that (U,S) cannot be an element of $\overrightarrow{E}_{total}$ under the condition of initial independence.
Let $U^{total} \cap S^{total} \neq \emptyset$. Here, an element of $U^{total} \cap S^{total}$ can act as a common cause. The example in Sec. A.1 instead uses dependence between disjoint autonomous systems in their initial distribution.

B. The Proof of Proposition 3.3

Since $\overrightarrow{E}_{\bar{T}} \subset \overrightarrow{E}_{total}$, it follows immediately that $\mathscr{G}^*_{\bar{T}} \subset \mathscr{G}^*_{total}$. It therefore remains to show that $\mathscr{G}^*_{total} \subset \mathscr{G}^*_{\bar{T}}$.
Consider $(U,S) \in \overrightarrow{E}_{total}$, and fix $K_S \in \overline{\mathscr{K}}_S$. Then, $T_{U\to S} \neq 0$. If $U \cap K_S \neq \emptyset$, then $(U,S) \in \overrightarrow{E}_{\bar{T}}$ by

definition. Let $U \cap K_S = \emptyset$. Here, $T_{U\to S}|_{K_S} := I^{\mathrm{co}}_{S'\cup U}|_{S\cup K_S}$, and its decomposition is obtained as:

$$T_{U\to S}|_{K_S} = T_{U\to S} + \sum_{i=1}^{|K_S|}(-1)^i \sum_{W\subset K_S, |W|=i} T_{U\cup W\to S}. \tag{B.1}$$

Since $T_{U\to S}|_{K_S} = 0$ and $T_{U\to S} \neq 0$, there is a nonzero term $T_{U\cup W\to S}$ for $\emptyset \neq W \subset K_S$ and $i = |W|$. Therefore, the hyperarc $(U\cup W, S) \in \overrightarrow{E}_{total}$, and one concludes that $(U\cup W, S) \in \overrightarrow{E}_{\bar{T}}$ from the fact that $(U\cup W)\cap K_S = W \neq \emptyset$. This leads to $\mathscr{G}^*_{total} \subset \mathscr{G}^*_{\bar{T}}$. ■

C. Examples of Hyperarc

The following examples illustrate local flow quantities with stochastic inputs. The focal updates are deterministic given their inputs, but backward reconstruction of all vertices is not assumed. Thus, these examples are not used as realizations of the reversible framework of the main propositions. The closed copying dynamics in Sec. 3.3 supplies such a realization.

C.1 *A Single B-arc*

Consider a three-component system a, b, and p with the dynamics:

$$\begin{aligned} \Gamma_a(t+\Delta t) &\sim B(\frac{1}{2}) \\ \Gamma_b(t+\Delta t) &\sim B(\frac{1}{2}) \\ \Gamma_p(t+\Delta t) &= \Gamma_a(t) \oplus \Gamma_b(t), \end{aligned} \tag{C.1}$$

where the updates of Γ_a and Γ_b are independent of each other and of the current state, $B(q)$ denotes Bernoulli distribution with probability q, and $\oplus$ denotes the XOR operation. We evaluate the quantities under the invariant distribution of independent fair bits.
Then,

$$\begin{aligned} T_{a\to p} &= T_{b\to p} = 0 \\ T^{\dagger}_{p\to a} &= T^{\dagger}_{p\to b} = 0 \\ T_{\{a,b\}\to\{p\}} &= -\ln 2 \\ T^{\dagger}_{\{p\}\to\{a,b\}} &= -\ln 2 \\ T_{\{p\}\to\{a,b\}} &= T^{\dagger}_{\{a,b\}\to\{p\}} = 0. \end{aligned} \tag{C.2}$$

These quantities describe a B-channel $\{a,b\} \to \{p\}$ and its reversed counterpart. At the focal vertex p, $K_p = \{a,b\}$, but $\alpha_1^{K_p\to p} = -\ln 2$. The displayed future variables do not reconstruct its past state. Thus, the local channel correspondence does not establish the reversible assumptions for the whole system.

C.2 *A Single F-arc*

Consider a four-component system p, ζ, d, and e with the dynamics:

$$\begin{aligned}
\Gamma_p(t+\Delta t) &\sim B(\frac{1}{2}) \\
\Gamma_\zeta(t+\Delta t) &\sim B(\frac{1}{2}) \\
(\Gamma_d,\Gamma_e)(t+\Delta t) &= \begin{cases} (\Gamma_\zeta(t),\Gamma_\zeta(t)) & \text{if } \Gamma_p(t)=1 \\ (\Gamma_\zeta(t),1-\Gamma_\zeta(t)) & \text{if } \Gamma_p(t)=0 \end{cases},
\end{aligned} \tag{C.3}$$

where the updates of Γ_p and Γ_ζ are independent of each other and of the current state. We evaluate the quantities under the invariant distribution of independent fair bits.
This dynamics provides a stochastic inverse of the XOR operation after relabeling the input as $1-\Gamma_p$. Then,

$$\begin{aligned}
T_{p\to d} &= T_{p\to e} = 0 \\
T^\dagger_{d\to p} &= T^\dagger_{e\to p} = 0 \\
T_{\{p\}\to\{d,e\}} &= -\ln 2 \\
T^\dagger_{\{d,e\}\to\{p\}} &= -\ln 2 \\
T_{\{d,e\}\to\{p\}} &= T^\dagger_{\{p\}\to\{d,e\}} = 0.
\end{aligned} \tag{C.4}$$

The remaining nonzero channels for the focal targets are

$$\begin{aligned}
T_{\zeta\to d} &= \ln 2, \\
T_{\{p,\zeta\}\to\{e\}} &= T_{\{p,\zeta\}\to\{d,e\}} = -\ln 2.
\end{aligned} \tag{C.5}$$

Their reversed counterparts are

$$\begin{aligned}
T^\dagger_{d\to\zeta} &= \ln 2, \\
T^\dagger_{\{e\}\to\{p,\zeta\}} &= T^\dagger_{\{d,e\}\to\{p,\zeta\}} = -\ln 2.
\end{aligned} \tag{C.6}$$

Thus, the F-channel $\{p\}\to\{d,e\}$ is part of a larger local support, including a B-channel and a channel with multiple vertices in both tail and head. These local quantities have the corresponding reversal of support, but the stochastic input system is not asserted to possess all backward dynamical-neighbor sets.

REFERENCES

1. Battiston F, Cencetti G, Iacopini I et al. Networks beyond pairwise interactions: structure and dynamics. *Phys Rep* 2020;**874**:1–92.
2. Battiston F, Amico E, Barrat A et al. The physics of higher-order interactions in complex systems. *Nat Phys* 2021;**17**(10):1093–98.
3. Rosas F. E, Mediano P. A. M, Luppi A. I et al. Disentangling high-order mechanisms and high-order behaviours in complex systems. *Nat Phys* 2022;**18**(5):476–7.
4. Gallo L, Muolo R, Gambuzza L. V et al. Synchronization induced by directed higher-order interactions. *Commun Phys* 2022;**5**:263.

5. Majhi S, Perc M, Ghosh D. Dynamics on higher-order networks: a review. *J R Soc Interface* 2022;**19**(188):20220043.

6. Zhang Y, Lucas M, Battiston F. Higher-order interactions shape collective dynamics differently in hypergraphs and simplicial complexes. *Nat Commun* 2023;**14**:1605.

7. Bick C, Gross E, Harrington H. A et al. What are higher-order networks?. *SIAM Rev* 2023;**65**(3):686–731.

8. Matsuda H. Physical nature of higher-order mutual information: intrinsic correlations and frustration. *Phys Rev E* 2000;**62**(3):3096–102.

9. Schneidman E, Still S, Berry M. J et al. Network information and connected correlations. *Phys Rev Lett* 2003;**91**(23):238701.

10. Rosas F. E, Mediano P. A. M, Gastpar M et al. Quantifying high-order interdependencies via multivariate extensions of the mutual information. *Phys Rev E* 2019;**100**(3):032305.

11. Williams P. L, Beer R. D. Nonnegative decomposition of multivariate information. 2010;.

12. Kolchinsky A. A novel approach to the partial information decomposition. *Entropy* 2022;**24**(3):403.

13. McGill W. J. Multivariate information transmission. *Trans IRE Prof Group Inf Theory* 1954;**4**(4):93–111.

14. Fano R. M. *The Transmission of Information*, volume 65. Massachusetts Institute of Technology, Research Laboratory of Electronics, Cambridge, MA 1949.

15. Ting H. K. On the amount of information. *Theory Probab Appl* 1962;**7**(4):439–47.

16. Han T. S. Multiple mutual informations and multiple interactions in frequency data. *Inf Control* 1980;**46**(1):26–45.

17. Yeung R. W. A new outlook on Shannon's information measures. *IEEE Trans Inf Theory* 1991;**37**(3):466–74.

18. Bell A. J. The co-information lattice. In *Proceedings of the Fifth International Conference on Independent Component Analysis and Blind Signal Separation*, pages 921–6, Nara, Japan 2003;.

19. Krippendorff K. Information of interactions in complex systems. *Int J Gen Syst* 2009;**38**(6):669–80.

20. Lee Y. Dynamical equations of mutual information and transfer entropy in discrete time. *AIP Adv* 2026;**16**(5):055108.

21. Gallo G, Longo G, Pallottino S et al. Directed hypergraphs and applications. *Discrete Appl Math* 1993;**42**(2–3):177–201.

22. Nguyen S, Pretolani D, Markenzon L. On some path problems on oriented hypergraphs. *RAIRO Theor Inform Appl* 1998;**32**(1-3):1–20.

23. Bretto A. *Hypergraph Theory: An Introduction*. Mathematical Engineering. Springer International Publishing, Cham 2013.

24. Steinmetz M, Torralba Á. Bridging the gap between abstractions and critical-path heuristics via hypergraphs. In *Proc. ICAPS*, volume 29, pages 473–81 2019;.

25. Lee Y. Information Circulation and External-flow Suppression under Finite Capacity. electronic preprint 2026;https://doi.org/10.2139/ssrn.7421118.

26. Diestel R. *Graph Theory*, volume 173 of *Graduate Texts in Mathematics*. Springer, Berlin, 5 edition 2017.

27. Kleinberg J, Tardos É. *Algorithm Design*. Pearson Education, Boston, MA 2005.

28. Shannon C. E, Weaver W. *The Mathematical Theory of Communication*. University of Illinois Press, Urbana, IL 1949.

29. Cover T. M, Thomas J. A. *Elements of Information Theory*. Wiley-Interscience, New York, NY 1991.

30. Barnett L, Barrett A. B, Seth A. K. Granger causality and transfer entropy are equivalent for Gaussian variables. *Phys Rev Lett* 2009;**103**(23):238701.

31. Schreiber T. Measuring information transfer. *Phys Rev Lett* 2000;**85**(2):461–4.

32. Bossomaier T, Barnett L, Harré M et al. *An Introduction to Transfer Entropy: Information Flow in Complex Systems*. Springer International Publishing, Cham 2016.

33. Ito S. Backward transfer entropy: informational measure for detecting hidden Markov models and its interpretations in thermodynamics, gambling, and causality. *Sci Rep* 2016;**6**:36831.

34. Spinney R. E, Lizier J. T, Prokopenko M. Transfer entropy in physical systems and the arrow of time. *Phys Rev E* 2016;**94**(2):022135.

35. Spinney R. E, Lizier J. T, Prokopenko M. Entropy balance and information processing in bipartite and nonbi-

partite composite systems. *Phys Rev E* 2018;**98**(3):032141.

36. Ausiello G, Laura L. Directed hypergraphs: introduction and fundamental algorithms—a survey. *Theor Comput Sci* 2017;**658**:293–306.

37. Jost J, Mulas R. Hypergraph Laplace operators for chemical reaction networks. *Adv Math* 2019;**351**:870–96.