

practice_exercise

Ekta Chaudhary

28/06/2020

```
library(tidyverse)
```

```
## -- Attaching packages ----- tidyverse 1.2
```

```
## v ggplot2 3.2.1    v purrr  0.3.2
## v tibble  2.1.3    v dplyr  0.8.3
## v tidyr   1.0.0    v stringr 1.4.0
## v readr   1.3.1    v forcats 0.4.0
```

```
## -- Conflicts ----- tidyverse_conflicts
```

```
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()    masks stats::lag()
```

```
library(readxl)
library(dplyr)
library(sqldf)
```

```
## Loading required package: gsubfn
```

```
## Loading required package: proto
```

```
## Loading required package: RSQLite
```

```
library(ggthemes)
```

```
practice_data = read_excel("./data/Practice_exercise.xlsx", sheet = "Data") %>%
  janitor::clean_names() %>%
  select(observation_number, quarter, employee_id, sex = sex_male_1, race, age, hospital_visit = hospital_visit_1)
  mutate(
    age_cat = case_when(
      age < 30 ~ 1,
      age <= 45 ~ 2,
      age > 45 ~ 3
    )
  )
```

```
#Checking for missing data
```

```
sapply(practice_data, function(x) sum(is.na(x)))
```

```
## observation_number      quarter      employee_id
##                0                0                0
##                sex                race                age
```

```
##           71           2123           0
##   hospital_visit      salary      health_score
##           0           0           0
##           age_cat
##           0
```

#Checking for missing data

```
practice_data %>%
  select(everything()) %>% # replace to your needs
  summarise_all(funs(sum(is.na(.))))
```

```
## Warning: funs() is soft deprecated as of dplyr 0.8.0
## Please use a list of either functions or lambdas:
##
##   # Simple named list:
##   list(mean = mean, median = median)
##
##   # Auto named with `tibble::lst()`:
##   tibble::lst(mean, median)
##
##   # Using lambdas
##   list(~ mean(., trim = .2), ~ median(., na.rm = TRUE))
## This warning is displayed once per session.
```

```
## # A tibble: 1 x 10
##   observation_num~ quarter employee_id sex race age hospital_visit
##             <int>   <int>       <int> <int> <int> <int>           <int>
## 1             0       0           0    71  2123    0             0
## # ... with 3 more variables: salary <int>, health_score <int>,
## #   age_cat <int>
```

#finding the minimum and maximum values of each variable

```
sapply(practice_data, function(x) min(x))
```

```
## observation_number      quarter      employee_id
##   1.000000e+00   1.000000e+00   1.000000e+00
##           sex           race           age
##           NA           NA       7.000000e+00
##   hospital_visit      salary      health_score
##   0.000000e+00   2.835070e+04   6.265991e-01
##           age_cat
##   1.000000e+00
```

```
sapply(practice_data, function(x) max(x))
```

```
## observation_number      quarter      employee_id
##   19103.00           12.00           2000.00
##           sex           race           age
##           NA           NA           172.00
```

```
##      hospital_visit      salary      health_score
##           1.00          68826.34           10.00
##           age_cat
##           3.00
```

#checking the number of employees with health score outside the range of data

```
practice_data %>%
  count(
    health_sc_6 = ifelse(health_score > 6, 1, 0)
  )
```

```
## # A tibble: 2 x 2
##   health_sc_6      n
##       <dbl> <int>
## 1           0 17865
## 2           1  1238
```

```
practice_data %>%
  select(
    employee_id, sex
  ) %>%
  filter(
    is.na(sex)
  ) %>%
  group_by(
    employee_id
  ) %>%
  summarise(
    missing = sum(is.na(sex))
  )
```

```
## # A tibble: 7 x 2
##   employee_id missing
##       <dbl>   <int>
## 1       1994      10
## 2       1995       9
## 3       1996      12
## 4       1997      11
## 5       1998      12
## 6       1999       7
## 7       2000      10
```

```
practice_data %>%
  select(
    employee_id, race
  ) %>%
  filter(
    is.na(race)
  ) %>%
  group_by(
    employee_id
```

```

) %>%
  summarise(
    miss = sum(is.na(race))
  )

```

```

## # A tibble: 220 x 2
##   employee_id miss
##         <dbl> <int>
## 1           8   10
## 2          10   12
## 3          13    9
## 4          22    9
## 5          36   12
## 6          38   12
## 7          48   10
## 8          49    7
## 9          51    8
## 10         55    9
## # ... with 210 more rows

```

#Calculating the number of employees in each age group for each quarter

```

emp_data = practice_data %>%
  mutate(
    quarter = factor(
      quarter),
    age_cat = factor(age_cat)
  )

```

```

emp_data = emp_data %>%
  select(
    employee_id, quarter, age_cat
  ) %>%
  group_by(
    quarter, age_cat
  ) %>%
  tally()


```

```

e <- ggplot(emp_data, aes(x = quarter, y = n, group = age_cat)) +
  geom_line(aes(color = age_cat)) +
  geom_point(aes(color = age_cat)) + labs(x = "Quarter", y = "Number of employees", title = "Number of employees by quarter and age group")
e

```

Number of employees in different age group for each quarter



#checking the trend in average salary over time

```
practice_data %>%
  select(
    salary, quarter
  ) %>%
  group_by(
    quarter
  ) %>%
  summarise(
    avg_salary = mean(salary)
  )
```

```
## # A tibble: 12 x 2
##   quarter avg_salary
##   <dbl>     <dbl>
## 1       1    43628.
## 2       2    44274.
## 3       3    45021.
## 4       4    45531.
## 5       5    46133.
## 6       6    46948.
## 7       7    47780.
## 8       8    48667.
## 9       9    49562.
## 10      10    50498.
## 11      11    51433.
```


```
## 12      12      52376.
```

```
salary_data = practice_data %>%  
  mutate(  
    age_cat = factor(age_cat),  
    quarter = factor(quarter))
```

```
#checking the trend in average salary over time by age group
```

```
salary_data = salary_data %>%  
  select(  
    salary, quarter, age_cat  
  ) %>%  
  group_by(  
    quarter, age_cat  
  ) %>%  
  summarise(  
    avg_salary = mean(salary)  
  )
```


```
p <- ggplot(salary_data, aes(x = quarter, y = avg_salary, group = age_cat)) +  
  geom_line(aes(color = age_cat)) +  
  geom_point(aes(color = age_cat)) + labs(x = "Quarter", y = "Average Salary", title = "Trend in average  
p
```



```
#checking the trend in mean health score over time
```

```
hc = practice_data %>%
  select(
    health_score, quarter
  ) %>%
  group_by(
    quarter
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
px <- ggplot(hc, aes(x = factor(quarter), y = avg_score)) +
  geom_bar(stat = "identity", position = "dodge") + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean Health Score over time")
px
```



#Mean health score over time by age group


```
practice_data %>%
  select(
    health_score, quarter, age_cat
  ) %>%
  group_by(
    quarter, age_cat
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
## # A tibble: 36 x 3
## # Groups:   quarter [12]
##   quarter age_cat avg_score
##   <dbl>   <dbl>   <dbl>
## 1       1       1       3.21
## 2       1       2       3.90
## 3       1       3       4.56
## 4       2       1       3.38
## 5       2       2       3.62
## 6       2       3       4.95
## 7       3       1       3.41
## 8       3       2       3.63
## 9       3       3       4.16
## 10      4       1       3.34
## # ... with 26 more rows
```

```
health_sc = practice_data %>%
  mutate(
    age_cat = factor(age_cat),
    quarter = factor(quarter))
```

```
health_sc = health_sc %>%
  select(
    health_score, quarter, age_cat
  ) %>%
  group_by(
    quarter, age_cat
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
p1 <- ggplot(health_sc, aes(x = quarter, y = avg_score, group = age_cat)) +
  geom_line(aes(color = age_cat)) +
  geom_point(aes(color = age_cat)) + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean Health Score by Quarter")
p1
```


#Mean health score over time by sex

```
health_sex = practice_data %>%
  mutate(
    sex = factor(sex),
    quarter = factor(quarter))
```

```
health_sex = health_sex %>%
  drop_na() %>%
  select(
    health_score, quarter, sex
  ) %>%
  group_by(
    quarter, sex
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
pq <- ggplot(health_sex, aes(x = quarter, y = avg_score, group = sex)) +
  geom_line(aes(color = sex)) +
  geom_point(aes(color = sex)) + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean Health Score over time by sex")
pq
```

Trend in Mean Health Score over time by Sex




#Mean health score over time by race

```
health_race = practice_data %>%
  mutate(
    race = factor(race),
    quarter = factor(quarter))
```

```
health_race = health_race %>%
  drop_na() %>%
  select(
    health_score, quarter, race
  ) %>%
  group_by(
    quarter, race
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
pe <- ggplot(health_race, aes(x = quarter, y = avg_score, group = race)) +
  geom_line(aes(color = race)) +
  geom_point(aes(color = race)) + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean H
pe
```

Trend in Mean Health Score over time by Race



#correcting the data

```
new_data = practice_data %>%
  drop_na() %>%
  filter(
    health_score <= 6,
    age >= 14, age <= 75
  )
```

#checking the trend in mean health score over time

```
hc1 = new_data %>%
  select(
    health_score, quarter
  ) %>%
  group_by(
    quarter
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
sqldf("select quarter, age_cat, avg(health_score)as avg_score
      from new_data
      group by quarter, age_cat")
```

```
##   quarter age_cat avg_score
```

```
## 1      1      1 2.796936
## 2      1      2 3.388951
## 3      1      3 3.425028
## 4      2      1 2.989041
## 5      2      2 3.195572
## 6      2      3 3.988790
## 7      3      1 2.957211
## 8      3      2 3.287013
## 9      3      3 3.764033
## 10     4      1 2.980156
## 11     4      2 3.332257
## 12     4      3 3.922990
## 13     5      1 2.953362
## 14     5      2 3.348889
## 15     5      3 3.963946
## 16     6      1 2.997310
## 17     6      2 3.310626
## 18     6      3 3.863035
## 19     7      1 3.012671
## 20     7      2 3.440885
## 21     7      3 3.899116
## 22     8      1 2.970070
## 23     8      2 3.298553
## 24     8      3 4.017854
## 25     9      1 2.956768
## 26     9      2 3.397101
## 27     9      3 3.911494
## 28    10      1 2.961097
## 29    10      2 3.319969
## 30    10      3 3.710119
## 31    11      1 3.036244
## 32    11      2 3.357914
## 33    11      3 3.924474
## 34    12      1 3.113301
## 35    12      2 3.404411
## 36    12      3 3.954721
```

#Mean health score over time by age group

```
new_data %>%
  select(
    health_score, quarter, age_cat
  ) %>%
  group_by(
    quarter, age_cat
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```


```
## # A tibble: 36 x 3
## # Groups:   quarter [12]
##   quarter age_cat avg_score
```

```
##      <dbl>   <dbl>   <dbl>
## 1         1         1     2.80
## 2         1         2     3.39
## 3         1         3     3.43
## 4         2         1     2.99
## 5         2         2     3.20
## 6         2         3     3.99
## 7         3         1     2.96
## 8         3         2     3.29
## 9         3         3     3.76
## 10        4         1     2.98
## # ... with 26 more rows
```

```
health_score = new_data %>%
  mutate(
    age_cat = factor(age_cat),
    quarter = factor(quarter))
```

```
health_score = health_score %>%
  select(
    health_score, quarter, age_cat
  ) %>%
  group_by(
    quarter, age_cat
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
pn <- ggplot(health_score, aes(x = quarter, y = avg_score, group = age_cat)) +
  geom_line(aes(color = age_cat)) +
  geom_point(aes(color = age_cat)) + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean Health Score by Quarter")
pn
```




```
hosp = practice_data %>%
```

```
  select(
    employee_id, hospital_visit, quarter
  ) %>%
  group_by(
    quarter
  ) %>%
  summarise(
    percent = (sum(hospital_visit)/n())*100
  )
```

```
s <- ggplot(hosp, aes(x = quarter, y = percent, color = percent)) + theme_classic() + geom_line() + geom_point()
  scale_x_discrete(name = "Quarter", limits = c("1","2","3","4","5","6","7","8","9","10","11","12")) +
  scale_y_continuous(name = "Percentage of employees visiting hospitals",
    breaks = seq(0, 25, 5),
    limits = c(0, 25)) + labs(x = "Quarter", y = "Percentage of employees visiting hospitals")
```

Trend in Hospital visits over time



```
w = practice_data %>%
  drop_na() %>%
  mutate(
    sex = factor(sex)
  ) %>%
  select(
    employee_id, quarter, sex
  ) %>%
  group_by(
    quarter, sex
  ) %>%
  summarise(n = n()) %>%
  mutate(freq = n / sum(n)*100)
w
```


```
## # A tibble: 24 x 4
## # Groups:   quarter [12]
##   quarter sex      n freq
##   <dbl> <fct> <int> <dbl>
## 1      1 0      305 50.8
## 2      1 1      295 49.2
## 3      2 0      399 50.9
## 4      2 1      385 49.1
## 5      3 0      503 49.8
## 6      3 1      507 50.2
## 7      4 0      632 49.1
## 8      4 1      655 50.9
```

```
## 9      5 0      726 48.9
## 10     5 1      759 51.1
## # ... with 14 more rows
```

```
t = w %>%
  ggplot(aes(x = quarter, y = freq, fill = sex)) + geom_bar(stat = "identity", legend = c("Female", "Male"),
  scale_y_continuous(name = "Percentage",
                      breaks = seq(0, 100, 10),
                      limits = c(0, 100)) + labs(x = "Quarter", y = "Percentage" , title = "Distribution by Gender over time")
```

```
## Warning: Ignoring unknown parameters: legend
```

```
t
```



```
j = practice_data %>%
  drop_na() %>%
  mutate(
    race = factor(race)
  ) %>%
  select(
    quarter, race
  ) %>%
  group_by(
    quarter, race
  ) %>%
  summarise(n = n()) %>%
```



```
mutate(freq = n / sum(n)*100)
j
```

```
## # A tibble: 36 x 4
## # Groups:   quarter [12]
##   quarter race      n freq
##   <dbl> <fct> <int> <dbl>
## 1      1 1      336  56.
## 2      1 2      167  27.8
## 3      1 3       97  16.2
## 4      2 1      432  55.1
## 5      2 2      222  28.3
## 6      2 3      130  16.6
## 7      3 1      561  55.5
## 8      3 2      293  29.0
## 9      3 3      156  15.4
## 10     4 1      718  55.8
## # ... with 26 more rows
```

```
k = j %>%
  ggplot(aes(x = quarter, y = freq, fill = race)) + geom_bar(stat = "identity") +
  theme_classic() + scale_fill_manual(values = c("dodgerblue4", "skyblue", "pink")) + scale_x_discrete(name = "Quarter",
  scale_y_continuous(name = "Percentage",
    breaks = seq(0, 100, 10),
    limits = c(0, 100)) + labs(x = "Quarter", y = "Percentage" , title = "Distribution by Race over time")
k
```

