

practice_exercise

Ekta Chaudhary

28/06/2020

```
library(tidyverse)
```

```
## -- Attaching packages ----- tidyverse 1.2
```

```
## v ggplot2 3.2.1    v purrr  0.3.2
## v tibble  2.1.3    v dplyr  0.8.3
## v tidyr   1.0.0    v stringr 1.4.0
## v readr   1.3.1    v forcats 0.4.0
```

```
## -- Conflicts ----- tidyverse_conflicts
```

```
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()    masks stats::lag()
```

```
library(readxl)
```

```
library(dplyr)
```

```
library(sqldf)
```

```
## Loading required package: gsubfn
```

```
## Loading required package: proto
```

```
## Loading required package: RSQLite
```

```
practice_data = read_excel("./data/Practice_exercise.xlsx", sheet = "Data") %>%
  janitor::clean_names() %>%
  select(observation_number, quarter, employee_id, sex = sex_male_1, race, age, hospital_visit = hospital)
  mutate(
    age_cat = case_when(
      age < 30 ~ 1,
      age <= 45 ~ 2,
      age > 45 ~ 3
    )
  )
```

```
#Checking for missing data
```

```
sapply(practice_data, function(x) sum(is.na(x)))
```

```
## observation_number      quarter      employee_id
##                0                0                0
##                sex          race          age
##                71          2123                0
##    hospital_visit      salary    health_score
##                0                0                0
##          age_cat
##                0
```

```
#Checking for missing data
```

```
practice_data %>%  
  select(everything()) %>% # replace to your needs  
  summarise_all(funs(sum(is.na(.))))
```

```
## Warning: funs() is soft deprecated as of dplyr 0.8.0  
## Please use a list of either functions or lambdas:  
##  
##   # Simple named list:  
##   list(mean = mean, median = median)  
##  
##   # Auto named with `tibble::lst()`:  
##   tibble::lst(mean, median)  
##  
##   # Using lambdas  
##   list(~ mean(., trim = .2), ~ median(., na.rm = TRUE))  
## This warning is displayed once per session.
```

```
## # A tibble: 1 x 10  
##   observation_num~ quarter employee_id sex race age hospital_visit  
##               <int>   <int>       <int> <int> <int> <int>         <int>  
## 1             0     0           0    71  2123     0             0  
## # ... with 3 more variables: salary <int>, health_score <int>,  
## #   age_cat <int>
```

```
#finding the minimum and maximum values of each variable
```

```
sapply(practice_data, function(x) min(x))
```

```
## observation_number      quarter      employee_id  
##      1.000000e+00      1.000000e+00      1.000000e+00  
##              sex              race              age  
##              NA              NA      7.000000e+00  
##      hospital_visit      salary      health_score  
##      0.000000e+00      2.835070e+04      6.265991e-01  
##              age_cat  
##      1.000000e+00
```

```
sapply(practice_data, function(x) max(x))
```

```
## observation_number      quarter      employee_id  
##      19103.00      12.00      2000.00  
##              sex              race              age  
##              NA              NA      172.00  
##      hospital_visit      salary      health_score  
##              1.00      68826.34      10.00  
##              age_cat  
##              3.00
```

```
#checking the number of employees with health score outside the range of data
```

```
practice_data %>%
  count(
    health_sc_6 = ifelse(health_score > 6, 1, 0)
  )
```

```
## # A tibble: 2 x 2
##   health_sc_6      n
##       <dbl> <int>
## 1           0 17865
## 2           1  1238
```

```
practice_data %>%
  select(
    employee_id, sex
  ) %>%
  filter(
    is.na(sex)
  ) %>%
  group_by(
    employee_id
  ) %>%
  summarise(
    missing = sum(is.na(sex))
  )
```

```
## # A tibble: 7 x 2
##   employee_id missing
##       <dbl>   <int>
## 1       1994       10
## 2       1995        9
## 3       1996       12
## 4       1997       11
## 5       1998       12
## 6       1999        7
## 7       2000       10
```

```
practice_data %>%
  select(
    employee_id, race
  ) %>%
  filter(
    is.na(race)
  ) %>%
  group_by(
    employee_id
  ) %>%
  summarise(
    miss = sum(is.na(race))
  )
```

```
## # A tibble: 220 x 2
##   employee_id miss
```


```
##           <dbl> <int>
##  1             8    10
##  2            10    12
##  3            13     9
##  4            22     9
##  5            36    12
##  6            38    12
##  7            48    10
##  8            49     7
##  9            51     8
## 10            55     9
## # ... with 210 more rows
```

#Calculating the number of employees in each age group for each quarter

```
emp_data = practice_data %>%
  mutate(
    quarter = factor(
      quarter),
    age_cat = factor(age_cat)
  )
```

```
emp_data = emp_data %>%
  select(
    employee_id, quarter, age_cat
  ) %>%
  group_by(
    quarter, age_cat
  ) %>%
  tally()
```

```
e <- ggplot(emp_data, aes(x = quarter, y = n, group = age_cat)) +
  geom_line(aes(color = age_cat)) +
  geom_point(aes(color = age_cat)) + labs(x = "Quarter", y = "Number of employees", title = "Number of employees by quarter and age group")
e
```



#checking the trend in average salary over time

```
practice_data %>%
  select(
    salary, quarter
  ) %>%
  group_by(
    quarter
  ) %>%
  summarise(
    avg_salary = mean(salary)
  )
```

```
## # A tibble: 12 x 2
##   quarter avg_salary
##   <dbl>    <dbl>
## 1       1    43628.
## 2       2    44274.
## 3       3    45021.
## 4       4    45531.
## 5       5    46133.
## 6       6    46948.
## 7       7    47780.
## 8       8    48667.
## 9       9    49562.
## 10      10    50498.
## 11      11    51433.
```


```
## 12      12      52376.
```

```
salary_data = practice_data %>%  
  mutate(  
    age_cat = factor(age_cat),  
    quarter = factor(quarter))
```

```
#checking the trend in average salary over time by age group
```

```
salary_data = salary_data %>%  
  select(  
    salary, quarter, age_cat  
  ) %>%  
  group_by(  
    quarter, age_cat  
  ) %>%  
  summarise(  
    avg_salary = mean(salary)  
  )
```


```
p <- ggplot(salary_data, aes(x = quarter, y = avg_salary, group = age_cat)) +  
  geom_line(aes(color = age_cat)) +  
  geom_point(aes(color = age_cat)) + labs(x = "Quarter", y = "Average Salary", title = "Trend in average  
p
```



```
#checking the trend in mean health score over time
```

```
hc = practice_data %>%
  select(
    health_score, quarter
  ) %>%
  group_by(
    quarter
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
px <- ggplot(hc, aes(x = factor(quarter), y = avg_score)) +
  geom_bar(stat = "identity", position = "dodge") + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean Health Score over time")
px
```



#Mean health score over time by age group


```
practice_data %>%
  select(
    health_score, quarter, age_cat
  ) %>%
  group_by(
    quarter, age_cat
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
## # A tibble: 36 x 3
## # Groups:   quarter [12]
##   quarter age_cat avg_score
##   <dbl>   <dbl>   <dbl>
## 1     1     1     3.21
## 2     1     2     3.90
## 3     1     3     4.56
## 4     2     1     3.38
## 5     2     2     3.62
## 6     2     3     4.95
## 7     3     1     3.41
## 8     3     2     3.63
## 9     3     3     4.16
## 10    4     1     3.34
## # ... with 26 more rows
```

```
health_sc = practice_data %>%
  mutate(
    age_cat = factor(age_cat),
    quarter = factor(quarter))
```

```
health_sc = health_sc %>%
  select(
    health_score, quarter, age_cat
  ) %>%
  group_by(
    quarter, age_cat
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
p1 <- ggplot(health_sc, aes(x = quarter, y = avg_score, group = age_cat)) +
  geom_line(aes(color = age_cat)) +
  geom_point(aes(color = age_cat)) + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean Health Score by Quarter")
p1
```


#Mean health score over time by sex

```
health_sex = practice_data %>%
  mutate(
    sex = factor(sex),
    quarter = factor(quarter))
```

```
health_sex = health_sex %>%
  drop_na() %>%
  select(
    health_score, quarter, sex
  ) %>%
  group_by(
    quarter, sex
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
pq <- ggplot(health_sex, aes(x = quarter, y = avg_score, group = sex)) +
  geom_line(aes(color = sex)) +
  geom_point(aes(color = sex)) + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean Health Score over time by sex")
pq
```

Trend in Mean Health Score over time by Sex



#Mean health score over time by race

```
health_race = practice_data %>%
  mutate(
    race = factor(race),
    quarter = factor(quarter))
```

```
health_race = health_race %>%
  drop_na() %>%
  select(
    health_score, quarter, race
  ) %>%
  group_by(
    quarter, race
  ) %>%
  summarise(
    avg_score = mean(health_score)
  )
```

```
pe <- ggplot(health_race, aes(x = quarter, y = avg_score, group = race)) +
  geom_line(aes(color = race)) +
  geom_point(aes(color = race)) + labs(x = "Quarter", y = "Mean Health Score", title = "Trend in Mean H
pe
```

