Barriers for measure, residue, corner and pairing methods in the lonely runner conjecture

Vico Bonfioli

18 October 2026. Preprint, not peer reviewed.

Abstract

We study which standard methods cannot prove the lonely runner inequality $\mathrm{ML}(v) \geq 1/(m+1)$ for positive integer speed vectors, and why. (i) At the target level, no integrable weight certifies a vector with $\mathrm{ML}(v) \leq 1/(m+1)$, so first-moment and Riesz-product certificates are blind to tight vectors; this is a formalized restatement of a lemma of Bedert. (ii) Five explicit families with 12 to 18 speeds defeat bounded-level base-point certificates of the kinds tested, and without a leave-one-out hypothesis no bound on denominators of lonely times depending only on m exists. (iii) For tight vectors of Goddyn–Wong type, $m+1$ divides no speed; a conjecture (T) extends this to all tight primitive vectors, and at a maximiser of a tight vector the denominator is $m+1$ times a divisor of the gcd of a corner pair. (iv) The doubling graph on the non-units in the window $[1, 2m]$ has an isolated vertex whenever $2m+1$ is composite and exceeds 9, hence no perfect matching. (v) Covering-radius bounds at the target value are false. Each statement carries a label; attributions to prior work are listed in Section 10. No progress on the conjecture is claimed.

1 Introduction

1.1 Setting

A speed vector is $v = (v_1, \ldots, v_m) \in \mathbb{Z}^m_{>0}$; there are $m+1$ runners, one of them stationary. Write $\|x\| = \mathrm{dist}(x, \mathbb{Z})$, $\mathrm{gap}_v(t) = \min_i \|v_i t\|$ and $\mathrm{ML}(v) = \max_{t \in \mathbb{R}} \mathrm{gap}_v(t)$ (a maximum, since gap_v is continuous and 1-periodic). The lonely runner conjecture (LRC) states $\mathrm{ML}(v) \geq 1/(m+1)$ for all v. A vector is *tight* if $\mathrm{ML}(v) = 1/(m+1)$; $(1, \ldots, m)$ is tight. The bad set of speed i at level δ is $B_i = \{t : \|v_i t\| < \delta\}$ and the good set is $G_\delta = \{t : \forall i,\ \|v_i t\| \geq \delta\}$; a time t is *lonely at level* δ if $t \in G_\delta$, that is if $\mathrm{gap}_v(t) \geq \delta$. A vector is *divisor-complete* (DC) if every $j \in \{2, \ldots, m+1\}$ divides some v_i. Throughout, m is the number of speeds; the literature cited uses n for m or for $m+1$, and each citation below is translated.

The inequality is known for $m \leq 12$ (Rosenfeld for eight runners [13]; Trakulthongchai for nine and ten [16], refereed; Sungkawichai and Trakulthongchai up to thirteen runners [14], unrefereed; Allikvere claims fourteen and fifteen runners [1], unrefereed, computer-assisted). The generic bounds are $1/(2m)$ by a union bound ([15, Prop. 1.1], [11, Section 5]) and $1/(2m) + m^{-5/3-o(1)}$ [3, Thm. 1.3].

The reduction studied in the accompanying archive reduces the conjecture for all m, given three finite statements, to the statement `FloorCruxU`: *if the floor holds for all smaller numbers of speeds, then for $m \geq 12$ every primitive v such that every leave-one-out subvector is primitive, v is DC, v admits a relation $\sum c_i v_i = 0$ with $c \in \{-1, 0, 1\}^m \setminus \{0\}$, and no*

Riesz-product certificate (Section 3) fires, satisfies $\mathrm{ML}(v) \geq 1/(m+1)$ (Lean: `FloorCruxU`, `universalFloor_of_four_atoms_U`). This statement is open. The methods below are those that were aimed at it.

1.2 Labels and provenance

VERIFIED: formalized in Lean 4 with Mathlib, zero `sorry`, axioms among `propext`, `Classical.choice`, `Quot.sound`, in the repository making the claim. All VERIFIED statements in this note are *carried over from the repository where they built; not rebuilt in this archive*: the build artefacts of the modules concerned are present there with timestamps matching their sources and contain no `sorry` (the modules holding the open statements (T), (T′), (T2) do contain `sorry`); the axiom prints of 1 October 2026 are recorded in `LEAN_MAP.md` and were not re-run. PROVED: complete written proof, reviewed adversarially, with conceptual abstract and zoom map, not formalized. HEURISTIC: computation or plausibility only; searches state their ground and termination status (RESOLVED if completed, UNRESOLVED if capped). CONJECTURE: no proof. FALSE: counterexample in hand. The strength of a result is the weakest label in its dependency chain.

1.3 Results

Item	Section	Statement	Label
Thm 2	3	If $\mathrm{ML}(v) \le \delta$ the good set is countable; no integrable weight can satisfy the certificate inequality at a tight vector	VERIFIED, carried over
Cor 3	3	No Riesz-product certificate fires on tight vectors	VERIFIED, carried over
Rem 4	3	Scope: what the barrier does not cover	HEURISTIC
Prop 5, 6	4	Escape lemma; missing divisor j gives $\|v_i/j\| \ge 1/j$ and a lonely time	VERIFIED, carried over (escape; weak form of the missing-divisor bound); strict form PROVED
Prop 7	4	(a) Times of positive gap of $v^{(Q)}$ have denominator $> Q$; (b) $\mathrm{ML}(v^{(Q)}) \ge 1/(m+1)$; so no bound $Q(m)$ on denominators of lonely times without leave-one-out primitivity	PROVED
Obs 8	4	Five families defeat bounded-level certificates	HEURISTIC (exact, finite)
Thm 9, Cor 10	5	Goddyn–Wong type vectors: $m+1$ divides no speed	VERIFIED, carried over
Prop 13	5	The implications (T′) ⇒ (T) and (T) ⇒ no tight primitive vector is DC; neither (T) nor (T′) is proved	PROVED (implications)
Prop 14	5	At a maximiser k/L of v with $\mathrm{ML}(v) = a/b < \frac{1}{2}$: $b \mid L$ and L/b divides the gcd of a corner pair	PROVED
Conj 12, 15	5	(T), (T′), (T2)	CONJECTURE
Prop 16	6	Isolated vertex in the doubling graph on the non-units of $[1, 2m]$ for composite $2m+1 > 9$	PROVED
Prop 18, Rem 19	7	Half-phase statement false; covering-radius target false	VERIFIED, carried over; cited computation

Theorem 2 and the escape lemma are known in substance (Tao, Bedert, Rosenfeld); Proposition 6 is known; Theorem 9 is, to the extent read, not stated in the literature, but its proof is a few lines. What is attributed and what is not is itemized in Section 10.

1.4 Conceptual abstract

The note tests five standard routes to the floor on the vectors where it is sharp. Averaging a fixed weight over the circle cannot certify a tight vector at the target level: its good set is a finite union of tent-function level sets, hence countable and without mass. Certificates of bounded denominator fail without a leave-one-out hypothesis, because a speed divisible by $\mathrm{lcm}(1, \dots, Q)$ vanishes at every time of denominator at most Q. Pairing by doubling fails because, for odd composite $2m+1$ above 9, the non-unit $2m+1-2p$, with p the least prime factor, has no doubling or halving partner in the window. Local corner analysis fails because the maximum at a corner depends on the residues of all other speeds. Covering-radius bounds fail because the shifted statement they imply is false. What remains is an explicit rational time or a classification of the tight vectors.

1.5 Zoom map for Theorem 2 and Corollary 3

L1.1 A level set of $\|at\|$ is countable (`nearestIntDist_eq_mem`).

L1.2 If $\mathrm{ML}(v) \le \delta$, then G_δ lies in the union of such level sets, hence is countable and null (`goodSet_countable`, `goodSet_null`).

L1.3 A nonnegative weight satisfies $\int F \le \sum_i \int_{B_i} F$ (`integral_le_sum_badSet`).

L1.4 Contrapositive: a strict inequality $\sum_i \int_{B_i} F < \int F$ implies $\delta < \mathrm{ML}(v)$ (`lt_ML_of_integral_unionBound`).

L1.5 Specialization to Riesz-product weights at $\delta = 1/(m+1)$ and non-vacuity on $(1, \dots, m)$ (`not_UDFires_of_tight`, `ML_consec_le`).

2 Methods against known theorems

Method	Strongest known statement it supports	Where it stops	Source
First moment on individual bad sets	$\mathrm{ML} \ge 1/(2m)$	ignores overlaps of bad sets; insensitive to phases, so cannot exceed the shifted infimum (Section 7)	[15], [11]
Weighted first moment, Riesz products	$\mathrm{ML} \ge 1/(2m) + m^{-5/3-o(1)}$; floor for dissociated sets	Thm 2: fixed integrable weights see no room at tight vectors	[3]
Atomic measures, rational witness times	floor when some $j \le m+1$ divides no speed; floor for $m \le 12$ by sieved computation	Prop 7, Obs 8: fixed-level checks are defeated	[15], [13], [16], [14]
Relation structure (Fourier on $\mathbb{Z}/q$, windows)	counterexample or tight $\Rightarrow$ integer relation of ℓ^1 norm $\le 2m+3$	tight vectors near $(1, \dots, m)$ satisfy it vacuously	[6], [2]
Finite checking, lattice width	counterexamples have bounded height; $n^{O(n^2)}$, $(n(n+1)/2)^{n-1}$	bounds not usable for all m	[15], [10], [7]
Covering radius	equals shifted loneliness (Henze–Malikiosis)	the shifted statement is false (Sec. 7)	[4]
Local analysis, Goddyn–Wong families	tight vectors of one-entry type are classified	transfers tightness, not a floor (Sec. 5)	[8], [17]
Doubling pairings	(internal to the covering framework)	isolated vertices (Sec. 6)	this note

3 Volume and smooth-weight certificates

For a nonnegative $F \in L^1[0,1]$ and a level δ consider

$$\sum_i \int_{[0,1]\cap B_i} F < \int_{[0,1]} F. \tag{1}$$

If $\mathrm{ML}(v) < \delta$, every t lies in some B_i, so $F \le \sum_i \mathbf{1}_{B_i} F$ pointwise and (1) fails. The weights $F = P^D$, $P(t) = \prod_i (1 - \cos 2\pi v_i t)$, are Riesz products; U_D *fires* if (1) holds for $F = P^D$

at $\delta = 1/(m+1)$. The case $F \equiv 1$ is the union bound of [15, Prop. 1.1]. For an arbitrary probability measure μ and closed bad sets the inequality $\int \Phi \, d\mu \geq 1$, $\Phi = \sum_i \mathbf{1}_{\{\|v_i x\| \leq \delta\}}$, at $\delta = \mathrm{ML}(v)$ is [3, Lemma 2.1].

Lemma 1 (Level sets)**.** *For $a \in \mathbb{Z} \setminus \{0\}$ and $\delta \in \mathbb{R}$, $\{t : \|at\| = \delta\} \subset \{(n \pm \delta)/a : n \in \mathbb{Z}\}$ is countable.*

Theorem 2. *Let $v \in (\mathbb{Z} \setminus \{0\})^m$, $\delta \in \mathbb{R}$ and $\mathrm{ML}(v) \leq \delta$. Then G_δ is countable, hence Lebesgue-null. Consequently, for every measurable $F \geq 0$ integrable on $[0,1]$,*

$$\int_{[0,1]} F \leq \sum_i \int_{[0,1] \cap B_i} F,$$

so (1) *implies $\delta < \mathrm{ML}(v)$.*

Proof. Let $t \in G_\delta$. Then $\mathrm{gap}_v(t) \geq \delta \geq \mathrm{ML}(v) \geq \mathrm{gap}_v(t)$, so $\mathrm{gap}_v(t) = \delta$. As gap_v is a minimum over finitely many terms, some i has $\|v_i t\| = \delta$. By Lemma 1, G_δ is a finite union of countable sets. Outside G_δ some i has $t \in B_i$, hence $F(t) \leq \sum_i \mathbf{1}_{B_i}(t) F(t)$ for almost every t; integrate over $[0,1]$. □

Label: VERIFIED, carried over (`goodSet_countable`, `goodSet_null`, `integral_le_sum_badSet`, `lt_ML_of_integral_unionBound`). The inequality for absolutely continuous measures is Bedert's Lemma 2.1 read with $\mu = F\,dt$; the countability of G_δ is elementary.

Corollary 3. *If $\mathrm{ML}(v) \leq 1/(m+1)$, in particular if v is tight, no U_D fires, $D \geq 0$. Conversely, if U_D fires then $1/(m+1) < \mathrm{ML}(v)$.*

Label: VERIFIED, carried over (`not_UDFires_of_tight`, `lt_ML_of_UDFires`); the hypothesis is not vacuous, $\mathrm{ML}(1, \ldots, m) \leq 1/(m+1)$ (`ML_consec_le`, `not_UDFires_consec`). The corollary is an immediate consequence of the cited lemma of Bedert.

Remark 4 (Scope). Theorem 2 concerns (1) at one fixed level and one fixed weight. It does not exclude a weight $F_{\delta'}$ chosen separately for each level $\delta' < \mathrm{ML}(v)$: the indicator of $\{\mathrm{gap}_v > \delta'\}$ satisfies (1) exactly for $\delta' < \mathrm{ML}(v)$, but constructing it requires knowing the lonely times. What is excluded is a weight given by a formula uniform in v, such as P^D, at the target level. Atomic measures are outside the theorem: a point mass at a rational time is the escape lemma (Proposition 5), and the uniform measure on $\{j/p\}$ with p a prime dividing no speed is [3, Lemma 5.3, Cor. 5.4]. Volume comparisons of geometry-of-numbers type (successive minima, covering radii bounded through volume) were not shown to reduce to (1); the claim that they do is not made. Label: HEURISTIC for the scope statements, as they delimit methods and are not theorems.

Obstruction for volume methods. A first-moment argument that uses only the measures of the individual bad sets gives $1/(2m)$, because each B_i has measure 2δ and the sum must reach 1; the factor two is the multiplicity with which bad sets overlap, which this argument discards. Sharper volume arguments must control the intersections $B_i \cap B_j$, and these depend on the starting phases of the runners, whereas the individual measures do not; since the shifted statement is false (Section 7), a volume argument can pass the exponent only by using the zero phase through the intersection pattern. At the target value itself the pattern is rigid: by Theorem 2 the bad sets of a tight vector cover the circle up to a null set, so a weight can gain nothing beyond equality.

Obstruction for smooth-weight certificates. A Riesz product or any other integrable weight turns the existence of a lonely time into a strict inequality between two Lebesgue integrals, and at a vector with $\mathrm{ML}(v) = 1/(m+1)$ the two integrals are forced by Theorem 2 to satisfy the opposite non-strict inequality, because the lonely set is countable and carries no mass. The mechanism is that gap_v is a minimum of finitely many tent functions, each of whose level sets is discrete; no choice of weight can see a set of measure zero. The obstruction persists off the tight set in the tested range: the five families of Table 1 have ML strictly above the target and $U_1, U_2 \geq 1$.

4 Residue and base-point certificates

Proposition 5 (Escape lemma)**.** *Let $q \geq 1$, $a \in \mathbb{Z}$, $\varepsilon \in \mathbb{R}$, $\delta > 0$. If for each i there are integers N_i, r_i with $v_i a = qN_i + r_i$ and $\delta \leq r_i/q + v_i\varepsilon \leq 1 - \delta$, then $\delta \leq \mathrm{ML}(v)$.*

Proof. At $t = a/q + \varepsilon$ we have $v_i t = N_i + r_i/q + v_i\varepsilon$, so $\|v_i t\| \geq \delta$ for all i. □

Label: VERIFIED, carried over (`ML_ge_of_escape`). With $\varepsilon = 0$ this is the definition of a proper tuple in [16, Def. 3], the witness-time ansatz of [13]; the extension to $\varepsilon \neq 0$ is a routine generalization.

Without a bound on q the statement is empty: $\delta \leq \mathrm{ML}(v)$ holds if and only if some (q, a, ε) works, with $q = 1$ (`lonely_iff_escape`, VERIFIED, carried over), and for $\delta < \mathrm{ML}(v)$ one can take $\varepsilon = 0$ (`rational_point_of_lt`, VERIFIED, carried over). The window lemma at levels $m+1 < q \leq 2m+2$ and the certificates used in the archive are instances (`window_via_escape`, `famF_via_escape`, `lemmaP_via_escape`). All content lies in the bound on q.

Proposition 6 (Missing divisor)**.** *If $2 \leq j \leq m+1$ and j divides no v_i, then $\|v_i/j\| \geq 1/j$ for all i, so $\mathrm{ML}(v) \geq 1/j$ with witness $t = 1/j$. In particular $\mathrm{ML}(v) > 1/(m+1)$ if $j \leq m$, and $\mathrm{ML}(v) \geq 1/(m+1)$ if $j = m+1$.*

Label: The bound $\|v_i/j\| \geq 1/j$ is PROVED in one line: $v_i \bmod j \in [1, j-1]$. The Lean lemma `witness_of_missing_divisor` (VERIFIED, carried over) has the signature $j \leq (k+2)\min(r, j-r)$ with $r = v_i \bmod j$ for $k+1 = m$ speeds, that is $\|v_i/j\| \geq 1/(m+1)$, the weak form; the strict form is not stated there, and no other part of that Lean statement was re-read in this revision. Known: [15, Lemma 5.1(i)]; also the survey [11] proves that the conjecture holds when some $k \leq n+1$ divides no speed, and [17, Lemma 2.3] is the case $t = 1/r$ for one-entry modifications. Not claimed as new.

Consequently a tight vector satisfies: every $j \in [2, m]$ divides some speed, since otherwise $\mathrm{ML}(v) \geq 1/j > 1/(m+1)$. This is why the crux is restricted to DC vectors.

Proposition 7 (Denominators of lonely times)**.** *Let $m \geq 2$, $Q \geq 1$, $L = \mathrm{lcm}(1, \ldots, Q)$ and $v^{(Q)} = (1, L, 2L, \ldots, (m-1)L) \in \mathbb{Z}^m_{>0}$. If $t = a/q \in \mathbb{Q}$ in lowest terms has $\mathrm{gap}_{v^{(Q)}}(t) > 0$, then $q > Q$. (b) $\mathrm{ML}(v^{(Q)}) \geq 1/(m+1)$, witnessed by $t = c/(L(m+1))$ for any integer c with $L \leq c \leq Lm$ and $c \equiv 1 \pmod{m+1}$; such c exists. Consequently, for every function $Q(m)$ there is a vector with m speeds that has lonely times at level $1/(m+1)$ but none of denominator at most $Q(m)$.*

Proof. If $q \leq Q$ then $q \mid L$, so $Lt \in \mathbb{Z}$ and $\|Lt\| = 0$. The speed L occurs in $v^{(Q)}$ because $m \geq 2$, so $\mathrm{gap}(t) = 0$. Hence $q > Q$ whenever $\mathrm{gap}(t) > 0$.

(b) The interval $[L, Lm]$ contains $L(m-1)+1$ integers. For $L=1$ it contains $1 \equiv 1$. For $L \geq 2$ it contains at least $m+1$ consecutive integers, hence one with $c \equiv 1 \pmod{m+1}$. Put $t = c/(L(m+1))$. Since $L \leq c \leq Lm$, $t \in [\frac{1}{m+1}, \frac{m}{m+1}]$ and $\|t\| \geq \frac{1}{m+1}$ (speed 1). For $1 \leq j \leq m-1$ the speed jL gives $jLt = jc/(m+1)$; $\gcd(c, m+1) = 1$ and $m+1 \nmid j$, so $m+1 \nmid jc$ and $\|jc/(m+1)\| \geq \frac{1}{m+1}$. Hence $\mathrm{gap}(t) \geq \frac{1}{m+1}$, and by (a) its denominator is $> Q$. For the consequence take $Q = Q(m)$. □

Label: PROVED. Idea: a speed divisible by every small denominator is invisible to every small-denominator test, because it sends all rational times of small denominator to 0 *mod* 1*. Zoom map: L1.1 reduction of $q \leq Q$ to $q \mid L$ (arithmetic, no lemma); L1.2 positivity of the gap forces $q > Q$; L1.3 existence of c and the bound* $\mathrm{gap}(t) \geq \frac{1}{m+1}$ *at $t = c/(L(m+1))$. For $Q \geq 2$ the subvector $(L, 2L, \ldots, (m-1)L)$ obtained by deleting the speed* 1 *has* $\gcd = L > 1$*, so $v^{(Q)}$ is outside the class of the crux, whose members have primitive leave-one-out subvectors. No prior statement was located in the literature read.*

Rationality of maximizers. The gap is piecewise linear with slopes $\pm v_i \neq 0$, so it has no flat piece and each local maximum is a vertex of its graph; every vertex is the solution of a linear equation with integer coefficients, hence rational. So every maximiser of gap_v is rational, and for $v = v^{(Q)}$ every maximiser, which has positive gap, has denominator $> Q$ by the proposition. This is not needed for the proposition.

Obstruction for residue-condition methods. A test at denominator q sees only the residues $v_i \bmod q$ of the speeds, and a speed with residue 0 at every level up to Q is invisible to all of them; Proposition 7 makes this exact, and the families of Table 1 realize it approximately inside the class of the crux by a few tall speeds (up to 4943) whose residues complete a covering of the small levels. The mechanism is that the residues of a few free tall speeds can be chosen independently modulo each small q while the maximal loneliness stays above the target; this explanation is consistent with the table and is not proved, label HEURISTIC.

4.1 Level drift

Let $\mathrm{TL}(Q)$ denote the statement: for $m \geq 12$ and every class member v of the crux, there are $q \leq Q(m)$, a unit a modulo q, $\sigma = \pm 1$ and $\varepsilon \geq 0$ such that Proposition 5 applies at $t = \sigma a/q + \varepsilon$ with the first-order arc condition on the non-multiples of q and a condition on the multiples of q through ε. The certificate kinds named in the table (WC, window points, window levels, exact multiple scan, first order) are the variants implemented in `cert_m` of `code/reduction/mdr`.

Observation 8. For each certificate kind and level range in Table 1, the stated class member has no certificate of that kind in that range.

Label: HEURISTIC. Each row is a finite exact computation (exact integer arithmetic; ML *recomputed by two independent programs, see Section 8), not checked in Lean. Class membership used U_1, U_2 computed in double precision with an a posteriori bound of 10^{-12}; the listed margins exceed it. Termination: rows 1 to 5 RESOLVED (exhaustive over the stated levels); row 6 UNRESOLVED.*

Last row of the table: a time-limited hunt of 120 s per m (one process, seed 1) performed 14 686, 8 962, 4 141, 2 859 evaluations for $m = 18, 20, 22, 24$, reached best levels $29, 37, 34, 37$

Certificate kind and range	Class member without certificate	U_1, U_2
WC, $q \le m+1$	$[1,4,5,6,7,9,11,13,40,60,238,2731]$, $m=12$, ML $=14/121$	1.003918, 1.001345
Window points, $q \le 2m+2$	$[1,3,4,5,7,9,11,13,40,48,100,1079]$, $m=12$, ML $=11/84$ (first certified at $q=13$ by WC)	1.002573, 1.000429
Window levels, exact, $q \le 2m+2$	$[1,5,6,7,8,9,11,13,17,19,23,25,29,31,366,1680]$, $m=16$, ML $=168/1697$; first level 35	1.483, 1.715
All levels, first order and exact, $q \le 2m+1$	$[1,3,4,5,7,11,13,17,19,23,31,70,74,116,180,272,3986,4943]$, $m=18$; ML $=28/191$; first point at $q=38$	1.007002, 1.003326
All levels, first order, $q \le 2m+2$	$[1,3,4,5,7,11,13,17,19,23,31,38,70,74,116,180,272,4422]$, $m=18$; ML $=30/203$; first level 39	1.024583, 1.014631
Exact multiple scan, $q \le 2m+2$, $m=18,\dots,24$	none found in time-limited hunts; UNRESOLVED (not exhaustive)	n/a

Table 1: Families defeating bounded-level certificates (script `code/obstructions/repro_04.sh`). The window-levels row also has a first-order version, first level 41, on $[1,3,4,5,7,8,9,11,12,13,17,19,23,29,31,37,224,1800]$, $m=18$. The $m=12$ vector in row 2 has least witness $(4,31)$.

against the cutoffs $38,42,46,50$, and produced no family; an earlier run of 480 s per m was reported with the same outcome and is not repeated here. Both are UNRESOLVED and support no statement about other budgets or seeds.

Positive ground, regenerated by `mdr enumw`: the window-only first-order version had no failure on the exhaustive supersets (DC, leave-one-out primitive, not strongly free) for $m=6$, $H \le 50$ (15 890 700 sets, 3 202 695 in the class); $m=8$, $H \le 24$ (735 471; 119 915); $m=10$, $H \le 30$ (30 045 015; 3 327 077); $m=12$, $H \le 26$ (9 657 700; 934 023); $m=14$, $H \le 24$ (1 961 256; 107 976), each box RESOLVED; and none on the 1067 census families (lists in `data/obstructions/census`) and the 147 adversarial families (`mdr cert`, certificate level at most $2m+2$ for all). The census families are residuals of boxes and are not independent of the boxes. The failures above thus appear only at larger m or height. The role of the condition "no U_D fires" in this ground is not understood.

5 Local corners: the Goddyn–Wong mechanism and statement (T)

Corners. ML(v) is attained at a time k/D with $D \in \{2v_i, v_i+v_j, |v_i-v_j|\}$ and $0 \le k < D$ ([11, Section 2]; the numerators are not restricted to residues coprime to D). A local argument fixes such a pair and controls the maximum from it; the remaining $m-2$ speeds enter only through their residues modulo D. For ML$(v) < \frac{1}{2}$ every maximizer is such a corner, of a rising and a falling tent (Proposition 14).

For multipliers $\mu = (\mu_1,\dots,\mu_{n-1}) \in \mathbb{N}^{n-1}$ write $[n-1]_\mu = (\mu_1 \cdot 1,\dots,\mu_{n-1}(n-1))$, with $n = m+1$. The *GCD condition* is: for each $r \in [n-1]$ with $\mu_r \ge 2$ and each b with $n-r \le b \le \mu_r(n-r)-1$, $\gcd(r,b) > 1$. Goddyn and Wong prove ML$([n-1]_\mu) = 1/n$ under this condition [8, Thm. 3.1], quoted as [17, Thm. 5.1]; for one accelerated entry it is also necessary [8, Thm. 2.3], quoted as [17, Thm. 1.1]. Zhang classifies all one-entry modifications $[n-1]_{r\to w}$, $w > n-1$: tight if and only if $2r > n-1$, $r \mid w$ and the GCD condition holds with $\mu = w/r$, or $(n,r,w) \in \{(5,2,7),(6,2,9)\}$ [17, Thm. 1.2].

Theorem 9. *If μ satisfies the GCD condition, then n divides no speed $\mu_r r$, $1 \le r \le n-1$.*

Proof. If $\mu_r = 1$ then $0 < r < n$. Let $\mu_r \ge 2$ and suppose $\mu_r(n-r) \ge n$. The interval $[n-r, \mu_r(n-r)-1]$ has length $\mu_r(n-r)-(n-r) \ge r$, so it contains r consecutive integers, one of which is $\equiv 1 \pmod r$ and hence coprime to r; this contradicts the GCD condition. So $0 < \mu_r(n-r) < n$. Now $\mu_r r + \mu_r(n-r) = \mu_r n$, so $n \mid \mu_r r$ would give $n \mid \mu_r(n-r)$, impossible for $0 < \mu_r(n-r) < n$. □

Label: VERIFIED, carried over (`not_dvd_speed`; axioms recorded as `propext`, `Quot.sound`). A statement of this form was not located in the sources read; the original proof of [8, Thm. 3.1] was not available for this audit and may contain the same step. The published argument obtains a sharper size bound; only the weaker one is used.

Corollary 10. *Under the GCD condition with $\mu_1 = 1$ (forced, since $\gcd(1,b)=1$), the vector $[n-1]_\mu$ is not DC: $j = n = m+1$ divides no speed. The same holds for the vectors $(1,3,4,7)$, $(1,3,4,5,9)$, $(1,4,5,6,7,11,13)$, $(1,2,3,4,5,7,12)$, $[1..11]\cup\{13,24\}$, $[1..17]\cup\{19,36\}$ and for the two-entry doubling instance $[1..73]\setminus\{70,72\}\cup\{140,144\}$ with $n=74$ [17, Cor. 5.2 and the example following Thm. 5.1].*

Label: VERIFIED, carried over for $[n-1]_\mu$ under the GCD condition (`not_divisorComplete`). For the listed vectors: HEURISTIC, an exact computation (script `code/obstructions/gwcheck.py`) of ML $= 1/n$ *and of the non-divisibility, RESOLVED.*

Remark 11. A vector built by the Goddyn–Wong construction is excluded from the class of the crux, since the proof of its tightness shows $m+1 \nmid v_i$. The mechanism uses three inputs: the baseline $[n-1]$, the identity $\mu_r r \equiv -\mu_r(n-r) \pmod n$ valid because the new speed is an integer multiple of an existing entry, and the GCD hypothesis reused through a size bound. A general class member has no such decomposition, and the generalizations in the literature (Zhang's one-entry classification, the multi-swap theorem) keep the multiplicative form. The route therefore gives no lower bound for divisor-complete vectors. The converse of the multi-acceleration theorem is not true in general [17, Question 9.3].

Obstruction for local-corner methods. The maximum of gap_v is a corner determined by two speeds, but whether the corner is a maximum depends on the residues of every other speed modulo the corner's denominator, and these residues carry no structure that a pair-local argument can use. In the Goddyn–Wong families the missing control is supplied by a global identity, namely that every speed is a multiple of an entry of $[n-1]$ and the sum $\mu_r r + \mu_r(n-r)$ is a multiple of n; for a vector without this decomposition the maximizer still sits at denominator $m+1$ in every tight case computed (Section 8), but no pair of speeds identifies which speed would carry the divisor $m+1$ if it existed. This is why a direct local proof of (T) below gets no purchase, label HEURISTIC.

5.1 Statement (T)

Conjecture 12 ((T), (T′))**.** *Let $v \in \mathbb{Z}^m_{>0}$ with $\gcd(v)=1$ and $\mathrm{ML}(v) = 1/(m+1)$. Then (T) $m+1$ divides no v_i; (T′) every t with $\mathrm{gap}_v(t) = \mathrm{ML}(v)$ has the form $a/(m+1)$, $a \in \mathbb{Z}$.*

Label: CONJECTURE. Provenance: observation on the tight vectors of Goddyn–Wong type and on exhaustive small boxes. Evidence: Section 8, exhaustive primitive search for $m = 3, \dots, 8$ in boxes finds the tight vectors listed there, all satisfying (T) and (T′). Consequence: by Proposition 6 a tight vector has every $j \in [2, m]$ dividing a speed, so (T) is equivalent to the statement that no tight primitive vector is DC. Lean statements `TightNoDvd`, `TightMaximisersAtMp1` *are defined without proof;* `tightNoDvd_of_strong` *has a* `sorry`.

Proposition 13. *Let $v \in \mathbb{Z}^m_{>0}$ and suppose $m+1 \mid v_i$ for some i. Then $\operatorname{gap}_v(a/(m+1)) = 0 < \mathrm{ML}(v)$ for every $a \in \mathbb{Z}$. Consequently (T′) implies (T), and (T) implies that no tight primitive vector is DC.*

Proof. Since $v_i/(m+1) \in \mathbb{Z}$, $v_i \cdot a/(m+1) \in \mathbb{Z}$ and $\operatorname{gap}_v(a/(m+1)) \le \|v_i t\| = 0$. Also $\mathrm{ML}(v) > 0$, as $\operatorname{gap}_v(t) > 0$ at any irrational t. Assume (T′) and let v be primitive and tight with $m+1 \mid v_i$. A maximizer t exists, since gap_v is continuous and 1-periodic; by (T′), $t = a/(m+1)$, so $\operatorname{gap}_v(t) = 0 \neq \mathrm{ML}(v)$, a contradiction. Hence (T) holds. A DC vector has $m+1 \in \{2, \dots, m+1\}$ dividing some v_i, which (T) excludes for tight primitive v. □

Label: PROVED (the implications only; neither (T) nor (T′) is proved). Idea: a speed divisible by $m+1$ vanishes at every candidate time of denominator $m+1$, so such a time cannot be a maximizer. Zoom map: L1.1 a divisible speed has gap 0 at $a/(m+1)$; L1.2 a maximizer exists and has positive gap; L1.3 divisor-completeness supplies the divisible speed. The proof does not use $\gcd(v) = 1$; that hypothesis matters for the truth of (T′), since the vector (2), $m = 1$, is tight and non-primitive and its maximizer $1/4$ is not in $\frac{1}{2}\mathbb{Z}$. Not formalized.

Proposition 14 (Tight corner)**.** *Let $v \in \mathbb{Z}^m_{>0}$ with $\mathrm{ML}(v) = a/b < \frac{1}{2}$ in lowest terms, and let t^* be a maximizer of gap_v. Then $t^* = k/L$ in lowest terms is rational, and there are indices i, j with $v_i t^* \equiv a/b$ and $v_j t^* \equiv -a/b \pmod 1$ such that*

$$b \mid L, \qquad L \mid b \gcd(v_i, v_j), \qquad L \mid v_i + v_j.$$

In particular $L = bd$ with $d \mid \gcd(v_i, v_j)$. If v is tight and $m \ge 2$ then $b = m+1$, so (T′) holds at t^* if and only if $d = 1$.

Proof. L1: (a) at a maximizer some speed sits at $+a/b$ and some at $-a/b$; (b) the two congruences give the divisibilities.

(a) Put $\delta = a/b$ and $S = \{i : \|v_i t^*\| = \delta\}$, nonempty since $\operatorname{gap}_v(t^*) = \delta$. As $0 < \delta < \frac{1}{2}$, each $i \in S$ has $v_i t^* \equiv \sigma_i \delta \pmod 1$ with a unique $\sigma_i \in \{+1, -1\}$. Suppose $\sigma_i = +1$ for all $i \in S$. For small $\varepsilon > 0$, $\|v_i(t^* + \varepsilon)\| = \delta + v_i \varepsilon$ for $i \in S$, and $\|v_i(t^* + \varepsilon)\| > \delta$ for $i \notin S$ by continuity. Then $\operatorname{gap}_v(t^* + \varepsilon) > \delta = \mathrm{ML}(v)$, which is absurd. The case $\sigma_i = -1$ for all $i \in S$ is the same with $t^* - \varepsilon$. So there are $i, j \in S$ with $\sigma_i = +1$, $\sigma_j = -1$. Then $v_i t^* = N_i + a/b$ and $v_j t^* = N_j - a/b$ with $N_i, N_j \in \mathbb{Z}$; in particular $t^* \in \mathbb{Q}$, say $t^* = k/L$ with $\gcd(k, L) = 1$.

(b) Adding, $(v_i + v_j)k/L = N_i + N_j \in \mathbb{Z}$, so $L \mid (v_i + v_j)k$ and $L \mid v_i + v_j$. Multiplying $v_i k/L = N_i + a/b$ by bL gives $bv_i k = L(bN_i + a)$. Hence $L \mid bv_i k$, so $L \mid bv_i$; and reducing modulo b gives $b \mid La$, so $b \mid L$ because $\gcd(a, b) = 1$. In the same way $bv_j k = L(bN_j - a)$ gives $L \mid bv_j$. Thus $L \mid \gcd(bv_i, bv_j) = b \gcd(v_i, v_j)$, and $L = bd$ gives $d \mid \gcd(v_i, v_j)$.

For tight v and $m \ge 2$, $\mathrm{ML}(v) = 1/(m+1) \le \frac{1}{3} < \frac{1}{2}$, so $a = 1$, $b = m+1$. (T′) at t^* says $L \mid m+1$; with $m+1 \mid L$ this is $L = m+1$, that is $d = 1$. □

Label: PROVED. Idea: at a maximizer one runner sits exactly at distance $+\mathrm{ML}$ *ahead of an integer and another exactly* ML *behind (the two residues* $\pm\mathrm{ML}$ *are distinct because* $\mathrm{ML} < \frac{1}{2}$*), otherwise the time could be moved to raise the gap; clearing the denominator of* ML *turns the two corner equations into congruences that bound the denominator of the maximizer by* b *times a common divisor of the corner pair. Zoom map: L1.1 opposite signs at a maximizer; L1.2 divisibilities from the two corner equations. That a maximizer lies at a time* k/D *with* D *a sum of two speeds is the vertex lemma [5, Lemma 2.2], [11, Section 2]; the divisibilities* $b \mid L$ *and* $L \mid b\gcd(v_i, v_j)$ *were not located in the sources read. For tight* v *with* $m+1 \mid v_l$ *and* $d = 1$ *at a maximizer,* $v_l t^* \in \mathbb{Z}$*, which is impossible; so a tight primitive vector with a speed divisible by* $m+1$ *would have* $d \geq 2$ *at every maximizer. Worked example:* $v = (1, 2, 4, 6)$*,* $\mathrm{ML} = 1/4$*,* $t^* = 3/8$*,* $L = 8$*,* $b = 4$*,* $d = 2$*, corner pair* $(6, 2)$ *with* $6t^* \equiv \frac{1}{4}$*,* $2t^* \equiv -\frac{1}{4}$*,* $L \mid v_i + v_j = 8$*,* $L \mid b\gcd(6, 2) = 8$ *(not tight:* $\mathrm{ML} = 1/4 > 1/5$*). Tight non-primitive example:* $v = (2, 4)$*,* $m = 2$*,* $\mathrm{ML} = 1/3$*, with maximizers* $1/6, 1/3, 2/3, 5/6$*; at* $t^* = 1/6$ *the corner pair is* $(2, 4)$*,* $L = 6$*,* $b = 3$*,* $d = 2$*. So* $d > 1$ *occurs, and the hypothesis* $\gcd(v) = 1$ *is needed for (T′). Not formalized.*

5.2 The family E_m and a related floor

For $E_m = (1, \ldots, m-1, m(m+1))$, $m \geq 3$, $\mathrm{ML}(E_m) = (m+1)/(m^2+m+1)$ (the case $s = m+1$ of [9, Thm. 5.1], reproved in Lean with a different argument, `ML_Em_m`, VERIFIED, carried over). Hence $(2m+1)\mathrm{ML}(E_m) > 2$ for $m \geq 3$ (`thmB_gt`, VERIFIED, carried over).

Conjecture 15 ((T2))**.** *If* v *is primitive and* $m+1$ *divides some* v_i*, then* $\mathrm{ML}(v) \geq 2/(2m+1)$*.*

Label: CONJECTURE. Evidence: in the exhaustive boxes of Section 8, the minimum of ML *over primitive DC vectors equals* $2/(2m+1)$ *for* $m = 3, \ldots, 7$*, attained for example by* $(1, 3, 4)$*, and* E_m *is not a minimizer. Consequence: every primitive vector in which* $m+1$ *divides a speed satisfies the floor with strict inequality. The vector* $(1, 5, 6, 11, 16, 17)$*,* $m = 6$*, has* $\mathrm{ML} = 5/33 < 2/13$*, but no speed is divisible by* 7*, so it lies outside the class. A statement that* E_m *is the unique minimizer is false.*

6 Doubling pairings

In the covering formulation a lonely time is ruled out by exhibiting bad sets that cover the window of integers $[1, 2m]$ for the modulus $q = 2m+1$; the smallest such sets are pairs. For q odd let $N = \{v \leq 2m : \gcd(v, q) > 1\}$ be the non-units in the window. The doubling graph on $[1, 2m]$ has edges $\{v, 2v\}$ with $2v \leq 2m$; it lives on the integers of the window, not on residues modulo q.

Proposition 16 (Chain parity)**.** *Let* $q = 2m+1 \geq 3$ *be odd,* $N = \{v \in [1, 2m] : \gcd(v, q) > 1\}$*, and let* Γ *be the graph on* N *with edges* $\{v, 2v\}$*,* $2v \leq 2m$*. If* q *is composite and* $q \neq 9$*, then* $s = q - 2p$*, with* p *the least prime factor of* q*, is an isolated vertex of* Γ*; in particular* Γ *has no perfect matching. If* q *is prime then* $N = \emptyset$*, and if* $q = 9$ *then* $N = \{3, 6\}$ *is a single edge. Hence every component of* Γ *has an even number of vertices if and only if* q *is prime or* $q = 9$*.*

Proof. Let q be composite and p its least prime factor. Then $q = pr$ with $r \geq p \geq 3$, so $q \geq 3p$ and $q \geq p^2$. Put $s = q - 2p$. Then s is odd, $p \leq s \leq q - 6 \leq 2m$, and $\gcd(s, q) = \gcd(2p, q) =$

$p > 1$, so $s \in N$. Being odd, s has no neighbour $s/2$. Its other possible neighbour $2s$ lies in $[1, 2m]$ if and only if $2s \le q - 1$, that is $q \le 4p - 1$. In that case $p^2 \le q \le 4p - 1$ forces $p \le 3$, hence $p = 3$ and $q \le 11$ is an odd composite multiple of 3, so $q = 9$. Thus for $q \ne 9$ the vertex s is isolated, a component with one vertex. If q is prime no $v \in [1, 2m]$ has $\gcd(v, q) > 1$. If $q = 9$, $N = \{3, 6\}$ and Γ is the edge $\{3, 6\}$. An isolated vertex is a component of odd size and admits no perfect matching. □

Label: PROVED. Idea: for composite $q > 9$ the non-unit $q - 2p$ is odd, so it has no halving partner, and its double exceeds the window, so it has no doubling partner; it is an isolated vertex of the doubling graph on the non-units of the window. Zoom map: L1.1 $q - 2p$ is a non-unit ($\gcd = p$*); L1.2 it is odd; L1.3 its double exceeds $2m$ unless $q = 9$. Independent check: brute-force chain lengths (the components of Γ) for odd $q \le 20001$, no mismatch (Section 8). Not formalized.*

Remark 17. The companion covering paper classifies every bad pair with non-unit endpoints as a doubling pair or a pair with $a + b \in \{2m + 1, 2m + 2\}$ (Lean-verified for the dichotomy; carried over), and pairs with $a + b = 2m + 1$ and both endpoints non-units do not occur. For $q \in \{25, 27, 33, 35, 49\}$ no non-unit bad pair with $a + b = 2m + 2$ was found in earlier computations, which are not regenerated here (HEURISTIC). If the same holds for all composite q, Proposition 16 implies that a partition of the window into bad pairs cannot cover the non-units, so parts of size at least three are forced; this implication is conditional and carries the label HEURISTIC.

Obstruction for doubling methods. A pairing of the window by $\{j, 2j\}$ treats the integers as a union of doubling chains, and a component with an odd number of vertices leaves one vertex unmatched. For a composite modulus $q > 9$ the vertex $q - 2p$ is an odd non-unit whose double exceeds the window, so it is an isolated vertex of the doubling graph on the non-units; this is the whole mechanism, and it does not depend on any property of the speeds. Passing to the cyclic group $\mathbb{Z}/q\mathbb{Z}$, where doubling is a permutation, does not repair it, since the bad sets of the covering formulation are multiples of j at modulus j and do not realize the wrap-around partner (earlier computations, not regenerated; HEURISTIC).

7 Covering radius and the zero phase

For phases $s \in \mathbb{R}^m$ put $\mathrm{gap}_{v,s}(t) = \min_i \|s_i + v_i t\|$. Henze and Malikiosis identify $\mathrm{ML}(v)$ with $\frac{1}{2} - \frac{1}{2}\kappa(Z(v))$ and the shifted supremum $\sup_t \mathrm{gap}_{v,s}(t)$, minimized over s, with $\frac{1}{2} - \frac{1}{2}\mu(Z(v))$, where κ is the first minimum and μ the covering radius of the lonely runner zonotope $Z(v)$ [4, Prop. 1.7]. Blanco, Criado and Santos show by exact computation that the shifted infimum for $v = (1, \ldots, n)$ is below $1/(n + 1)$ for $5 \le n \le 17$ speeds, for instance $15/94 < 1/6$ for $(1, 2, 3, 4, 5)$ [4, Prop. 5.11, 5.14, Thm. 1.8]; Poliakova shows that the defect grows linearly and is positive for $n \ge 95$ [12, Thm. 1, 2].

Proposition 18. *There is no $t \in \mathbb{R}$ with $\|t\| \ge \frac{1}{7}$, $\|2t - \frac{1}{2}\| \ge \frac{1}{7}$, $\|3t - \frac{1}{2}\| \ge \frac{1}{7}$, $\|4t - \frac{1}{2}\| \ge \frac{1}{7}$, $\|6t - \frac{1}{2}\| \ge \frac{1}{7}$ and $\|9t\| \ge \frac{1}{7}$. That is, for speeds $(1, 2, 3, 4, 6, 9)$ and phases $(0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0)$ the shifted lonely runner bound for seven runners fails.*

Label: VERIFIED, carried over (`halfShift_not_lonely`). Computed value (HEURISTIC, exact): the supremum of the shifted gap is $3/22$, attained at $t = 7/22$, while the zero-phase

value for the same speeds is ML $= 1/5$ *(`code/obstructions/shift_grid.py`, exact rational scan of all* $t = p/q$*,* $q \le 400$*, and independent recomputation of* ML*).*

Remark 19. By the dictionary above, a bound $\mu(Z(v)) \le (n-1)/(n+1)$ for all v with n speeds is false for $5 \le n \le 17$ (equivalent to the failure of the shifted bound, cited computation, label HEURISTIC as a computation by others, reproduced for one instance by the exact scan above). The covering-radius route of [10], whose Proposition 2.7 bounds $\mu(C) \le \lambda_1(C-C) + \mu(\pi(C), \pi(\mathbb{Z}^d))$ for a convex body C, can therefore be used for the conjecture only through κ, never as a bound at the target value for μ. [10, Lemma 2.5] (if $|C \cap \mathbb{Z}^d| > \ell^d$ then $\lambda_1(C-C) \le 1/\ell$) is a statement on an arbitrary convex body; its conclusion is unchanged by translating C, and the zero phase enters only through the lattice-point count [10, Cor. 2.3].

Obstruction for covering-radius methods. Any argument whose steps remain valid when the zero phase is replaced by a phase vector in $\{0, \frac{1}{2}\}^m$ proves a false statement for $v = (1, 2, 3, 4, 6, 9)$, so a proof of the conjecture must use the zero phase, that is the common starting point of all runners. The covering radius is a translation-invariant quantity and is the right invariant for the shifted problem only; the target quantity is the first minimum. The mechanism is the failure of the shifted statement, proved above for one explicit instance.

8 Evidence and regeneration

All commands are run from the archive root; searches report RESOLVED or UNRESOLVED. Cases counted are independent vectors in distinct boxes.

Item	Regeneration command	Status
ML of the five rows of Table 1, of E_{14} and of $(1, 2, 3, 4, 6, 9)$	`mlcheck` (Rust) via `code/obstructions/repro_04.sh`; `python3 code/obstructions/mlexact.py` (independent, Python, all numerators, exact integers)	RESOLVED; both programs agree on 14/121, 11/84, 168/1697, 28/191, 30/203, 1/5
Certificate levels, U_1, U_2	`code/obstructions/repro_04.sh`	RESOLVED
Census and boxes	`mdr enumw M H`, `mdr cert 80 FILE` (see `code/obstructions/README.md`)	RESOLVED
Last row of Table 1	`MDR_WX=1 mdr hunt2` (README)	UNRESOLVED
Tight vectors, (T), (T′)	`python3 code/obstructions/tightscan.py`	see below
Minimum of ML on DC vectors (T2)	`python3 code/obstructions/t2box.py`	see below
Goddyn–Wong type vectors	`python3 code/obstructions/gwcheck.py`	RESOLVED
Chain parity	`python3 code/obstructions/chainparity.py 20001`	RESOLVED, $q \le 20001$
Shifted supremum	`python3 code/obstructions/shift_grid.py`	RESOLVED

Tight vectors. For each m and bound H, all primitive $v \in [1, H]^m$ with distinct entries such that every $j \in [2, m]$ divides a speed (necessary for tightness by Proposition 6) were enumerated and ML computed exactly, together with all maximizing times.

m	H	vectors enumerated	tight vectors found
3	60	17 432	$(1, 2, 3)$
4	40	47 078	$(1, 2, 3, 4)$, $(1, 3, 4, 7)$
5	30	63 880	$(1, \ldots, 5)$, $(1, 3, 4, 5, 9)$
6	24	57 111	$(1, \ldots, 6)$
7	20	23 212	$(1, \ldots, 7)$, $(1, 2, 3, 4, 5, 7, 12)$, $(1, 4, 5, 6, 7, 11, 13)$
8	16	4 481	$(1, \ldots, 8)$

In every box, no tight vector has a speed divisible by $m + 1$, and every maximizing time of every tight vector lies in $\frac{1}{m+1}\mathbb{Z}$. Each box is RESOLVED; the boxes are small (toy only), the heights for $m \geq 6$ are at most 24, and the boxes for different m are independent instances (different numbers of speeds). The evidence for (T) and (T′) is confined to this ground and says nothing for $m \geq 9$ or beyond these heights; both remain CONJECTURE.

Minimum on divisor-complete vectors. In the same boxes (primitive, DC including $j = m + 1$) the minimum of ML is $2/7$ ($m = 3$, $H \leq 60$; 10 946 vectors), $2/9$ ($m = 4$, $H \leq 40$; 27 600), $2/11$ ($m = 5$, $H \leq 30$; 46 316), $2/13$ ($m = 6$, $H \leq 24$; 28 835), $2/15$ ($m = 7$, $H \leq 20$; 12 673), each equal to $2/(2m + 1)$, and strictly below $\mathrm{ML}(E_m) = 4/13, 5/21, 6/31, 7/43, 8/57$. Each box is RESOLVED; the boxes are small, and the evidence is HEURISTIC and toy only: no statement is made for $m \geq 8$. The tight and DC counts in neighbouring boxes are not independent.

9 Open statements

- (T) and (T′), Conjecture 12. Evidence: the exhaustive boxes above and every tight vector of Goddyn–Wong type (Corollary 10). Consequence: (T) is equivalent to the statement that no tight primitive vector is DC (Conjecture 12 and Proposition 6); (T′) implies (T) (Proposition 13).
- (T2), Conjecture 15. Evidence: equality $2/(2m+1)$ is the minimum on DC vectors in the boxes for $m = 3, \ldots, 7$. Consequence: strict floor for primitive vectors with $(m + 1) \mid v_i$.

Whether the class of the crux contains a tight vector is open. Neither statement is used elsewhere in this note.

10 Attribution and prior art

Each item was compared with the sources cited; “known” means the statement or its proof is in the cited place.

Item	Status	Reference location
Thm 2, inequality for weights $F \geq 0$	known	[3, Lemma 2.1] with $\mu = F\,dt$ (closed bad sets, ML $= \delta$); union bound [15, Prop. 1.1]
Thm 2, countability of G_δ; formalization	ours	elementary; the Lean formalization is ours
Cor 3	known in substance	consequence of the same lemma; the vanishing $H \equiv 0$ at tight instances is used in [2, proof of Thm. 2.1]
Prop 5	known in substance	[16, Def. 3], [13] for $\varepsilon = 0$; $\varepsilon \neq 0$ routine
Prop 6	known	[15, Lemma 5.1(i)]; [17, Lemma 2.3]
Prop 7	ours	no statement located in the sources read
Obs 8, Table 1	ours	computation; no comparable table located
Thm 9, Cor 10	partially known	the tightness statements are [8, Thm. 3.1, 2.3], [17, Thm. 1.1, 1.2, 5.1]; the non-divisibility corollary was not located (original proof of [8] not read)
Prop 13	ours	elementary; (T) is not stated in [9], [17], [5], [11] as read. [5, Thm. 4.2] uses the non-divisibility of the four tight instances with $n \leq 6$ as an observation
Conj 12, 15	ours	see above; [9, Cor. 2.3 for E_m] is [9, Thm. 5.1]
Prop 16	ours	elementary; no prior statement located
Prop 18	ours (instance)	the shifted conjecture is false: [4, Thm. 1.8]; the instance $(1, 2, 3, 4, 6, 9)$ with half-phases is a second, independently checked counterexample, not claimed as new in kind
Rem 19	known	[4, Prop. 1.7], [10, Prop. 2.7, Lemma 2.5]

The qualitative statement that counterexamples and tight vectors lie on finitely many explicit hyperplanes is [2, Thm. 1.1, 2.1] (norm $2m + 3$), derivable with weaker bounds from [7]; the random-speed precursor is [6]. Nothing in this note claims it.

References

[1] Jaan Allikvere. Fourteen and fifteen lonely runners, 2026. arXiv:2609.02604, version 2.

[2] Matthias Beck and Samuel Everett. Lonely runner relations, 2026. arXiv:2609.06259, version 2.

[3] Benjamin Bedert. Riesz products and the lonely runner conjecture: a wider gap of loneliness, 2025. arXiv:2511.16636.

[4] Mónica Blanco, Francisco Criado, and Francisco Santos. Coloopless zonotopes and counterexamples to the shifted lonely runner conjecture, 2026. arXiv:2603.24784, version 3.

[5] Francesco Cordella. Odd denominators in the lonely runner spectrum for six speeds, 2026. arXiv:2609.03444, version 2.

[6] Sebastian Czerwiński. Random runners are very lonely. *J. Combin. Theory Ser. A*, 119:1194–1199, 2012. arXiv:1102.4464.

[7] Vishesh Giri and Noah Kravitz. The structure of lonely runner spectra. *Math. Proc. Cambridge Philos. Soc.*, 180:343–361, 2026. arXiv:2304.01462.

[8] Luis Goddyn and Erick B. Wong. Tight instances of the lonely runner. *Integers*, 6, 2006. Article A38.

[9] Noah Kravitz. Barely lonely runners and very lonely runners: a refined approach to the lonely runner problem. *Combinatorial Theory*, 1, 2021. Article 17; arXiv:1912.06034.

[10] Romanos Diogenes Malikiosis, Francisco Santos, and Matthias Schymura. Linearly-exponential checking is enough for the lonely runner conjecture and some of its variants. *Forum of Mathematics, Sigma*, 13:e164, 2025. arXiv:2411.06903.

[11] Guillem Perarnau and Oriol Serra. The lonely runner conjecture turns 60. *Computer Science Review*, 58:100798, 2025. arXiv:2409.20160.

[12] Daria Poliakova. More (shifted) runners, less loneliness, 2026. arXiv:2609.23952.

[13] Matthieu Rosenfeld. The lonely runner conjecture holds for eight runners, 2025. arXiv:2509.14111.

[14] Touch Sungkawichai and Tanupat Trakulthongchai. Eleven, twelve, and thirteen lonely runners, 2026. arXiv:2604.23906, version 2.

[15] Terence Tao. Some remarks on the lonely runner conjecture. *Contributions to Discrete Mathematics*, 13(2):1–31, 2018. arXiv:1701.02048.

[16] Tanupat Trakulthongchai. Nine and ten lonely runners. *Electron. J. Combin.*, 33(2), 2026. Article P2.46; arXiv:2511.22427.

[17] Yuhan Zhang. Tight instances of the lonely runner conjecture: complete classification of one-entry modifications, a new infinite family, and the growth bound, 2026. arXiv:2608.13599, version 2.