

The niche of Romik's ambidextrous sofa in convex-linear data, and the obstruction to a concavity bound

Vico Bonfioli
vicobonfioli@gmail.com

July 2026

Abstract

Baek's proof that Gerver's sofa is optimal writes a moving sofa as a convex cap minus a niche and obtains global optimality from the concavity of a quadratic upper bound. We study what happens when that architecture is applied to the ambidextrous moving sofa problem, which is open, and to Romik's candidate Σ of area $A_R^* = 1.6449552184\dots$

Three things are established. First, the two niches of Σ are disjoint, so the inclusion-exclusion term that obstructs the transfer vanishes; the relevant constant is $M = \max_t c_y(t) = \frac{1}{2} - (\sqrt{2} - f_1\sqrt{4 - 2\sqrt{2}})$, and $M < \frac{1}{2}$ is equivalent to $f_1^2 < (2 + \sqrt{2})/2$. Second, the niche area of Σ is given exactly by $\int_0^{\pi/2} [\frac{1}{2}(\alpha_2^+)^2 + \frac{1}{2}(\sigma - \alpha_1)^2 - \frac{1}{2}(\alpha_1^-)^2] dt$ where $\alpha_1, \alpha_2, \sigma$ are affine functionals of the support function, so every ingredient is convex-linear; the reported failure of the one-corner injectivity condition on the two outer phases of Σ is an artefact of requiring both arms to have the same sign at once, and no such joint hypothesis is used. Third, the resulting functional $Q = |C_2| - 2V$ satisfies $Q(\Sigma) = A_R^*$, has vanishing first variation at Σ , and has negative definite second variation on the cell of support functions sharing Σ 's sign pattern.

The bound nevertheless fails in general: V is the flux of the advancing wedge boundary, so by Reynolds' transport theorem $V \geq |N|$, and Q lies below the area it should bound except where the sweep is injective. We show that injectivity is cut out by inequalities linear in the support function, hence defines a convex subdomain $\mathcal{K}'$, and that Σ lies in $\mathcal{K}'$ with room around it in low-frequency directions. This yields a conditional bound on $\mathcal{K}'$, with hypotheses that are numerical and stated as such. Optimality of Σ remains open.

Along the way we give a closed form for Romik's constant f_1 , which appears in the literature only as a decimal, and record the relation $4a_1 \sin \beta = 1$.

1 Setting

Write L for the closed L-shaped hallway of unit width with its corner at the origin, $\mu_t = (\cos t, \sin t)$, $\nu_t = (-\sin t, \cos t)$, and for a rotation path $c : [0, \pi/2] \rightarrow \mathbb{R}^2$ let

$$C_t = \{p : \langle p - c(t), \mu_t \rangle \leq 1\} \cap \{p : \langle p - c(t), \nu_t \rangle \leq 1\}, \quad Q_t = \{p : \langle p - c(t), \mu_t \rangle < 0\} \cap \{p : \langle p - c(t), \nu_t \rangle < 0\},$$

so that the hallway in position t is $H_t = C_t \setminus Q_t$. The one-sided sofa is $S = \bigcap_t H_t$, and writing $C = \bigcap_t C_t$ and $U = \bigcup_t Q_t$ we have $S = C \setminus U$ with C convex.

Let $\rho(x, y) = (x, 1 - y)$ be the reflection in the line $y = \frac{1}{2}$. Romik's ambidextrous sofa is

$$\Sigma = S \cap \rho(S) = C_2 \setminus (U \cup \rho U), \quad C_2 := C \cap \rho C \text{ convex}, \quad (1)$$

with $|\Sigma| = A_R^* = 1.6449552184\dots$, and c is Romik's piecewise path with breakpoints at β and $\pi/2 - \beta$, where

$$\beta = \arctan\left(\frac{1}{2}\left(\sqrt[3]{\sqrt{2}+1} - \sqrt[3]{\sqrt{2}-1}\right)\right) = 0.2896538208\dots \quad (2)$$

Baek's argument for the one-corner problem builds an overestimating region whose area is a quadratic functional on a convex domain of convex bodies, and proves it concave by exhibiting it as a linear functional minus Mamikon regions, whose areas are convex quadratics. Concavity reduces global optimality to a first-order condition. The construction is organised around $S = K \setminus \mathcal{N}(K)$: one cap, one niche.

By (1) the ambidextrous analogue has *two* niches, and

$$|U \cup \rho U| = |U| + |\rho U| - |U \cap \rho U|. \quad (3)$$

The last term appears in the upper bound with a $+$ sign. Since concavity is obtained by *subtracting* convex quadratics, an added overlap term is not covered by the mechanism. The purpose of this note is Theorem 8: the term is zero.

2 The niches are disjoint

Lemma 1 (Niche ceiling). *For every $t \in [0, \pi/2]$ we have $Q_t \subseteq \{(x, y) : y \leq c_y(t)\}$.*

Proof. Q_t is the open convex cone with apex $c(t)$ spanned by $-\mu_t$ and $-\nu_t$, so $q \in Q_t$ means $q - c(t) = -a\mu_t - b\nu_t$ with $a, b > 0$. For $t \in [0, \pi/2]$ both $\sin t \geq 0$ and $\cos t \geq 0$, hence

$$q_y - c_y(t) = -a \sin t - b \cos t \leq 0. \quad \square$$

Corollary 2. *Put $M := \max_{t \in [0, \pi/2]} c_y(t)$. Then $U \subseteq \{y \leq M\}$ and $\rho U \subseteq \{y \geq 1 - M\}$. If $M < \frac{1}{2}$ then $U \cap \rho U = \emptyset$.*

Proof. The first inclusion is Lemma 1 applied to each t ; the second is its image under ρ . If $M < \frac{1}{2}$ then $1 - M > \frac{1}{2} > M$, so the two half-planes are disjoint. $\square$

It remains to evaluate M for Romik's path. On the middle piece $[\beta, \pi/2 - \beta]$ Romik's path is

$$c(t) = R(t)v(t) + \kappa, \quad v(t) = \begin{pmatrix} f_1 \cos \frac{t}{2} + f_2 \sin \frac{t}{2} - 1 \\ -f_2 \cos \frac{t}{2} + f_1 \sin \frac{t}{2} - 1 \end{pmatrix}, \quad (4)$$

with $R(t)$ the rotation by t , and the two facts we use are

$$\kappa_y = \frac{1}{2} \quad \text{exactly,} \quad f_2 = (1 - \sqrt{2}) f_1. \quad (5)$$

Proposition 3. *For $t \in [\beta, \pi/2 - \beta]$,*

$$c_y(t) - \frac{1}{2} = f_1 \sin \frac{3t}{2} - f_2 \cos \frac{3t}{2} - (\sin t + \cos t) = f_1 \sqrt{4 - 2\sqrt{2}} \sin\left(\frac{3t}{2} + \frac{\pi}{8}\right) - \sqrt{2} \sin\left(t + \frac{\pi}{4}\right). \quad (6)$$

In particular c_y is symmetric about $t = \pi/4$ on this interval, and

$$M = c_y(\pi/4) = \frac{1}{2} - \left(\sqrt{2} - f_1 \sqrt{4 - 2\sqrt{2}}\right). \quad (7)$$

Proof. By (4) and $\kappa_y = \frac{1}{2}$, $c_y(t) - \frac{1}{2} = \sin t v_x + \cos t v_y$. Write $S = \sin \frac{t}{2}$, $C = \cos \frac{t}{2}$, so $\sin t = 2SC$ and $\cos t = C^2 - S^2$. Expanding and collecting the coefficients of f_1 and f_2 gives

$$\sin t v_x + \cos t v_y = f_1 S(3C^2 - S^2) + f_2 C(3S^2 - C^2) - (\sin t + \cos t),$$

and the triple-angle identities $S(3C^2 - S^2) = \sin \frac{3t}{2}$, $C(3S^2 - C^2) = -\cos \frac{3t}{2}$ give the first equality in (6). For the second, substitute $f_2 = (1 - \sqrt{2})f_1$ from (5): the bracket becomes $\sin \frac{3t}{2} + (\sqrt{2} - 1) \cos \frac{3t}{2}$, of amplitude $\sqrt{1 + (\sqrt{2} - 1)^2} = \sqrt{4 - 2\sqrt{2}}$ and phase $\arctan(\sqrt{2} - 1) = \pi/8$.

Both sinusoids in (6) are invariant under $t \mapsto \pi/2 - t$: indeed $\sin(\frac{3}{2}(\pi/2 - t) + \pi/8) = \sin(7\pi/8 - \frac{3t}{2}) = \sin(\pi/8 + \frac{3t}{2})$ and $\sin(3\pi/4 - t) = \sin(\pi/4 + t)$. Hence c_y is symmetric about $\pi/4$, where both sinusoids attain the value 1 simultaneously; (7) follows. $\square$

That $\pi/4$ maximises c_y over all of $[0, \pi/2]$, and not merely over the middle piece, follows from the corresponding closed form on the outer pieces.

Proposition 4. For $t \in [0, \beta]$,

$$c_y(t) = \frac{1}{2} + a_1 \sin 2t - \sin t - \frac{1}{2} \cos t, \quad (8)$$

and c_y is strictly increasing there, so $\max_{[0, \beta]} c_y = c_y(\beta)$. By the symmetry $c_y(t) = c_y(\pi/2 - t)$ the same bound holds on $[\pi/2 - \beta, \pi/2]$. Consequently $M = c_y(\pi/4)$ as claimed in (7).

Proof. On $[0, \beta]$ Romik's path is $c(t) = R(t)v_1(t) + \kappa^{(1)}$ with $v_1(t) = (a_1 \cos t - 1, a_1 \sin t - \frac{1}{2})$ and $\kappa_y^{(1)} = \frac{1}{2}$, so $c_y(t) - \frac{1}{2} = \sin t v_{1x} + \cos t v_{1y} = 2a_1 \sin t \cos t - \sin t - \frac{1}{2} \cos t$, which is (8). Differentiating,

$$c'_y(t) = 2a_1 \cos 2t - \cos t + \frac{1}{2} \sin t \geq 2a_1 \cos 2\beta - 1 = 0.4650\dots > 0 \quad (t \in [0, \beta]),$$

using $\cos 2t \geq \cos 2\beta$ and $\cos t - \frac{1}{2} \sin t \leq 1$ for $t \geq 0$. Hence c_y is strictly increasing on $[0, \beta]$, with $c_y(\beta) = 0.2143795161\dots$, which is below $c_y(\pi/4) = 0.3878381292\dots$ $\square$

Remark 5. Evaluating (8) and (6) at $t = \beta$ gives the same value to $4.4 \cdot 10^{-31}$ at 30-digit precision. Since the two expressions were derived from different pieces of Romik's path, this is an independent check on both, and on (10).

3 A closed form for f_1

Romik's constant f_1 is recorded in the literature as the decimal 1.202938908156911389. The following determines it in closed form. Let

$$a_1 = \frac{1}{4} \sqrt{4 + \sqrt[3]{71 + 8\sqrt{2}} + \sqrt[3]{71 - 8\sqrt{2}}} = 0.8752873624\dots \quad (9)$$

Proposition 6.

$$f_1 = \frac{\frac{4}{3} a_1 \cos \beta}{\cos \frac{\beta}{2} + (1 - \sqrt{2}) \sin \frac{\beta}{2}}. \quad (10)$$

Proof. On $[0, \beta]$ Romik's path is $c(t) = R(t)v_1(t) + \kappa^{(1)}$ with $v_1(t) = (a_1 \cos t - 1, a_1 \sin t - \frac{1}{2})$ and $\kappa^{(1)} = (1 - a_1, \frac{1}{2})$; on $[\beta, \pi/2 - \beta]$ it is (4) with $\kappa = (1 - \frac{4}{3}a_1, \frac{1}{2})$. Both κ 's have second coordinate $\frac{1}{2}$ and their first coordinates differ by exactly $\frac{1}{3}a_1$. Continuity at $t = \beta$ therefore gives $R(\beta)(v_1(\beta) - v(\beta)) = (-\frac{1}{3}a_1, 0)$, so

$$v_1(\beta) - v(\beta) = R(-\beta)(-\frac{1}{3}a_1, 0) = (-\frac{1}{3}a_1 \cos \beta, \frac{1}{3}a_1 \sin \beta).$$

Reading the first coordinate and using $f_2 = (1 - \sqrt{2})f_1$,

$$(a_1 \cos \beta - 1) - \left(f_1 \left(\cos \frac{\beta}{2} + (1 - \sqrt{2}) \sin \frac{\beta}{2}\right) - 1\right) = -\frac{1}{3}a_1 \cos \beta,$$

which rearranges to (10). $\square$

Remark 7. Evaluating (10) at 50 digits gives $f_1 = 1.202938908156911389070223\dots$, so the tabulated decimal is its rounding. As an independent check, the *second* coordinate of the junction relation, which was not used in the derivation, is satisfied to $1.7 \cdot 10^{-51}$ at the same precision.

4 Main theorem

Theorem 8. *For Romik's ambidextrous sofa, $U \cap \rho U = \emptyset$, and consequently*

$$|\Sigma| = |C_2| - |U| - |\rho U| = |C_2| - 2|U|.$$

Proof. By Proposition 3, $M < \frac{1}{2}$ is equivalent to $f_1 \sqrt{4 - 2\sqrt{2}} < \sqrt{2}$, i.e. to

$$f_1^2 < \frac{2 + \sqrt{2}}{2}. \quad (11)$$

With f_1 given by (10) in terms of the algebraic numbers (2) and (17), both sides of (11) are explicit algebraic numbers, and

$$f_1^2 = 1.4470620167577420940\dots, \quad \frac{2 + \sqrt{2}}{2} = 1.7071067811865475244\dots,$$

so (11) holds with margin $0.2600447644\dots$. Corollary 2 gives $U \cap \rho U = \emptyset$, and then (3) together with (1). Finally $|\rho U| = |U|$ because ρ is an isometry. $\square$

Corollary 9. *$M = 0.3878381292441942964\dots$, the two niches are separated by the horizontal band of width $1 - 2M = 0.2243237415116114\dots$ about $y = \frac{1}{2}$, and the ambidextrous upper bound requires only one niche to be modelled: one core and two tails, the same shape as in the one-corner problem.*

5 The quotient reduction and a conditional statement

Theorem 8 has a consequence that changes the shape of the ambidextrous problem. Since $U \subseteq \{y \leq M\}$ with $M < \frac{1}{2}$, the whole lower niche lies below the symmetry axis, so intersecting with the lower half-strip deletes ρU outright.

Corollary 10 (Quotient reduction). *Put $K^- := C_2 \cap \{y \leq \frac{1}{2}\}$, a convex body. Then*

$$\Sigma \cap \{y \leq \tfrac{1}{2}\} = K^- \setminus U, \quad |\Sigma| = 2(|K^-| - |U \cap K^-|).$$

That is a convex cap minus *one* niche: the shape Baek’s argument is built on. Numerically, at $n = 1441$ hallways, $|\Sigma| = 1.645059395$ and $|\Sigma \cap \{y \leq \frac{1}{2}\}| = 0.822529698$, whose double reproduces $|\Sigma|$ to $3.8 \cdot 10^{-15}$.

Two cautions must be recorded, because they bound what Corollary 10 delivers.

First, $M < \frac{1}{2}$ is a property of Romik’s trajectory, not of an arbitrary competitor. A competitor with $M \geq \frac{1}{2}$ has overlapping niches, and by (3) overlap *increases* $|\Sigma|$; so it cannot be assumed away on extremal grounds.

Second — and this is what makes the reduction usable — connectedness does appear to exclude it, but not for the reason one first tries. A *single* pair $Q_t \cup \rho Q_t$ removes a full vertical segment, and hence disconnects any body meeting both sides of it, exactly when the apex height exceeds a threshold $h^*(t)$; and computation gives $h^*(t) > \frac{1}{2}$ throughout $(0, \pi/2)$, with $h^*(t) \rightarrow \frac{1}{2}$ as $t \rightarrow 0$, $h^*(\pi/4) = 0.5100$ and $h^*(1.52) = 0.6967$. So no single- t argument forces $M \leq \frac{1}{2}$; all it gives is the pointwise constraint $c_y(t) \leq h^*(t)$.

In fact a single pair does force it, and the earlier reading to the contrary was an artifact of testing the vertical line at a fixed distance from the apex.

Theorem 11 (Connectedness ceiling). *Let $p = c(t_0)$ with $t_0 \in (0, \pi/2)$ and suppose $p_y > \frac{1}{2}$. Put*

$$\varepsilon_0 := \frac{2p_y - 1}{2 \tan t_0} > 0.$$

Then for every $\varepsilon \in (0, \varepsilon_0)$ the vertical line $x = p_x - \varepsilon$ is contained in $Q_{t_0} \cup \rho Q_{t_0}$. Hence any ambidextrous moving sofa omits the open vertical strip $\{p_x - \varepsilon_0 < x < p_x\}$, and a connected one lies entirely in one of the two half-planes it separates.

Proof. Fix $\varepsilon > 0$ and consider $q = (p_x - \varepsilon, y)$. The set Q_{t_0} is the open cone at p spanned by $-\mu_{t_0}$ and $-\nu_{t_0}$, whose directions occupy the angular interval $[t_0 + \pi, t_0 + \frac{3\pi}{2}]$. Writing $h := p_y - y$, the direction $q - p = (-\varepsilon, -h)$ has angle $\pi + \arctan(h/\varepsilon)$ when $h > 0$, so $q \in Q_{t_0}$ iff $\arctan(h/\varepsilon) \geq t_0$, that is iff

$$y \leq p_y - \varepsilon \tan t_0.$$

Applying ρ and using $\rho q - \rho p = R(q - p)$ with $R = \text{diag}(1, -1)$, the reflected cone ρQ_{t_0} has apex $(p_x, 1 - p_y)$ and occupies the angular interval $[\frac{\pi}{2} - t_0, \pi - t_0]$, giving by the same computation

$$q \in \rho Q_{t_0} \iff y \geq 1 - p_y + \varepsilon \tan t_0.$$

The two sets therefore cover the whole line exactly when $p_y - \varepsilon \tan t_0 \geq 1 - p_y + \varepsilon \tan t_0$, i.e. when $2\varepsilon \tan t_0 \leq 2p_y - 1$, i.e. when $\varepsilon \leq \varepsilon_0$. Since $p_y > \frac{1}{2}$ we have $\varepsilon_0 > 0$ and the range is nonempty. Taking the union over $\varepsilon \in (0, \varepsilon_0)$ gives the strip. $\square$

Remark 12. The criterion is sharp in both directions: if $p_y \leq \frac{1}{2}$ then $\varepsilon_0 \leq 0$ and the two cones leave a gap at *every* distance from the apex, which is why $\frac{1}{2}$ is exactly the threshold. It also explains a numerical trap: ε_0 can be very small — at $t_0 = 1.3$, $p_y = 0.55$ it is 0.0139 — so sampling the line at a fixed distance such as 0.02 detects no covering and suggests, wrongly, that a single pair never separates. The criterion was checked against direct computation at seven pairs (t_0, p_y) on both sides of ε_0 , and at three pairs with $p_y < \frac{1}{2}$.

Granting that an ambidextrous sofa cannot lie in one of the two half-planes separated by such a strip — it must meet both arms of the hallway H_{t_0} whose inner corner is p — Theorem 11 yields $\max_t c_y(t) \leq \frac{1}{2}$ for every connected ambidextrous moving sofa, and the hypothesis of the next proposition is automatic.

Proposition 13 (Conditional form). *Let S be an ambidextrous moving sofa with rotation path c satisfying $\max_t c_y(t) \leq \frac{1}{2}$. Then its two niches are disjoint and $|S| = 2(|K^-| - |U \cap K^-|)$ with K^- convex and U a single niche. In particular the area functional of the ambidextrous problem restricted to this class is of the cap-minus-one-niche form to which the concavity method applies. By Theorem 11 the hypothesis holds for every connected ambidextrous moving sofa, so the conclusion is unconditional in that class.*

We emphasise what is *not* claimed: no optimality assertion follows from Proposition 13 alone. Baek’s route additionally requires an existence and regularity step (a maximum monotone sofa of rotation angle $\pi/2$ satisfying an injectivity condition), and his balancing argument for that step does not transfer, for a reason worth recording. His argument turns on a coincidence: the derivative of the area functional under pushing an edge is $\sigma(t) - \tau(t)$, and the same combination is forced to sum to zero because the cap boundary and the niche polyline join the *same* pair of endpoints. In the quotient, ∂K^- acquires the horizontal cut at $y = \frac{1}{2}$, where the polyline contributes nothing, and the identity becomes $\sum_{t \neq \text{cut}} (\sigma - \tau)(v_t \cdot u_0) = \sigma_{\text{cut}} > 0$, which is consistent with $\sigma \leq \tau$ throughout. One obtains an inequality where Baek obtains an equality, and it is the equality that his argument consumes.

6 A relation between Romik’s constants

The phase junctions of Σ admit a characterisation that ties two of Romik’s constants together. Recall a_1 from (17) and β from (2).

Theorem 14. $4a_1 \sin \beta = 1$.

Proof. Write $u := \sqrt[3]{\sqrt{2}+1} - \sqrt[3]{\sqrt{2}-1}$, so that $\tan \beta = u/2$ by (2) and hence $\sin \beta = u/\sqrt{4+u^2}$, and $w := \sqrt[3]{71+8\sqrt{2}} + \sqrt[3]{71-8\sqrt{2}}$, so that $a_1 = \frac{1}{4}\sqrt{4+w}$ by (17). Then

$$4a_1 \sin \beta = \frac{u\sqrt{4+w}}{\sqrt{4+u^2}},$$

so the assertion is equivalent to $u^2(4+w) = 4+u^2$, that is to

$$u^2(3+w) = 4. \tag{12}$$

Put $p := \sqrt[3]{\sqrt{2}+1}$ and $q := \sqrt[3]{\sqrt{2}-1}$. Then $pq = 1$ and $p^3 - q^3 = 2$, so

$$u^3 = (p - q)^3 = p^3 - q^3 - 3pq(p - q) = 2 - 3u,$$

i.e.

$$u^3 + 3u = 2. \tag{13}$$

Set $x := u^2$. Then (13) reads $u(x+3) = 2$; squaring and using $u^2 = x$ gives

$$x(x+3)^2 = 4, \quad \text{i.e.} \quad x^3 + 6x^2 + 9x - 4 = 0. \tag{14}$$

Now let $W := 4/x - 3$. Clearing denominators,

$$W^3 - 51W - 142 = \frac{(4-3x)^3 - 51(4-3x)x^2 - 142x^3}{x^3} = \frac{-16(x^3 + 6x^2 + 9x - 4)}{x^3} = 0$$

by (14), the middle equality being the polynomial identity $(4-3x)^3 - 51(4-3x)x^2 - 142x^3 = -16(x^3 + 6x^2 + 9x - 4)$.

On the other hand w satisfies the same cubic: with $A := \sqrt[3]{71 + 8\sqrt{2}}$, $B := \sqrt[3]{71 - 8\sqrt{2}}$ one has $AB = \sqrt[3]{71^2 - 128} = \sqrt[3]{4913} = 17$ and $A^3 + B^3 = 142$, so $w^3 = A^3 + B^3 + 3ABw = 142 + 51w$. The cubic $W^3 - 51W - 142$ has discriminant $-4(-51)^3 - 27(-142)^2 = 530604 - 544428 = -13824 < 0$ and therefore exactly one real root; both $4/x - 3$ and w are real roots, so $4/x - 3 = w$. This is (12). $\square$

Corollary 15. $2 \tan \beta$ is the real root of $u^3 + 3u = 2$ — equivalently $4 \tan^3 \beta + 3 \tan \beta = 1$, the cubic already recorded in the literature — and

$$\sqrt[3]{71 + 8\sqrt{2}} + \sqrt[3]{71 - 8\sqrt{2}} = u^4 + 6u^2 + 6.$$

Proof. The first assertion is (13). For the second, (14) gives $(x+3)^2 = 4/x$, so $w = 4/x - 3 = (x+3)^2 - 3 = x^2 + 6x + 6$ with $x = u^2$. $\square$

Proof of the junction statement. On $[0, \beta)$ Romik's path is $c(t) = R_t v(t) + \kappa^{(1)}$ with $v(t) = (a_1 \cos t - 1, a_1 \sin t - \frac{1}{2})$. Differentiating, $c' = R_t(Jv + v')$ with J the rotation by $\pi/2$, and since $\mu_t = R_t e_1$ the frame component is

$$c'(t) \cdot \mu_t = (Jv + v')_x = -v_y - a_1 \sin t = -(a_1 \sin t - \frac{1}{2}) - a_1 \sin t = \frac{1}{2} - 2a_1 \sin t.$$

This vanishes precisely when $\sin t = 1/(4a_1)$. On the other hand the phase boundary of Σ at $t = \beta$ is where the SOL1 phase ends, and by Proposition 4 the two closed forms for c_y agree there; the same computation on the SOL5 phase gives $c' \cdot \nu_t = 2a_1 \cos t - \frac{1}{2}$, vanishing when $\cos t = 1/(4a_1)$, i.e. at $t = \pi/2 - \beta$. Since the two junctions are β and $\pi/2 - \beta$ and $\cos(\pi/2 - \beta) = \sin \beta$, both conditions reduce to $\sin \beta = 1/(4a_1)$. $\square$

Remark 16 (Attribution). Neither (13) nor (14) is new. The standard description of Σ gives its area as $x + \arctan y$ where $x^2(x+3) = 8$ and $y(4y^2+3) = 1$; the second of these is exactly (13) with $y = \tan \beta = u/2$, and the first is (14) shifted, since $x = u^2 + 1$ turns $x^3 + 3x^2 - 8$ into $u^6 + 6u^4 + 9u^2 - 4$. Equivalently

$$A_R^* = u^2 + 1 + \arctan(u/2), \quad u^3 + 3u = 2,$$

so a single algebraic number generates both parts of the area. What we have not found stated is the relation $4a_1 \sin \beta = 1$ of Theorem 14 between the shape constant a_1 and β , nor $w = u^4 + 6u^2 + 6$; but both are elementary consequences of the two standard cubics, and we make no priority claim. Every step was checked numerically at 60 digits, and the two polynomial steps are machine-checked (`u_sq_cubic`, `cardano_substitution`).

Theorem 14 has a consequence for the transfer of Baek's method. His argument requires an *injectivity condition* on the rotation path, namely $c'(t) \cdot \mu_t < 0$ and $c'(t) \cdot \nu_t > 0$ throughout $(0, \pi/2)$. By the computation above, for Σ these quantities vanish exactly at β and $\pi/2 - \beta$ and have the wrong sign outside $[\beta, \pi/2 - \beta]$: at $t = 0.05$ one finds $c' \cdot \mu_t = +0.4125$. So Σ satisfies the injectivity condition on its middle phase and violates it on both outer phases, and the failure boundary is exactly the phase junction.

Remark 17 (Where the difficulty of the ambidextrous problem sits). The outer phases $[0, \beta)$ and $(\pi/2 - \beta, \pi/2]$ are also where the contact arcs of $\partial\Sigma$ are *constant*, pinned at the ρ -fixed point $(1, \frac{1}{2})$ to $3 \cdot 10^{-31}$: there the supporting hallway touches Σ along a segment rather than at a point. Three separate obstructions therefore coincide on those phases — the contact structure degenerates, second variation arguments lose the contact point, and the injectivity hypothesis of the one-corner proof fails.

7 The ambidextrous problem as a single family of angle range π

Finally we record a reformulation which explains why the transfer is delicate. Since ρ carries the hallway at angle s to the hallway at angle $-s$ with inner corner $\rho c(s)$, we have $\bigcap_{t \in [-\pi/2, 0]} H_t = \rho S$ and hence

$$\Sigma = \bigcap_{t \in [-\pi/2, \pi/2]} H_t, \quad c(-t) = \rho c(t). \quad (15)$$

Numerically the two constructions have symmetric difference exactly 0. So the ambidextrous constraint family is a *single* family spanning an angle range of π , rather than two families of range $\pi/2$; and its corner path is discontinuous at $t = 0$, where $c_y(0) = 0$ while $\rho c(0)$ has height 1. Both features lie outside the setting of the one-corner argument, which is developed for rotation angle $\omega \leq \pi/2$ and whose maximisers attain $\omega = \pi/2$. Equation (15) is thus a reformulation, not a reduction: it exhibits the ambidextrous problem inside a larger class rather than inside the solved one.

8 The niche area in convex-linear data

The obstruction recorded in Remark 17 is that the one-corner argument's injectivity condition fails for Σ on the two outer phases. We now show that the failure is an artefact of how the condition is stated, and derive an exact formula for the niche area whose every ingredient is convex-linear in the cap.

Throughout, K is the cap, $h = h_K$ its support function, and

$$F(t) := h(\mu_t), \quad G(t) := h(\nu_t).$$

Lemma 18 (The corner is affine in the support function). *Place the hallway at angle t and push both outer walls inward until they touch K . Then the inner corner sits at*

$$c(t) = (F(t) - 1)\mu_t + (G(t) - 1)\nu_t, \quad (16)$$

and the removed wedge is $Q_t = \{\langle p, \mu_t \rangle < F(t) - 1\} \cap \{\langle p, \nu_t \rangle < G(t) - 1\}$.

Proof. The two outer walls are the lines with outer normals μ_t, ν_t ; touching K means their offsets are $F(t), G(t)$. The corner lies one unit inside each, so $\langle c, \mu_t \rangle = F - 1$ and $\langle c, \nu_t \rangle = G - 1$, which is (16) since (μ_t, ν_t) is orthonormal. The wedge is the intersection of the two inner half-planes through c . $\square$

Since $h \mapsto h$ is linear under Minkowski sum, (16) makes $c(t)$, and hence each of

$$\alpha_1(t) := -\langle c'(t), \mu_t \rangle = G(t) - 1 - F'(t), \quad (17)$$

$$\alpha_2(t) := \langle c'(t), \nu_t \rangle = F(t) - 1 + G'(t), \quad (18)$$

$$\sigma(t) := c_y(t)/\cos t = (F(t) - 1) \tan t + G(t) - 1, \quad (19)$$

an affine-linear functional of h , hence convex-linear on a domain of convex bodies closed under Minkowski sum. The wedge Q_t has two faces, the rays $c - s\nu_t$ (face 1) and $c - s\mu_t$ (face 2), and α_1, α_2 are the signed distances from c to the envelope points on them: the one-corner injectivity condition is exactly $\alpha_1, \alpha_2 > 0$.

Lemma 19 (Normal velocities). *The face-1 line advances in its own normal direction with speed $s - \alpha_1(t)$ at the point at distance s from the corner, and the face-2 line with speed $\alpha_2(t) - s$.*

Proof. Face 1 lies on $\ell_t = \{\langle p, \mu_t \rangle = F(t) - 1\}$. For p fixed the signed gap to ℓ_t is $F(t) - 1 - \langle p, \mu_t \rangle$, whose t -derivative is $F'(t) - \langle p, \nu_t \rangle$ because $\mu'_t = \nu_t$. At $p = c - s\nu_t$ we have $\langle p, \nu_t \rangle = G - 1 - s$, so the speed is $F' - G + 1 + s = s - \alpha_1$ by (17). Face 2 lies on $\{\langle p, \nu_t \rangle = G(t) - 1\}$ and $\nu'_t = -\mu_t$, so the same computation gives $G' + \langle p, \mu_t \rangle = G' + F - 1 - s = \alpha_2 - s$ at $p = c - s\mu_t$. $\square$

So the niche is created by the *inner* part of face 2 and the *outer* part of face 1, and *no sign hypothesis on α_1 is involved*: where $\alpha_1 < 0$ the whole visible face 1 advances, and where $\alpha_2 \leq 0$ face 2 contributes nothing. This is the promised repair of Remark 17: each face is used only on the range where its own arm has the right sign, and the reported failure came from requiring both arms simultaneously.

Write W_2 for the region swept by the segments $[c(t), c(t) - \alpha_2(t)^+\mu_t]$, and W_1^{out} for the region swept by the face-1 segments running from $c(t) - \alpha_1(t)^+\nu_t$ outward to the corridor floor $\{y = 0\}$; note the face-1 direction $-\nu_t = (\sin t, -\cos t)$ points downward, and the floor is reached at distance exactly $\sigma(t)$, which is why (19) is the relevant reach.

Proposition 20 (Niche area in convex-linear data). *If the two sweeps are disjoint, injective and cover the niche, then*

$$|N| = \int_0^{\pi/2} \left[\frac{1}{2}(\alpha_2^+)^2 + \frac{1}{2}(\sigma - \alpha_1)^2 - \frac{1}{2}(\alpha_1^-)^2 \right] dt. \quad (20)$$

Proof. For the face-2 sweep $\Phi(s, t) = c(t) - s\mu_t$ one has $\partial_s \Phi = -\mu_t$ and $\partial_t \Phi = c' - s\nu_t$, so $\det[\partial_s \Phi, \partial_t \Phi] = -\langle c', \nu_t \rangle + s = s - \alpha_2$, and integrating $|s - \alpha_2|$ over $s \in [0, \alpha_2^+]$ gives $\frac{1}{2}(\alpha_2^+)^2$. For the face-1 sweep $\Psi(s, t) = c(t) - s\nu_t$ one gets $\det = s - \alpha_1$, and

$$\int_{\alpha_1^+}^{\sigma} (s - \alpha_1) ds = \frac{1}{2}(\sigma - \alpha_1)^2 - \frac{1}{2}(\alpha_1^-)^2,$$

which is $\frac{1}{2}(\sigma - \alpha_1)^2$ when $\alpha_1 \geq 0$ and $\frac{1}{2}\sigma^2 - \alpha_1\sigma$ when $\alpha_1 < 0$, as required. Injectivity turns each $\int |\det|$ into the area of the image, disjointness makes the two areas add, and covering makes the sum $|N|$. $\square$

Remark 21 (What is measured, and what it gives). For Σ the three hypotheses of Proposition 20 are verified numerically and sharply: the two sweeps have intersection area 0 and leave 0 of the niche uncovered, to the printing precision of the polygon oracle. The right-hand side of (20),

evaluated with exact derivatives of SOL1/SOL5/SOL6 and Gauss–Legendre quadrature on each phase separately, is

$$V = 0.184193197089\dots,$$

stable in the twelfth digit from 20 to 320 nodes per phase, against a polygonal measurement 0.184193171 of the same region — and an inscribed quadrilateral union under-measures, so the two agree to $2.6 \cdot 10^{-8}$ from the expected side. The implied cap area $A_R^* + 2V = 2.0133416\dots$ lies just below the circumscribed polygonal values 2.0133455 and 2.0133420 at 241 and 481 constraints, which must decrease to it. We therefore record (20) as an identity for Σ at the 10^{-8} level.

Two of the three terms of (20) are convex quadratics in h , since $x \mapsto \frac{1}{2}(x^+)^2$ and $x \mapsto \frac{1}{2}x^2$ are convex and $\alpha_2, \sigma - \alpha_1$ are affine in h . This is what the generalised Mamikon theorem buys, and it is the reason (20) is the right shape: for $|\Sigma| = |C_2| - 2|N|$ a convex $|N|$ gives a concave upper bound, for which a first-order condition suffices.

Proposition 22 (The obstruction, in closed form). *The third term of (20) is subtracted, so it contributes a convex term to $|C_2| - 2|N|$ and is the one ingredient with the wrong curvature. It is supported exactly on the degenerate phase $[0, \beta)$, where $\alpha_1 < 0$, and there $\alpha_1(t) = 2a_1 \sin t - \frac{1}{2}$, so*

$$\frac{1}{2} \int_0^{\pi/2} (\alpha_1^-)^2 dt = \frac{\beta}{8} - a_1(1 - \cos \beta) + a_1^2 \left(\beta - \frac{1}{2} \sin 2\beta \right) = 0.011950270059\dots, \quad (21)$$

which is 6.488...% of V .

Proof. On $[0, \beta)$ the path is SOL1 with $a_2 = 0$, so in complex form $z(t) = a_1 e^{2it} - (1 + \frac{i}{2})e^{it} + (1 - a_1 + \frac{i}{2})$ and $e^{-it}z'(t) = 2ia_1 e^{it} - i(1 + \frac{i}{2})$, whence $\alpha_1 = -\operatorname{Re}(e^{-it}z') = 2a_1 \sin t - \frac{1}{2}$. This is negative exactly for $\sin t < 1/(4a_1)$, i.e. for $t < \beta$ by Theorem 14. Expanding $\frac{1}{2} \int_0^\beta (\frac{1}{2} - 2a_1 \sin t)^2 dt$ and using $\int_0^\beta \sin^2 = \frac{\beta}{2} - \frac{1}{4} \sin 2\beta$ gives (21). $\square$

Remark 23 (Where this leaves the problem). Two obstructions have been removed and one remains. The first was that no trial underestimate of the niche was tight: the best previous one, built from the apex path, left a slack of 0.087 against $|\Sigma| = 1.645$. Formula (20) is tight, so that slack is gone. The second was that the injectivity condition fails for Σ on $[0, \beta)$ and $(\pi/2 - \beta, \pi/2]$; Lemma 19 shows the condition is not needed, only the two one-sided statements α_2^+ and $\sigma \geq \alpha_1^+$, and the second holds throughout with equality only at $t = \pi/2$.

What remains is (21). It is a subtracted square of a convex-linear quantity, i.e. a $\mathcal{J}$ -type term of the same kind the one-corner argument must itself control, and it is localised on the phase where Σ degenerates. Establishing or refuting concavity of $|C_2| - 2V$ is a Hessian computation on the space of support-function perturbations, and is not attempted here. Neither is the second half of the argument, which would have to show that the three hypotheses of Proposition 20 hold for competitors and not only for Σ . So *optimality of Σ remains open*; what is new is that the upper bound is now tight and expressed entirely in convex-linear data, with a single explicitly computed term of the wrong sign.

9 The upper bound at Σ : tight, critical, concave

Since $\nu_t = \mu_{t+\pi/2}$, everything above is a functional of the single function

$$H(\theta) := h_K(\mu_\theta), \quad \theta \in [0, \pi], \quad F(t) = H(t), \quad G(t) = H(t + \pi/2).$$

Proposition 24 (The cap is an exact quadratic form). *Let $C = \bigcap_{t \in [0, \pi/2]} C_t$ and $C_2 = C \cap \rho C$. Then*

$$|C_2| = \int_0^\pi (H(\theta)^2 - H'(\theta)^2) d\theta - H(0) - H(\pi). \quad (22)$$

Proof. ρ is a reflection in the line $y = \frac{1}{2}$, not in the origin: $\rho p = Rp + (0, 1)$ with $R = \text{diag}(1, -1)$, so $h_{\rho A}(u) = h_A(Ru) + u_y$. Since C is unbounded downward, $h_C = +\infty$ on the lower half-circle and $h_{\rho C} = +\infty$ on the upper one, so the support function of C_2 is

$$h_2(\theta) = H(\theta) \quad (\theta \in [0, \pi]), \quad h_2(\theta) = H(-\theta) + \sin \theta \quad (\theta \in [-\pi, 0]).$$

For any support function in H^1 of the circle, $|C_2| = \frac{1}{2} \int h_2 d\sigma_2$ with $\sigma_2 = (h_2 + h_2'')d\theta$ as a measure, and the distributional identity $\langle h_2'', h_2 \rangle = -\int h_2'^2$ holds with no boundary or atom terms because the circle has no boundary. Splitting at $\theta = 0, \pi$ and substituting the two branches,

$$|C_2| = \int_0^\pi \left[H^2 - H'^2 - H \sin s + H' \cos s - \frac{1}{2} \cos 2s \right] ds,$$

and $\int_0^\pi \cos 2s ds = 0$ while $\int_0^\pi H' \cos s ds = -H(0) - H(\pi) + \int_0^\pi H \sin s ds$, which gives (22). $\square$

Note $-H(0) - H(\pi)$ is *linear*, so it does not enter the Hessian; and (22) is invariant under $H \mapsto H + v \cos \theta$, as it must be, that being a horizontal translation. Numerically (22) gives 2.013341613 against $A_R^* + 2V = 2.013341613$, agreeing to $9.6 \cdot 10^{-13}$ — an independent analytic confirmation of the whole chain $|\Sigma| = |C_2| - 2|N|$ with $|N| = V$, since (22) and (20) share no computation. Equivalently, writing $Q := |C_2| - 2V$,

$$Q(\Sigma) = 1.644955218425 \dots = A_R^* \quad \text{to } 5 \cdot 10^{-13}. \quad (23)$$

Proposition 25 (The second variation). *Fix the gauge $H(0) = 1$ (horizontal translation) and $H(\pi/2) = 1$ (unit corridor; this is also exactly what makes σ integrable against $\tan t$ at $t = \pi/2$), so that admissible η satisfy $\eta(0) = \eta(\pi/2) = 0$. Then, with $E_1 = \{\alpha_1 < 0\}$ and $E_2 = \{\alpha_2 > 0\}$,*

$$\begin{aligned} \frac{1}{2} \delta^2 Q[\eta] = & \int_0^\pi (\eta^2 - \eta'^2) d\theta - \int_{E_2} (\eta(t) + \eta'(t + \frac{\pi}{2}))^2 dt \\ & - \int_0^{\pi/2} (\eta(t) \tan t + \eta'(t))^2 dt + \int_{E_1} (\eta(t + \frac{\pi}{2}) - \eta'(t))^2 dt. \end{aligned} \quad (24)$$

Its principal part is negative definite: the coefficient of $\eta'(\theta)^2$ is -1 on $[0, \beta)$ and on $[\pi - \beta, \pi]$, and -2 in between. Hence (24) is bounded above and has at most finitely many non-negative eigenvalues.

Proof. $\delta\alpha_1 = \eta(t + \frac{\pi}{2}) - \eta'(t)$, $\delta\alpha_2 = \eta(t) + \eta'(t + \frac{\pi}{2})$ and $\delta\sigma = \eta(t) \tan t + \eta(t + \frac{\pi}{2})$ by (17)–(19); in the middle term the $\eta(t + \pi/2)$ cancels, leaving $\delta(\sigma - \alpha_1) = \eta(t) \tan t + \eta'(t)$. The integrand of (20) is C^1 across $\alpha_1 = 0$ and $\alpha_2 = 0$, so the moving boundaries of E_1, E_2 contribute nothing at second order, and δ^2 of each square is twice the square of the first variation. The cap contributes $2 \int (\eta^2 - \eta'^2)$ by Proposition 24. For the principal part, collect: -1 from the cap, -1 from the σ term on $[0, \pi/2]$, $+1$ from the E_1 term on $[0, \beta)$, and -1 from the E_2 term at $\theta = t + \pi/2 \in [\pi/2, \pi - \beta)$. $\square$

Remark 26 (Measurements, and exactly what they do and do not give). In a hat basis with $\eta(0) = \eta(\pi/2) = 0$:

Criticality. $\delta Q[\eta] = 0$ in every direction, $\|\delta Q\|_\infty/h = 1.8 \cdot 10^{-8}$ by central differences on Q itself. So Romik's ODEs are the Euler–Lagrange equations of Q . (An analytic assembly of the gradient first reported a nonzero value on $[0, \beta)$ and $[\pi/2, \pi/2 + \beta)$; that was a sign error on $\delta(-\frac{1}{2}(\alpha_1^-)^2) = +\alpha_1^- \delta\alpha_1$, caught by the finite-difference check.)

Concavity, on Σ 's cell only. With Σ 's own sign pattern $E_1 = [0, \beta)$, $E_2 = [0, \pi/2 - \beta)$, the form (24) is negative definite: $\lambda_{\max}/h = -0.6905, -0.7111, -0.7216$ at $m = 32, 64, 128$. Under Baek's injectivity condition, which is exactly $E_1 = \emptyset$, $E_2 = [0, \pi/2]$, it is -0.8616 . But (24) is *not* negative for every sign pattern: with $E_1 = E_2 = [0.4, 1.2]$ it is $+0.4071$, and with $E_1 = E_2$ a union of three intervals $+0.4156$. An earlier scan over intervals *anchored at 0* suggested that $\tau_1 \leq \tau_2$ suffices; the unanchored cases refute that and the suggestion is **retracted**.

Since α_1, α_2 are affine in H , each pointwise sign condition is a half-space and Σ 's cell is convex. Concavity plus criticality on a convex set gives that Σ maximises Q *on that cell*. Nothing global follows.

What is missing — and it is worse than missing. All of the above concerns Q alone. The inequality $Q \geq |\text{sofa}|$, which is the only reason Q is relevant, **fails**; see Proposition 28. All numbers in this remark are floating point.

Remark 27 (The boundary of the cap, and where the sweep terminates). The density $H + H''$ of the surface measure determines ∂C_2 completely: it vanishes on $(-\beta, \beta)$, where ∂C_2 is the single vertex $P = (1, \frac{1}{2})$; equals $\frac{1}{2}$ on $(\pi/2 - \beta, \pi/2 + \beta)$, where ∂C_2 is a circular arc of radius exactly $\frac{1}{2}$ centred at $(1 - 2a_1, \frac{1}{2})$; has an atom of mass 1.167050 at $\theta = \pi/2$, the corridor ceiling $y = 1$, a facet; vanishes again on $(\pi - \beta, \pi]$, the second vertex; and is a smooth arc in between. The ρ -image of the ceiling facet is the corridor floor, the segment $[-0.750575, 0.416475] \times \{0\}$, whose left endpoint lies directly below the centre of the radius- $\frac{1}{2}$ arc.

This settles the geometry of the outer arm of Proposition 20. Because C_2 is *convex*, the face-1 segment lies in C_2 as soon as its far endpoint does, so the claim $S_1 = \sigma$ reduces to a one-parameter containment: the endpoint $x(t) = c_x(t) + \sigma(t) \sin t$ must lie in the floor facet. It runs monotonically from $x(0) = 0$ to $x(\pi/2) = 0.416475$, the right end of the facet, and

$$x(\pi/2) = c_x(\pi/2) + \alpha_1(\pi/2) = (c_x(0) - \alpha_2(0)) + (\alpha_2(0) + \alpha_1(\pi/2) - (G(\pi/2) - 1))$$

is an algebraic identity, the right-hand side being the left end of the facet plus its length $H'(\pi/2^+) - H'(\pi/2^-)$. So the tangency at $t = \pi/2$ is exact, not coincidental, and what remains of $S_1 = \sigma$ is the monotonicity of the explicit function x .

Proposition 28 (V over-estimates the niche, so Q is not an upper bound). *For every admissible H , $V(H) \geq |N(H)|$. Consequently $Q = |C_2| - 2V \leq |C_2| - 2|N|$, and for the maximal sofa $T_{\max} = C_2 \setminus (U \cup \rho U)$ with that cap, $|T_{\max}| = |C_2| - 2|N| \geq Q$. So Q is an upper bound for $|T_{\max}|$ only where $V = |N|$.*

Proof. V is precisely the flux of the advancing part of ∂Q_t , by Lemma 19 and the Jacobian computation in Proposition 20. By Reynolds' transport theorem the area gained by the sweeping region $N_T = \bigcup_{t \leq T} Q_t \cap C_2$ satisfies $\frac{d}{dT}|N_T| \leq \int_{\text{advancing part of } \partial Q_T} v_n ds$, because advance into already-swept territory contributes to the flux but gains no area. Integrating over $T \in [0, \pi/2]$ gives $|N| \leq V$. $\square$

Remark 29 (The size of the gap, measured). Equality in Proposition 28 holds exactly when nothing is re-swept, which is what the three hypotheses of Proposition 20 assert, and which is why $V(\Sigma) = |N(\Sigma)|$. Perturbing H by a smooth bump η supported where $H + H'' \geq \frac{1}{2}$ (so convexity survives) and measuring $|N|$ with a polygon oracle against V from the same H , the excess $V - |N|$, corrected for the oracle's known $O(1/n)$ bias, is

ε	−0.20	−0.10	−0.05	0.05	0.10	0.20
$V - N $	$5.5 \cdot 10^{-3}$	$1.4 \cdot 10^{-4}$	$4.2 \cdot 10^{-6}$	$9.3 \cdot 10^{-5}$	$8.4 \cdot 10^{-4}$	$5.0 \cdot 10^{-3}$

— strictly positive on *both* sides and growing faster than ε^2 . So Σ is an isolated zero of $V - |N|$: the functional Q touches the true bound at Σ and falls strictly below it in every direction. Therefore Q cannot certify Σ , and the fact that Q is tight, critical and concave there (Remark 26) does not yield an optimality statement.

This locates the difficulty exactly. The repair is the one the one-corner argument performs: replace the exact flux V by the flux over a sub-family on which the sweep is *provably* injective, so that the integral is the area of a genuine subset of N and the inequality runs the right way, while the restriction is vacuous at Σ . Producing such a sub-family is the content of that argument's Chapters 3–6 and is not done here.

One reformulation that may help. A point p lies on the face-2 line at parameter t iff $\phi(t) := \langle p, \nu_t \rangle - (G(t) - 1)$ vanishes, and with $s(p, t) := F(t) - 1 - \langle p, \mu_t \rangle$ one computes $\phi' = s - \alpha_2$ and $s' = -\phi - \alpha_1$, i.e. in complex form

$$z := s + i\phi, \quad z' = iz - (\alpha_1 + i\alpha_2).$$

So the whole face-2 sweep is governed by one linear ODE whose homogeneous part is an exact rotation, $z(t) = \zeta(t) + e^{it}w_0$ with w_0 an affine coordinate for p ; the sweep is injective iff no solution meets $\{\phi = 0, 0 \leq s \leq \alpha_2\}$ twice, and at every such meeting $\phi' = s - \alpha_2 \leq 0$, so the crossing is always downward. A second meeting therefore forces an upward crossing in between, at $s \geq \alpha_2$. Since the total rotation over $[0, \pi/2]$ is only $\pi/2$, the homogeneous part alone cannot supply that extra crossing; whether the forcing ζ can is the open point.

Proposition 30 (Monotonicity of the outer arm). *The face-1 segment's far endpoint has x -coordinate*

$$x(t) = c_x(t) + \sigma(t) \sin t = \frac{F(t) - 1}{\cos t}, \quad (25)$$

the x -intercept of the face-1 line, and $\cos^2 t x'(t) = F'(t) \cos t + (F(t) - 1) \sin t$. For Σ :

- on $[0, \beta)$, where $A(t) \equiv P$ so $F(t) = \cos t + \frac{1}{2} \sin t$,

$$x = 1 + \frac{1}{2} \tan t - \sec t, \quad \cos^2 t x' = \frac{1}{2} - \sin t > 0,$$

the last because $\sin \beta = 1/(4a_1) = 0.2856 \dots < \frac{1}{2}$ by Theorem 14, i.e. $\beta < \pi/6$;

- on $(\pi/2 - \beta, \pi/2]$, where $F(t) - 1 = A \cos t + \frac{1}{2} \sin t - \frac{1}{2}$ with $A = 1 - \frac{2}{3}a_1$,

$$x = A - \frac{1}{2} \tan\left(\frac{\pi}{4} - \frac{t}{2}\right), \quad \cos^2 t x' = \frac{1}{2}(1 - \sin t) > 0 \quad (t < \pi/2),$$

and in particular $x(\pi/2) = A = 1 - \frac{2}{3}a_1 = 0.4164750917 \dots$, which is exactly the right endpoint of the corridor floor facet of Remark 27;

- on $[\beta, \pi/2 - \beta]$, substituting $u = t/2 - \pi/8$ and using $-f_2/f_1 = \sqrt{2} - 1 = \tan(\pi/8)$,

$$\cos^2 t \, x' = W(u) := \frac{3K}{4} \sin u + \frac{K}{4} \cos 3u + \frac{1}{2} - \sin\left(2u + \frac{\pi}{4}\right), \quad K = f_1 \sqrt{4 - 2\sqrt{2}},$$

$$\text{on } |u| \leq \pi/8 - \beta/2.$$

Hence x is strictly increasing, given $W > 0$.

Proof. The face-1 line is $\{\langle p, \mu_t \rangle = F(t) - 1\}$, so its x -intercept is $(F - 1)/\cos t$; that it agrees with $c_x + \sigma \sin t$ is $c_x + c_y \tan t = (F - 1)/\cos t$, immediate from $c = (F - 1)\mu_t + (G - 1)\nu_t$. The three closed forms follow by substituting $F(t) - 1 = \operatorname{Re}(e^{-it}z(t))$ with the complex forms of SOL1, SOL6, SOL5. For the middle phase the κ_6 terms cancel and the triple-angle identities $\sin(3t/2) = s(3c^2 - s^2)$, $\cos(3t/2) = c(c^2 - 3s^2)$ with $s = \sin \frac{t}{2}$, $c = \cos \frac{t}{2}$ reduce the result to the stated three-term form. $\square$

Proposition 31 ($W > 0$, by a finite certificate). $W > 0$ on $|u| \leq u_0 := \pi/8 - \beta/2 = 0.2478721712\dots$, so x is strictly increasing on all of $[0, \pi/2]$ and $S_1 = \sigma$.

Proof. $K = 1.3020516916172893452\dots$, and $3u_0 = 0.7436\dots < \pi$, so $\cos 3u$ is unimodal on $[-u_0, u_0]$ with maximum at 0 and its minimum over any subinterval is attained at an endpoint. Also $\sin u$ is increasing, and $2u + \pi/4$ ranges over $[0.2896\dots, 1.2812\dots] \subset [0, \pi/2]$, so $\sin(2u + \pi/4)$ is increasing. Therefore on $[p, q] \subseteq [-u_0, u_0]$,

$$W \geq L(p, q) := \frac{3K}{4} \sin p + \frac{K}{4} \min(\cos 3p, \cos 3q) + \frac{1}{2} - \sin\left(2q + \frac{\pi}{4}\right).$$

Dividing $[-u_0, u_0]$ into 32 equal pieces gives $L > 0$ on every piece, with $\min L = 6.13 \cdot 10^{-3}$; the computation is at 40 working digits, so the margin exceeds the arithmetic error by 37 orders of magnitude. (16 pieces do not suffice: $\min L = -8.6 \cdot 10^{-3}$.) $\square$

Remark 32. W is in fact strictly decreasing, with $W(u_0) = \frac{1}{2}(1 - \cos \beta) = 0.020828595599846555\dots$ at the right junction, matching the last phase's $\frac{1}{2}(1 - \sin t)$ there to $1.7 \cdot 10^{-41}$, as C^1 gluing requires. The three closed forms of Proposition 30 were also checked against numerical differentiation of the path to 10^{-10} .

Remark 33 (The identity at 61 digits). With all constants from their closed forms ($u^3 + 3u = 2$, $\beta = \arctan(u/2)$, $a_1 = \sqrt{4 + u^2}/(4u)$, f_1 from Proposition 6) and 60 working digits,

$$V = 0.184193197088768988251655564606\dots, \quad |C_2| = 2.013341612602978828172120476813\dots,$$

$$Q(\Sigma) - A_R^* = -1.56 \cdot 10^{-61}.$$

The two sides share no computation, so this rules out the possibility that the double-precision agreement of (23) was a rounding coincidence. It remains floating point, not an interval enclosure.

10 A convex subdomain on which the bound is valid

Proposition 28 says $V \geq |N|$ with equality exactly when nothing is re-swept, i.e. when the combined sweep is injective. We now show that injectivity is governed by inequalities that are *linear* in H , so the set where the bound is valid is convex.

Each sweep lies on a line: face 2 at parameter t on $\ell_2(t) = \{\langle p, \nu_t \rangle = G(t) - 1\}$, face 1 on $\ell_1(t) = \{\langle p, \mu_t \rangle = F(t) - 1\}$. Two distinct lines meet in one point, so a double cover forces that point into both segments. Solving $\ell_2(t) \cap \ell_2(t')$ for the coordinate s on $\ell_2(t)$, and using $\langle \mu_t, \nu_{t'} \rangle = \sin(t - t')$,

$$s_{22}(t, t') = \frac{\langle c(t), \nu_{t'} \rangle - (G(t') - 1)}{\sin(t - t')}, \quad s_{11}(t, t') = \frac{\langle c(t), \mu_{t'} \rangle - (F(t') - 1)}{\sin(t' - t)}. \quad (26)$$

The numerator of s_{22} vanishes at $t' = t$ with derivative $-\alpha_2(t)$, so $s_{22}(t, t') \rightarrow \alpha_2(t)$: nearby face-2 lines meet at the *far* end of the face-2 segment $[0, \alpha_2]$. Likewise $s_{11}(t, t') \rightarrow \alpha_1(t)$, the *near* end of the face-1 segment $[\alpha_1^+, \sigma]$.

Definition 34. Let $\mathcal{K}'$ be the set of admissible H satisfying, for all $0 \leq t' < t \leq \pi/2$,

$$(C22) \quad \langle c(t), \nu_{t'} \rangle - (G(t') - 1) \geq \alpha_2(t) \sin(t - t'), \quad (27)$$

$$(C11) \quad \langle c(t), \mu_{t'} \rangle - (F(t') - 1) \leq \alpha_1(t)^+ \sin(t' - t) \quad (\text{read with } \sin(t' - t) < 0), \quad (28)$$

together with the analogous condition for $\ell_2(t) \cap \ell_1(t')$.

Proposition 35. $\mathcal{K}'$ is convex, and on $\mathcal{K}'$ the combined sweep is injective, hence $V = |N|$ and $Q = |C_2| - 2|N| \geq |T|$ for every ambidextrous sofa T with cap data H .

Proof. By Lemma 18, $c(t)$ is affine in H , hence so are $\langle c(t), \nu_{t'} \rangle$, $G(t') - 1$, $F(t') - 1$, $\alpha_1(t)$ and $\alpha_2(t)$; $\sin(t - t')$ does not depend on H . So (27) and (28) are linear inequalities in H and cut out a convex set. Given (27), for any pair $t' < t$ the intersection point of $\ell_2(t)$ and $\ell_2(t')$ has $s_{22}(t, t') \geq \alpha_2(t)$, so it is not interior to the segment at t , and the pair contributes no double cover; (28) does the same for face 1 with the inequality reversed because the relevant endpoint is the near one. With the cross condition, no point is swept twice.

For $V = |N|$: every point of N has a first capture time $\tau(p)$, at which $\partial Q_{\tau(p)}$ is advancing at p (otherwise p would have been captured earlier), so by Lemma 19 it lies on face 2 with $s \leq \alpha_2$ or on face 1 with $s \geq \alpha_1$. Hence N is covered by the two sweeps, and the sweeps lie in N because an advancing face point belongs to $Q_{t+\varepsilon}$. Injectivity turns each $\int |\det|$ into the area of its image, and the two images are disjoint, so the sum is $|N|$. $\square$

Remark 36 ($\Sigma \in \mathcal{K}'$, and $\mathcal{K}'$ has nonempty interior). Measured on a grid of 401 parameters, Σ satisfies all three conditions: of the 158 802 ordered pairs with $|t - t'|$ above the diagonal cutoff, *zero* give a point interior to both face-2 segments and *zero* interior to both face-1 segments, and of 58 081 cross pairs, *zero* land in both. The margins in (27), (28) vanish linearly as $t' \rightarrow t$, which is forced: that is the envelope limit. Away from the diagonal they are strictly positive.

$\mathcal{K}'$ is not a single point. For $H = H_\Sigma + \varepsilon \sin(k\theta)$, which respects the gauge for even k , all conditions still hold with zero violations at $k = 2$ for $|\varepsilon| \leq 3 \cdot 10^{-2}$ and at $k = 4$ for $|\varepsilon| \leq 10^{-2}$, in both signs, and the two margins trade off smoothly. So Σ lies in $\mathcal{K}'$ with low-frequency room around it.

The mechanism of failure is identified: for the compactly supported bump perturbations of Remark 29, where $V - |N|$ was found positive, the face-2 condition continues to hold and it is the *face-1* condition that fails, by $-4.7 \cdot 10^{-2}$ at $\varepsilon = 0.05$ and $-2.3 \cdot 10^{-1}$ at $\varepsilon = 0.20$. Those perturbations have large higher derivatives, which is what creates face-1 envelope self-intersections.

Theorem 37 (Conditional). *Assume:*

- (i) $Q(\Sigma) = A_R^*$, $\delta Q(\Sigma) = 0$ in every admissible direction, and $\delta^2 Q \preceq 0$ on the cell $\mathcal{C}$ of support functions sharing Σ 's sign pattern;
- (ii) the three conditions of Definition 34 hold on $\mathcal{K}' \cap \mathcal{C}$;
- (iii) competitors satisfy the niche separation $M < \frac{1}{2}$ of Corollary 2.

Then $|T| \leq A_R^*$ for every ambidextrous sofa T whose cap data lies in $\mathcal{K}' \cap \mathcal{C}$.

Proof. $\mathcal{C}$ is convex because α_1, α_2 are affine in H , so each pointwise sign condition is a half-space; $\mathcal{K}'$ is convex by Proposition 35; hence so is the intersection. By (i), Q is a concave function on that convex set with a critical point at Σ , so $Q(H) \leq Q(\Sigma) = A_R^*$ throughout. By (ii) and Proposition 35, $Q(H) \geq |T|$ there, using (iii) to get the factor 2 in $|T| \leq |C_2| - 2|N|$. $\square$

Remark 38 (Honest status of Theorem 37). Hypothesis (i) is *numerical*: $Q(\Sigma) = A_R^*$ to 61 digits (Remark 33), $\|\delta Q\|_\infty/h = 1.8 \cdot 10^{-8}$, and $\lambda_{\max}/h \rightarrow -0.72$ for a discretised form. Hypothesis (ii) is measured on a grid, not proved. Hypothesis (iii) is the conditional part of Theorem 11. So the effective label of Theorem 37 is HEURISTIC, and it does not assert optimality of Σ : the class $\mathcal{K}' \cap \mathcal{C}$ is not shown to contain every competitor, and by Remark 29 bodies outside $\mathcal{K}'$ genuinely violate $Q \geq |T|$. What the theorem does is exhibit an explicit convex class, with nonempty interior around Σ , on which the one-corner architecture transfers. Proving (i) and (ii) is the remaining work, and is finite: (i) reduces to a Hessian sign and a first-order identity in closed form, and (ii) to the linear inequalities (27), (28).

11 One curvature condition implies injectivity

The conditions of Definition 34 were verified on a grid. We now prove them, for all pairs at once, from a single linear condition on the cap.

Definition 39. Say H satisfies (RC) if the absolutely continuous part of the surface measure obeys

$$H(\theta) + H''(\theta) \leq 1 \quad \text{for a.e. } \theta \in [0, \pi],$$

the only atoms being the corridor ceiling and floor facets at $\theta = \pm\pi/2$. Equivalently: the cap is nowhere flatter than a circle whose radius is the corridor width.

Since $H + H'' \leq 1$ is linear in H , (RC) cuts out a convex set.

Theorem 40. *If H satisfies (RC), then (27) and (28) hold for every pair $0 \leq t' < t \leq \pi/2$. Hence each of the two sweeps is injective.*

Proof. Fix t and set $\tau = t - t' \in (0, t]$. For (27) put $n(t') = \langle c(t), \nu_{t'} \rangle - (G(t') - 1)$ and

$$\Phi(\tau) = n(t - \tau) - \alpha_2(t) \sin \tau,$$

so that (27) reads $\Phi \geq 0$. Now $n(t) = 0$ because $\langle c(t), \nu_t \rangle = G(t) - 1$, and $n'(t') = -\langle c(t), \mu_{t'} \rangle - G'(t')$ equals $-(F(t) - 1) - G'(t) = -\alpha_2(t)$ at $t' = t$ by (18); hence $\Phi(0) = \Phi'(0) = 0$. Differentiating again and using $\nu'' = -\nu$,

$$n''(t') = -\langle c(t), \nu_{t'} \rangle - G''(t') = -(n(t') + G(t') - 1) - G''(t'),$$

so with $n(t - \tau) = \Phi(\tau) + \alpha_2(t) \sin \tau$ the α_2 terms cancel and

$$\Phi''(\tau) + \Phi(\tau) = 1 - (G + G'')(t - \tau) =: R(\tau). \quad (29)$$

With $\Phi(0) = \Phi'(0) = 0$, (29) gives $\Phi(\tau) = \int_0^\tau \sin(\tau - u) R(u) du$. Since $\tau \leq t \leq \pi/2 < \pi$ we have $\sin(\tau - u) \geq 0$ for $u \in [0, \tau]$, and $R \geq 0$ by (RC) because $G(s) = H(s + \pi/2)$. Therefore $\Phi \geq 0$.

The ceiling atom of H sits at $\theta = \pi/2$, i.e. at $t - u + \pi/2 = \pi/2$, i.e. $u = t$. For $\tau < t$ that is outside $[0, \tau]$, and for $\tau = t$ it is the endpoint, where the kernel $\sin(\tau - u)$ vanishes. So the atom contributes nothing.

For (28) put $m(t') = \langle c(t), \mu_{t'} \rangle - (F(t') - 1)$ and $\Psi(\tau) = m(t - \tau) + \alpha_1(t) \sin \tau$. Then $m(t) = 0$, $m'(t) = \alpha_1(t)$ by (17), so $\Psi(0) = \Psi'(0) = 0$, and $\mu'' = -\mu$ gives in the same way

$$\Psi''(\tau) + \Psi(\tau) = 1 - (F + F'')(t - \tau),$$

whence $\Psi(\tau) = \int_0^\tau \sin(\tau - u) [1 - (H + H'')(t - u)] du \geq 0$. Multiplying by $-\sin \tau < 0$ turns $\Psi \geq 0$ into $s_{11} \leq \alpha_1$, which is (28). Here the atom needs $t - u = \pi/2$, i.e. $u = t - \pi/2 \leq 0$, reachable only at $t = \pi/2$, where $\sigma(\pi/2) = \alpha_1(\pi/2)$ makes the face-1 segment empty and the condition vacuous. $\square$

Remark 41 (Verification). For Σ the absolutely continuous part of $H + H''$ has maximum $0.8385682\dots$, at $\theta = \pi - \beta$, so (RC) holds with margin $0.1614\dots$; the atom at $\theta = \pi/2$ has mass $1.1670498\dots$, the ceiling facet length. The reduction (29) was checked against direct evaluation of $n(t - \tau) - \alpha_2(t) \sin \tau$ at several (t, τ) , agreeing to $2.4 \cdot 10^{-5}$, the finite-difference floor for H'' .

Theorem 40 also explains the failure mode of Remark 36. Across twelve perturbations, (RC) holds exactly when (28) holds: the compactly supported bumps give $\max(H + H'')_{\text{ac}} = 1.08, 1.45, 2.18, 2.67, 4.58, 8.41$ and all violate (28); $H_\Sigma + \varepsilon \sin(2\theta)$ at $|\varepsilon| = 0.03$ gives 0.888 and holds; $H_\Sigma + \varepsilon \sin(4\theta)$ gives 0.976 at $|\varepsilon| = 0.01$ and holds, 1.267 at $|\varepsilon| = 0.03$ and fails. The agreement is exact on both sides of the transition.

Remark 42 (What Theorem 40 does and does not give). It replaces hypothesis (ii) of Theorem 37, for the two self-intersection conditions, by (RC): a single explicit linear inequality with a geometric reading. The cross condition, that $\ell_2(t) \cap \ell_1(t')$ avoid both segments, is not covered; it is measured to hold at Σ (0 of 58 081 pairs) and remains a hypothesis. So the hypothesis class of Theorem 37 may be replaced by $\{(\text{RC})\} \cap \{\text{cross}\} \cap \mathcal{C}$, an intersection of two convex sets with one measured condition.

Remark 43 (The first variation, pointwise). Hypothesis (i) of Theorem 37 contains $\delta Q(\Sigma) = 0$. Integrating every η' in δQ by parts and collecting the coefficient of $\eta(\theta)$ gives the Euler–Lagrange equations of Q :

$$H + H'' = \alpha_2^+ + (\sigma - \alpha_1) \tan \theta - (\sigma - \alpha_1)' + (\alpha_1^-)' \quad (0 < \theta < \pi/2),$$

$$H + H'' = -(\alpha_2^+)'(s) + \alpha_1^-(s), \quad s = \theta - \pi/2 \quad (\pi/2 < \theta < \pi).$$

Evaluated on Σ at eighteen values of θ spanning all six regions, the residual is at most $9.4 \cdot 10^{-6}$, which is the finite-difference floor. So $\delta Q(\Sigma) = 0$ has an explicit pointwise form; proving it amounts to substituting the closed forms of SOL1, SOL5, SOL6 phase by phase, and is not carried out here.

12 The cross condition, the first variation, and a Gårding bound

Theorem 44. *(RC) implies the cross condition: for all t, t' the point $\ell_2(t) \cap \ell_1(t')$ is not interior to both sweep segments. Consequently, under (RC) the combined sweep is injective and $V = |N|$, so $Q = |C_2| - 2|N| \geq |T|$.*

Proof. Work in the frame at t and write $p = c(t) + a\mu_t + b\nu_t$. The equation $\langle p, \nu_t \rangle = G(t) - 1$ forces $b = 0$, so p moves along $\ell_2(t)$ with $s_2 = -a$; the equation $\langle p, \mu_{t'} \rangle = F(t') - 1$ then gives, for $t' = t - \tau$,

$$s_2 = \frac{m(t - \tau)}{\cos \tau}, \quad s_1 = -\Phi(\tau) - (\alpha_2(t) - s_2) \sin \tau,$$

using $\langle \mu_t, \nu_{t-\tau} \rangle = \sin \tau$ and $n(t - \tau) = \Phi(\tau) + \alpha_2(t) \sin \tau$. If $\tau > 0$ and p is interior to the face-2 segment, then $0 < s_2 < \alpha_2(t)$, and $\Phi \geq 0$ by Theorem 40 with $\sin \tau \geq 0$ because $\tau \leq \pi/2$; hence $s_1 < 0 \leq \alpha_1(t')^+$ and p lies below the face-1 segment.

For $t' > t$ work in the frame at t' : the same computation with the roles of the two families exchanged gives, for $t = t' - \tau'$ with $\tau' > 0$,

$$s_2 = -\Psi(\tau') - (s_1 - \alpha_1(t')) \sin \tau',$$

so if p is interior to the face-1 segment then $s_1 > \alpha_1(t')^+ \geq \alpha_1(t')$ and $s_2 < 0$, outside $[0, \alpha_2(t)]$. At $t' = t$ the two lines are perpendicular and meet at $c(t)$, where $s_1 = s_2 = 0$, interior to neither. Both displayed identities were checked against a direct linear solve, agreeing to 10^{-16} . $\square$

So hypothesis (ii) of Theorem 37 may be replaced throughout by (RC).

Theorem 45. $\delta Q(\Sigma) = 0$. *More precisely the Euler–Lagrange equations of Remark 43 hold identically on each of the six regions, as trigonometric identities in a_1, f_1, f_2 — no relation among the constants is used.*

Proof. On each phase, $F(t) - 1 = \operatorname{Re}(e^{-it}z(t))$ and $G(t) - 1 = \operatorname{Im}(e^{-it}z(t))$ evaluate to

$$\begin{array}{ll} \text{SOL1} & F - 1 = \cos t + \frac{1}{2} \sin t - 1, & G - 1 = (2a_1 - 1) \sin t + \frac{1}{2} \cos t - \frac{1}{2}, \\ \text{SOL6} & F - 1 = f_1 \cos \frac{t}{2} + f_2 \sin \frac{t}{2} - 1 + \kappa \cos t + \frac{1}{2} \sin t, & G - 1 = -f_2 \cos \frac{t}{2} + f_1 \sin \frac{t}{2} - 1 + \frac{1}{2} \cos t - \kappa \sin t, \\ \text{SOL5} & F - 1 = (1 - \frac{2}{3}a_1) \cos t + \frac{1}{2} \sin t - \frac{1}{2}, & G - 1 = (\frac{8}{3}a_1 - 1) \sin t + \frac{1}{2} \cos t - 1, \end{array}$$

with $\kappa = 1 - \frac{4}{3}a_1$. Substituting these into $\alpha_1 = G - 1 - F'$, $\alpha_2 = F - 1 + G'$, $\sigma = (F - 1) \tan t + G - 1$ and into the two Euler–Lagrange equations, with the sign pattern of each region fixed ($\alpha_2 > 0$ except on the last phase, $\alpha_1 < 0$ only on the first), every residual reduces identically to 0. $\square$

Remark 46. The computation is a computer algebra verification, six residuals, each returning 0 with a_1, f_1, f_2 free. It was controlled against five deliberately perturbed variants of the middle-phase equation (sign flip on the tan term, each of the two terms dropped, α_1 and α_2 exchanged, $H + H''$ replaced by $H - H''$), all of which return nonzero; and the three pairs of phase formulas were checked against the reference path to $2.2 \cdot 10^{-16}$. The content of Theorem 45 is that Romik's ODE families are exactly the critical-point equations of Q .

Remark 47 (Towards $\delta^2 Q \preceq 0$). The remaining part of hypothesis (i) is the sign of the second variation. The step that makes it accessible is an exact algebraic cancellation. On $[0, \beta]$ write $p = \eta'(t)$, $q = \eta(t)$, $r = \eta(t + \pi/2)$, $T = \tan t$; the three terms of (24) that carry $\eta'(t)$ combine as

$$-p^2 - (qT + p)^2 + (r - p)^2 = -(p + qT + r)^2 + 2r^2 + 2qTr, \quad (30)$$

so the added term is absorbed exactly and the remainder carries no derivative of η . What is left is a lower-order estimate. With $\eta(0) = \eta(\pi/2) = 0$ the relevant Poincaré constants are 4 on $[0, \pi/2]$, 1 on $[\pi/2, \pi]$ (Dirichlet–Neumann, $\eta(\pi)$ free) and $(\pi/2\beta)^2 = 29.41$ on $[0, \beta]$. Retaining a fraction δ of $-\int_0^\beta \eta'^2$ before applying (30) and splitting $(qT + r)^2$ with a parameter κ , the coefficient of $\int_0^\beta \eta'^2$ is

$$\left(\frac{1+\kappa}{1-\delta} - 1\right) \tan^2 \beta + 1 - \delta \left(\frac{\pi}{2\beta}\right)^2 = -1.83 \dots < 0 \quad (\delta = \frac{1}{10}, \kappa = 1),$$

and the r^2 remainder, of coefficient $3.22 \dots$, is absorbed by the E_2 term of (24), which supplies $-\frac{1}{2} \int_{\pi/2}^{\pi-\beta} \eta'^2$: since $\eta(\pi/2) = 0$ gives $\int_{\pi/2}^{\pi-\beta} \eta'^2 \leq (\beta^2/2) \int \eta'^2$, the requirement is $3.22 \cdot 0.0420 = 0.135 \leq \frac{1}{2}$. So every ingredient of a Gårding inequality $\delta^2 Q \leq -c \|\eta\|_{L^2}^2$ is in place; assembling the chain with explicit constants is not carried out here.

Measured, the constant is $c = 0.7285 \dots$: the supremum of $\delta^2 Q[\eta] / \|\eta\|_{L^2}^2$ over the hat space, as a symmetric generalised eigenvalue problem against the exact mass matrix, is $-0.7059, -0.7188, -0.7255, -0.7285$ at $m = 32, 64, 128, 256$.

13 Concavity, and the area identity

Theorem 48. *On the cell $\mathcal{C}$ of Σ 's sign pattern, $\delta^2 Q \leq 0$, with equality only at $\eta = 0$. Explicitly, writing $J_1 = \int_0^\beta \eta'^2$, $J_2 = \int_\beta^{\pi/2} \eta'^2$, $J_3 = \int_{\pi/2}^\pi \eta'^2$,*

$$\frac{1}{2} \delta^2 Q[\eta] \leq -1.6013 J_1 - 0.3710 J_2 + 0 \cdot J_3 - 0.005 \int_{\pi/2}^{\pi-\beta} \eta'^2. \quad (31)$$

Proof. On $[0, \beta]$ retain a fraction δ of $-\int_0^\beta \eta'^2$ and apply (30) with the remaining $-(1-\delta)\eta'^2$; completing the square in $p = \eta'(t)$ and discarding the resulting negative square,

$$-(1-\delta)p^2 - (qT + p)^2 + (r - p)^2 \leq \frac{(qT + r)^2}{1-\delta} - q^2 T^2 + r^2 \leq A q^2 T^2 + B r^2,$$

with $A = \frac{1+\kappa}{1-\delta} - 1$, $B = \frac{1+1/\kappa}{1-\delta} + 1$ from $(qT + r)^2 \leq (1+\kappa)q^2 T^2 + (1+1/\kappa)r^2$, and $T = \tan t \leq \tan \beta$ on $[0, \beta]$. The three Poincaré constants for $\eta(0) = \eta(\pi/2) = 0$ are $P_1 = (\pi/2\beta)^2 = 29.4135$ on $[0, \beta]$, $P_2 = (\pi/2(\pi/2 - \beta))^2 = 1.5035$ on $[\beta, \pi/2]$ and $P_3 = 1$ on $[\pi/2, \pi]$, each with a Dirichlet endpoint and a free one. Finally $-(\eta(t) + \eta'(t + \pi/2))^2 \leq -\lambda \eta'(t + \pi/2)^2 + \frac{\lambda}{1-\lambda} \eta(t)^2$ turns the E_2 term into $-\lambda \int_{\pi/2}^{\pi-\beta} \eta'^2 + \frac{\lambda}{1-\lambda} (J_1 + J_2)$, and $\eta(\pi/2) = 0$ gives $\int_{\pi/2}^{\pi-\beta} \eta'^2 \leq (\beta^2/2) \int_{\pi/2}^{\pi-\beta} \eta'^2$, so the $B r^2$ remainder is absorbed as soon as $\lambda \geq B\beta^2/2$. Collecting,

$$\begin{aligned} \text{coefficient of } J_1 &= A \tan^2 \beta - \delta P_1 + 1 + \frac{\lambda}{1-\lambda}, & \text{of } J_2 &= 1 - P_2 + \frac{\lambda}{1-\lambda}, \\ \text{of } J_3 &= 1 - P_3 = 0, & \text{of } \int_{\pi/2}^{\pi-\beta} \eta'^2 &= B\beta^2/2 - \lambda. \end{aligned}$$

At $\delta = \frac{1}{10}$, $\kappa = 2$, $\lambda = 0.117$ these are -1.6013 , -0.3710 , 0 and -0.005 , giving (31). For strictness, equality forces $\eta \equiv 0$ on $[0, \pi/2]$ and η the first Dirichlet–Neumann mode on $[\pi/2, \pi]$; but with $\eta \equiv 0$ on $[0, \pi/2]$ the E_2 term contributes $-\int_0^{\pi/2-\beta} \eta'(t + \pi/2)^2 < 0$. $\square$

The bound $\delta^2 Q \leq -0.7419 \|\eta\|_{L^2(0, \pi/2)}^2$ that (31) yields may be compared with the measured supremum of $\delta^2 Q / \|\eta\|_{L^2(0, \pi)}^2$, which is $-0.7285\dots$; the two agree to the extent the proof’s J_3 coefficient is slack.

Proposition 49. $\sigma - \alpha_1 = \cos t \, x'(t)$ on each phase, with $x = (F - 1)/\cos t$ as in (25). Hence the middle term of (20) is $\frac{1}{2} \int \cos^2 t \, x'(t)^2 dt$, an energy of the intercept path.

Proof. $\sigma - \alpha_1 = (F - 1) \tan t + F'$ by (17) and (19), and $\cos^2 t \, x' = F' \cos t + (F - 1) \sin t$ by (25); dividing by $\cos t$ gives the claim. Verified symbolically on SOL1, SOL5 and SOL6, residual 0 in each case, which also recovers the first and third closed forms of Proposition 30. $\square$

Remark 50 (The area identity). Since $u = 2 \tan \beta$, Romik’s area is

$$A_R^* = u^2 + 1 + \arctan(u/2) = 1 + 4 \tan^2 \beta + \beta.$$

Using Proposition 49 to make every integrand elementary, the six blocks of $|C_2|$ and the three blocks of V evaluate at 50 working digits to $|C_2| = 2.0133416126029788281721\dots$, $V = 0.1841931970887689882517\dots$, and

$$Q(\Sigma) - (1 + 4 \tan^2 \beta + \beta) = -2.7 \cdot 10^{-51}.$$

A symbolic evaluation of the same integrals was attempted and did not terminate; the identity therefore remains HEURISTIC, now against a closed-form target and with a phase-by-phase decomposition.

Remark 51 (Status of Theorem 37). Hypothesis (ii) is Theorems 40 and 44, proved from (RC). Hypothesis (i) splits: $\delta Q(\Sigma) = 0$ is Theorem 45, $\delta^2 Q \preceq 0$ is Theorem 48, and $Q(\Sigma) = A_R^*$ is Remark 50, still numerical. Hypothesis (iii), the separation $M < \frac{1}{2}$ for competitors, is untouched; across a family of low-frequency perturbations no competitor satisfying (RC) was found with $M \geq \frac{1}{2}$, but no implication is proved. So the conditional theorem now rests on one 51-digit numerical identity and one geometric hypothesis, rather than on four measured conditions. Optimality of Σ remains open.

14 Formalisation

Lemma 1, Corollary 2 and the collapse of (3) are machine-checked in Lean 4 without Mathlib as `niche_below_apex`, `niche_disjoint` and `union_area_of_disjoint`. The identity underlying Σ ’s middle-phase equation is also formalised: writing D for the formal derivative in the half-angle, $D(Dv) = -(v + 1)$ for every arc of constant term -1 , which is $4v'' = -(v + (1, 1))$ for the bracket of (4) (`Trig.sol6_bracket`, proved by `rfl` and depending on no axioms), as is the frame-speed computation behind Theorem 14 (`Trig.sol1_speed_mu`) and the covering criterion of Theorem 11 (`strip_covers_iff`). `lake build` is clean with no `sorry`, and `#print axioms` reports nothing beyond `propext` and `Quot.sound`. Trigonometric quantities are carried as formal symbols with the sign hypotheses $\sin t \geq 0$, $\cos t \geq 0$, which is all Lemma 1 uses.

References

- [1] J. Baek, *Optimality of Gerver's sofa*, arXiv:2411.19826, 2024.
- [2] J. L. Gerver, *On moving a sofa around a corner*, Geom. Dedicata **42** (1992), 267–283.
- [3] D. Romik, *Differential equations and exact solutions in the moving sofa problem*, Exp. Math. **27** (2018), 316–330.