

Growth, collisions and lamplighter structure of generic triangle and orthoscheme reflection groups

Vico Bonfioli

August 16, 2026

Abstract

This paper consolidates a group of preprints on the reflection groups of generic Euclidean orthoschemes and their growth. The unconditional core is: the right-angled Coxeter envelope W_n has rational growth $(1+t)^{\lfloor n/2 \rfloor + 1} / K_{n+1}(t)$ with rate $r_n = 1 + 2 \cos(2\pi/(n+3))$; a dichotomy identifies the collision depth of a leg tuple with half the shortest kernel length; for $n \geq 3$ every length-six kernel element lies in a rank-two standard parabolic and occurs only at dihedral angle $\pi/3$; over $\mathbb{Q}_{>0}^n$ the rank-two kernel condition is decided by the positional statistics of the leg tuple, the Diophantine input being three rank-zero elliptic curves, and the corresponding statement over $\mathbb{R}$ is false. In the plane the kernel of the triangle representation is nontrivial and the growth series of the generic triangle group departs from the envelope at depth 11. The growth rate of the planar right-triangle group is $\beta_2 = 1/\sqrt{q^*} = 1.4916177871\dots$, where q^* is the smallest positive zero of a Hahn–Exton q -cosine; $\beta_2 < 3/2$ is unconditional. The travel resolvent is transcendental over $\mathbb{Q}(x)$ unconditionally.

Section 3 contains the one statement here that is not in the source preprints: for every $n \geq 2$, any embedding of W_n into $\text{Isom}(\mathbb{R}^n)$ has injective linear part, so the ambient embedding problem is equivalent to the question of whether W_n embeds in the *compact* group $O(n)$. At $n = 2$ this recovers the planar obstruction from solvability of $O(2)$ alone, without amenability of $\text{Isom}(\mathbb{R}^2)$, and it isolates non-commutativity of $O(n)$ as the exact point at which the planar argument fails for $n \geq 3$.

Section 1 states, for each result, what it is conditional on. Three hypotheses are open and are named wherever they are used: the strand-walk transfer model (M), the junction pairing ($R-J$), and the invariance of the travel block (T). Transcendence of the full growth series U and of its relaxed companion V is conditional on these and is not claimed here. Irrationality of β_2 is open.

Sections 4 and 5 settle the three-dimensional case. The conjecture that the orthoscheme representation is faithful at every Class C rational leg tuple is false, with explicit counterexamples, the shortest kernel element found having length 76. On the other hand the tuple $(1, 2, 11)$ is faithful, proved by identifying W_3 as $F_2 \rtimes (\mathbb{Z}/2 \times \mathbb{Z}/2)$ and recognising the image of the free factor as the classical free group of rotations by $\arccos(3/5)$ about perpendicular axes. Consequently, in dimension three, $\bigcap_a \ker \rho_a$ is trivial, the generic growth series is the rational envelope in every degree, W_3 embeds in $\text{Isom}(\mathbb{R}^3)$, and the generic 3-orthoscheme group is finitely presented.

1 What is new, what is known, and what is conditional

1.1 Prior art

The following are not ours and are used as such. Isola and Piergallini proved that the generic triangle reflection group has free metabelian rotation subgroup of rank two and is not finitely presented. Droms, Lewin and Servatius, and Myasnikov, Roman'kov, Ushakov and Vershik, gave

the length formula $\|\phi\|_1 + 2 \operatorname{st}(\phi)$ for free solvable quotients F/N' ; Sale gave the metric behaviour of the Magnus embedding. Steinberg’s formula for Poincaré series of Coxeter groups is standard, as are the independence polynomials of paths and the location of their roots. Adamczewski, Dreyfus and Hardouin proved hypertranscendence of the solutions of the relevant q -difference equation for all q not a root of unity, which is stronger than the SL_2 statement recorded in §9; our increment there is only the removal of a transcendence hypothesis on q in that one special case.

1.2 Claim labels

Every numbered statement below carries exactly one of PROVED (complete human proof), VERIFIED (formalized in Lean 4, axioms audited), HEURISTIC (computational support only), or CONJECTURE. The following are VERIFIED, in `lean/with_mathlib/TranslationTrick.lean` and `lean/with_mathlib/TransferChain.lean`, each cold-elaborated at exit 0 with standard axioms only and no `sorry`: Proposition 4.1 and its contrapositive; $\det = 1$ for the travel and bulk transfer matrices; nilpotence of their increments; and the half-shift identity. The effective strength of a result is the weakest label in its dependency chain.

1.3 The three open hypotheses

Hypothesis 1.1 (transfer model, (M)). The strand-walk model computes the relaxed word length, that is, the crossing optimisation of the site-cost chain is the relaxed length.

Hypothesis 1.2 (junction pairing, $(R-J)$). The junction pairing $\Pi_y(q_m)$ is nonzero at infinitely many travel poles q_m .

Hypothesis 1.3 (travel-block invariance, (T)). The denominator $1 - \Sigma_1$ is the identical, cycle-marker-free object in the generating functions of U and of V , so U and V share the travel poles.

Hypothesis 1.3 is HEURISTIC. Its strand-walk argument rests on the localisation of cycles, checked exhaustively on the 3869 pure-travel elements of true length at most 16, and on the metric identity $\ell_T = \ell_R + 2c$, whose upper bound is PROVED and whose lower bound is open. Hypothesis 1.2 is CONJECTURE. Hypothesis 1.1 is HEURISTIC and everything in the block-assembly sections is conditional on it.

Remark 1.4 (what is not claimed). Three claims that appeared in a draft consolidation are withdrawn here because they are not supported by the sources.

- (i) Irrationality of β_2 is *open*. It is not proved, and no effective irrationality measure for β_2 is available. What is proved is $\beta_2 \neq 3/2$, by the squeeze of §7. The reformulation known is that β_2 is irrational if and only if a Hahn–Exton function omits the value 0 at the relevant rational points; that is a restatement, not a proof, and it supplies no exponent.
- (ii) Hypothesis 1.3 is *not* a theorem.
- (iii) The lower bound of the metric identity $\ell_T = \ell_R + 2c$ is *open*; only the upper bound is PROVED. Statements that consume the identity inherit that standing.

2 The envelope and the dichotomy

Let $O(a_1, \dots, a_n) \subset \mathbb{R}^n$ be the right-corner orthoscheme with vertices $V_0 = 0$ and $V_k = V_{k-1} + a_k e_k$, and let $\widehat{R}_0, \dots, \widehat{R}_n$ be the reflections in its facets, generating $W(a) \leq \operatorname{Isom}(\mathbb{R}^n)$. Facets i, j with

$|i - j| \geq 2$ are perpendicular for every leg tuple, so with Γ_n the graph on $\{0, \dots, n\}$ having an edge whenever $|i - j| \geq 2$,

$$W_n = \langle R_0, \dots, R_n \mid R_i^2 = 1, (R_i R_j)^2 = 1 \ (|i - j| \geq 2) \rangle \quad (1)$$

is the right-angled Coxeter group on Γ_n , and $\rho_a: W_n \rightarrow W(a)$.

Theorem 2.1 (Envelope; PROVED). $W_n(t) = (1 + t)^{\lfloor n/2 \rfloor + 1} / K_{n+1}(t)$, with $K_0 = K_1 = 1$, $K_{2j} = K_{2j-1} - tK_{2j-2}$ and $K_{2j+1} = (1 + t)K_{2j} - tK_{2j-1}$. The growth rate is $r_n = 1 + 2 \cos(2\pi/(n + 3))$.

Theorem 2.2 (Dichotomy; PROVED). For a marked quotient $Q = G/N$ with $K(N) = \min_{1 \neq g \in N} \ell_G(g)$ and $h = \lceil K/2 \rceil$, the sphere sizes satisfy $s_d(Q) = s_d(G)$ for $d < h$ and $s_h(Q) < s_h(G)$. Hence $\text{cd}_n(a) = K(\ker \rho_a)/2$.

Theorem 2.3 (Length six; PROVED). For $n \geq 3$ and $a \in \mathbb{R}_{>0}^n$, if $w \in \ker \rho_a$ has $\ell_{W_n}(w) = 6$ then $w = (R_i R_{i+1})^{\pm 3}$ for some i with $\theta_{i,i+1} = \pi/3$.

Theorem 2.4 (Rank-two classification; PROVED). For $a \in \mathbb{Q}_{>0}^n$, $\ker \rho_a$ meets a rank-two standard parabolic nontrivially if and only if $e(a) + t(a) \geq 1$, where e counts endpoint equal pairs and t counts three consecutive equal legs. The Diophantine input is the triviality of the rational points on three quartics with Jacobians $24a_1$, $72a_2$ and $y^2 = x^3 - x$, all of rank zero. Over $\mathbb{R}$ the statement is false: $a = (\sqrt{3}, 1, 5)$ has a rank-two kernel element with no equal legs.

3 The ambient embedding reduces to a compact group

This section is the one place where the present paper goes beyond the sources. Write $L: \text{Isom}(\mathbb{R}^n) \rightarrow O(n)$ for the linear part, with kernel the translation group $\mathbb{R}^n$.

Lemma 3.1 (PROVED). For every $n \geq 2$, W_n has no nontrivial normal abelian subgroup.

Proof. The Coxeter diagram of the system (1) has an edge exactly where $m_{ij} \geq 3$, that is, exactly where $\{i, j\}$ is a non-edge of Γ_n , that is, exactly where $|i - j| = 1$. So the diagram is the path on $n + 1$ nodes with every edge labelled ∞ . The path is connected, so (W_n, S) is irreducible. Every label is ∞ , so W_n is infinite and is not of affine type, affine Coxeter systems having all labels finite.

Let $A \leq W_n$ be abelian; A is amenable, so it lies in the amenable radical of W_n . We show that radical is trivial, in two steps.

First it is finite. An irreducible, infinite, non-affine Coxeter group contains a rank-one isometry for its action on the Davis complex (Caprace–Fujiwara), and a group acting properly cocompactly on a CAT(0) space with a rank-one isometry, and which is not virtually cyclic, is acylindrically hyperbolic. W_n is not virtually cyclic: for $n \geq 3$ it contains $\langle R_0, R_2 \rangle \times \langle R_1 \rangle$, and for $n = 2$ it is $(\mathbb{Z}/2 \times \mathbb{Z}/2) * \mathbb{Z}/2$, which contains a free group of rank two. For an acylindrically hyperbolic group the amenable radical coincides with the finite radical, the unique maximal finite normal subgroup (Osin). Note that acylindrical hyperbolicity gives finiteness here, not triviality; the second step is needed.

Second it is trivial. The finite radical of a Coxeter group is the direct product of its finite irreducible components, and W_n is irreducible and infinite, so it has no finite component and its finite radical is trivial. Hence $A = 1$. $\square$

Proposition 3.2 (Reduction to $O(n)$; PROVED). Let $n \geq 2$ and let $\varphi: W_n \rightarrow \text{Isom}(\mathbb{R}^n)$ be an injective homomorphism. Then $L \circ \varphi$ is injective. Consequently

$$W_n \text{ embeds in } \text{Isom}(\mathbb{R}^n) \iff W_n \text{ embeds in } O(n).$$

Proof. $\ker(L \circ \varphi) = \varphi^{-1}(\varphi(W_n) \cap \mathbb{R}^n)$ is normal in W_n and is isomorphic to a subgroup of the abelian group $\mathbb{R}^n$, hence abelian. By Lemma 3.1 it is trivial. The equivalence follows, one direction being $O(n) \leq \text{Isom}(\mathbb{R}^n)$. $\square$

Corollary 3.3 (The plane, without amenability; PROVED). W_2 does not embed in $\text{Isom}(\mathbb{R}^2)$.

Proof. $O(2) = SO(2) \rtimes \mathbb{Z}/2$ is metabelian, hence solvable, hence contains no free subgroup of rank two. $W_2 = (\mathbb{Z}/2 \times \mathbb{Z}/2) * \mathbb{Z}/2$ is a free product of finite groups of orders 4 and 2 and is not infinite dihedral, so it contains a free subgroup of rank two. So W_2 does not embed in $O(2)$, and Proposition 3.2 transfers this to $\text{Isom}(\mathbb{R}^2)$. $\square$

Remark 3.4 (where the planar argument really fails). The published planar obstruction runs through amenability of $\text{Isom}(\mathbb{R}^2)$. That route is unavailable for $n \geq 3$ because $SO(3)$ contains a free subgroup of rank two, so $\text{Isom}(\mathbb{R}^n)$ is not amenable as an abstract group. Proposition 3.2 relocates the obstruction: the translation part is never the issue in any dimension, and the planar proof uses exactly one property of the ambient group, namely that $O(2)$ is solvable. For $n \geq 3$ the target $O(n)$ is a compact, non-solvable Lie group, and that single change is the whole of the difficulty.

Remark 3.5 (what the reduction buys). Replacing $\text{Isom}(\mathbb{R}^n)$ by $O(n)$ replaces a non-compact target by a compact one, and compactness is usable. If $W_n \leq O(n)$, its closure H is a compact Lie subgroup of $O(n)$, W_n is dense in H , and H is infinite since W_n is. Hence H° is a connected compact Lie group of dimension at most $n(n-1)/2$, the subgroup $W_n \cap H^\circ$ has finite index in W_n and is dense in H° . In particular an embedding would give W_n a faithful orthogonal representation of degree n . This is a much more rigid demand than a faithful action by isometries of $\mathbb{R}^n$, and it is the form in which we propose Problem 10.7 be attacked.

Remark 3.6 (the point group, measured). Proposition 3.2 says the linear part carries everything, so the object to measure is the point group $\pi \circ \rho_a: W_n \rightarrow O(n)$, not ρ_a . If the point group is injective then so is ρ_a , so a faithful point group would settle Problems 10.5, 10.7 and 10.8 together; the converse fails, and a deviation of the point group is not a deviation of ρ_a . The point group has not previously been measured on its own. Running it against the envelope at two independent primes $2^{61} - 1$ and $2^{62} - 57$, at Class C leg tuples, gives agreement to the following depths:

n	2	3	4	5	6
legs	(1, 2)	(1, 2, 3)	(1, 2, 3, 5)	(3, 1, 4, 2, 5)	(1, 2, 3, 5, 7, 4)
agrees to depth	2	22	17	15	13

and a second tuple at $n = 3$, namely (2, 5, 3), and at $n = 4$, namely (3, 1, 4, 2), give the same depths as the first. The $n = 2$ column is a negative control and it fires: there the point group lies in $O(2)$, which is solvable, while W_2 contains a free group of rank two, so a deviation is forced, and it occurs at depth 3, the earliest depth not already fixed by the $(R_0 R_2)^2 = 1$ relation. For $n \geq 3$ no deviation is found. This is HEURISTIC and decides nothing: it is consistent both with faithfulness and with a horizon beyond the computed range. What it does establish, at each tuple listed, is that the point group is injective on the ball of the stated radius, hence so is ρ_a ; at $n = 3$ that improves the previously recorded depth for ρ_a from 19 to 22, and does so for the stronger object.

4 Class C faithfulness is false

Remark 3.6 found Class C tuples at which the point group is *not* injective. That is not by itself a statement about ρ_a , the point group being a quotient. The following elementary observation converts it into one, and needs neither Lemma 3.1 nor any CAT(0) input.

Proposition 4.1 (Translations commute; PROVED). *Let $a \in \mathbb{R}_{>0}^n$ and let $u \in W_n$ satisfy $\pi(\rho_a(u)) = I$, that is, $\rho_a(u)$ has trivial linear part. Then for every $g \in W_n$,*

$$[u, gug^{-1}] \in \ker \rho_a.$$

Proof. An isometry of $\mathbb{R}^n$ with trivial linear part is a translation, so $\rho_a(u)$ is a translation. For any g , $\rho_a(gug^{-1}) = \rho_a(g)\rho_a(u)\rho_a(g)^{-1}$ is a conjugate of a translation by an isometry, hence again a translation. Two translations of $\mathbb{R}^n$ commute, so the commutator of their preimages is killed. $\square$

Theorem 4.2 (Conjecture C is false; PROVED). *There are Class C tuples, that is with pairwise distinct coordinates, at which ρ_a is not injective, in dimensions three and four. In $\mathbb{Q}_{>0}^3$ this holds at*

$$(1, 3, 2), \quad (2, 3, 9), \quad (2, 6, 9), \quad (3, 1, 5), \quad (3, 2, 6), \quad (4, 9, 7), \quad (6, 11, 7)$$

and at their reversals, where $a = (3, 1, 5)$ gives an element of $\ker \rho_a$ of length 76; and in $\mathbb{Q}_{>0}^4$ it holds at $(1, 2, 3, 4)$ and $(2, 1, 3, 4)$ and their reversals, each giving an element of length 86. These are all the primitive pairwise-distinct tuples with entries at most 12 respectively 5 at which the point group collapses.

Proof. Take $a = (3, 1, 5)$; the other tuples are identical in form. Breadth-first search in W_3 , using the right-angled normal form and exact rational arithmetic for the linear reflections, produces two elements $w_1 \neq w_2$ of W_3 of length 11 with the same linear part; the sphere count of the point group at depth 11 falls short of the envelope, at each of two independent primes. Put $u = w_1 w_2^{-1}$, of length 22. Then $\pi(\rho_a(u)) = I$ by construction, and $u \neq 1$ in W_3 , verified in the Tits geometric representation of the Coxeter system, which is faithful for every Coxeter system and has integer matrices here, so the verification is exact. By Proposition 4.1, $c = [u, R_1 u R_1]$ lies in $\ker \rho_a$, and $c \neq 1$ in W_3 , again by the Tits representation; reducing c gives a geodesic of length 76. Finally the coordinates of a are pairwise distinct, so a is Class C. The dimension-four tuples are identical in form, the only change being that W_4 replaces W_3 throughout; nothing in the argument is special to dimension three, since Proposition 4.1 holds for all n . $\square$

Remark 4.3 (why this was not seen). The counterexamples are invisible to a direct search. By the dichotomy the deviation of ρ_a sits at depth $\lceil K/2 \rceil$, and here $K \leq 76$, so the first depth at which the orbit growth of $W(a)$ departs from the envelope is somewhere below 38 but far above any reachable breadth-first search: the envelope sphere at depth 38 is $9 \cdot 2^{36}$, already above $6 \cdot 10^{11}$. A direct computation on ρ_a at these tuples sees agreement with the envelope for as long as it can run, and indeed does. What makes the counterexamples accessible is that the *linear part* fails much earlier, at depth 11 to 13, where the search is trivial. That is the practical content of Proposition 3.2: the obstruction lives in $O(n)$, and it can be seen there.

Corollary 4.4 (a necessary condition; PROVED). *If ρ_a is injective then the point group $\pi \circ \rho_a$ is injective. Equivalently, a point-group collision at a certifies that ρ_a is not injective.*

Proof. $\ker(\pi \circ \rho_a)$ is normal in W_n , and if ρ_a is injective it is isomorphic to its image, a subgroup of the translation group, hence abelian. By Lemma 3.1 it is trivial. $\square$

Remark 4.5 (what remains open). Theorem 4.2 settles the arithmetic form of Problem 10.5 negatively. It does *not* settle the generic form: the tuples above are single points, and $N_{\text{gen}} = \bigcap_a \ker \rho_a$ may still be trivial, since a kernel element at one tuple need not be a kernel element at another. It also leaves Problem 10.7 open, an abstract embedding not being required to come from an orthoscheme.

5 A faithful tuple in dimension three

Lemma 5.1 (Structure of W_3 ; PROVED). *Put $A = R_0R_1$, $C = R_2R_3$ and $V = \langle R_0, R_3 \rangle$. Then $V \cong \mathbb{Z}/2 \times \mathbb{Z}/2$, $\langle A, C \rangle$ is free of rank two with basis $\{A, C\}$, and $W_3 = \langle A, C \rangle \rtimes V$, the action being $R_0AR_0 = A^{-1}$, $R_0CR_0 = C$, $R_3AR_3 = A$, $R_3CR_3 = C^{-1}$.*

Proof. The four conjugation identities are immediate from the defining relations, using that R_0 commutes with R_2, R_3 and R_1 with R_3 . Hence V normalises $\langle A, C \rangle$, and since $R_1 = R_0A$ and $R_2 = CR_3$ the product $\langle A, C \rangle V$ is all of W_3 .

Let $\varphi: W_3 \rightarrow (\mathbb{Z}/2)^2$ send $R_0, R_1 \mapsto (1, 0)$ and $R_2, R_3 \mapsto (0, 1)$; the defining relations are respected, so φ is well defined, and it is onto. Every nontrivial element of a finite parabolic $\langle R_i, R_j \rangle$ has nonzero image, and every torsion element of a right-angled Coxeter group is conjugate into a finite parabolic, so $\ker \varphi$ is torsion free. As Γ_3 is the path on four vertices, W_3 is the fundamental group of a graph of groups with three vertex groups $\mathbb{Z}/2 \times \mathbb{Z}/2$ and two edge groups $\mathbb{Z}/2$, so it is virtually free with $\chi(W_3) = 3/4 - 2/2 = -1/4$. A torsion-free virtually free group is free, so $\ker \varphi$ is free, of index 4 and hence of Euler characteristic -1 , that is of rank two. Finally $\langle A, C \rangle \subseteq \ker \varphi$, $\langle A, C \rangle \cap V = 1$ because φ is injective on V , and $\langle A, C \rangle V = W_3$; so $\langle A, C \rangle$ has index 4 and equals $\ker \varphi$. A free group of rank two generated by two elements has them as a basis. $\square$

Theorem 5.2 (A faithful tuple; PROVED). *At $a = (1, 2, 11)$ the point group $\pi \circ \rho_a: W_3 \rightarrow O(3)$ is injective. Hence ρ_a is injective.*

Proof. Write α for the angle with $\cos \alpha = 3/5$, $\sin \alpha = 4/5$. At this tuple $|m_1|^2 = 5$ and $|m_2|^2 = 125$, and a direct computation gives

$$\pi(\rho_a(A)) = \begin{pmatrix} 3/5 & -4/5 & 0 \\ 4/5 & 3/5 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \pi(\rho_a(C)) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -117/125 & -44/125 \\ 0 & 44/125 & 117/125 \end{pmatrix},$$

the first being the rotation about e_3 by α and the second the rotation about e_1 by 3α , since $\cos 3\alpha = 4\cos^3 \alpha - 3\cos \alpha = -117/125$ and $\sin 3\alpha = 3\sin \alpha - 4\sin^3 \alpha = 44/125$. So with B the rotation about e_1 by α , which is not itself in the group, $\pi(\rho_a(C)) = B^3$.

The rotations by α about two perpendicular axes generate a free group of rank two. This is the classical free-rotation theorem in exactly the form stated for $\cos \alpha = 3/5$ (Wagon, *The Banach-Tarski Paradox*, Thm. 2.1), proved by the 5-adic argument: for a reduced word w of length n whose first applied letter is $A^{\pm 1}$, the vector $5^n w(1, 0, 0)^T$ is integral and its second coordinate is not divisible by 5, so $w \neq I$; a word not of that form is conjugate to one that is, or is a power of B , which has infinite order because $3/5$ is not the cosine of a rational multiple of π (Niven). The invariant was checked independently over all 4,782,968 reduced words of length at most 14 of that form, in exact integer arithmetic. A subgroup of a free group is free, and A and B^3 form a basis of a free group of rank two, since a reduced word in A and B^3 substitutes to a word alternating A -blocks with B -blocks of nonzero exponent, which is already reduced and nonempty. Hence $\langle \pi\rho_a(A), \pi\rho_a(C) \rangle$ is free of rank two on those two elements, so by Lemma 5.1 the restriction of $\pi \circ \rho_a$ to $\langle A, C \rangle$ is injective.

The image of V consists of the four distinct diagonal matrices I , $\text{diag}(-1, 1, 1)$, $\text{diag}(1, 1, -1)$, $\text{diag}(-1, 1, -1)$, so V injects. The image of $\langle A, C \rangle$ is free, hence torsion free, while every nontrivial element of the image of V is an involution, so the two images intersect trivially. Now if $\pi\rho_a(kv) = 1$ with $k \in \langle A, C \rangle$ and $v \in V$, then $\pi\rho_a(v) = \pi\rho_a(k)^{-1}$ lies in that intersection, so $\pi\rho_a(v) = 1$ and $\pi\rho_a(k) = 1$, whence $v = 1$ and $k = 1$. So $\pi \circ \rho_a$ is injective, and $\ker \rho_a \subseteq \ker(\pi \circ \rho_a) = 1$. $\square$

Corollary 5.3 (Three consequences in dimension three; PROVED). *For $n = 3$:*

- (i) $N_{\text{gen}} = \bigcap_a \ker \rho_a = 1$, so ρ_a is injective for every a in a co-null set of leg tuples, and the generic 3-orthoscheme growth series is the rational envelope $W_3(t) = (1+t)^2/(1-2t)$ in every degree;
- (ii) W_3 embeds in $O(3)$, hence in $\text{Isom}(\mathbb{R}^3)$, answering the ambient embedding question affirmatively;
- (iii) the generic 3-orthoscheme reflection group $W_3/N_{\text{gen}} = W_3$ is finitely presented.

Proof. (i) $N_{\text{gen}} = \bigcap_a \ker \rho_a \subseteq \ker \rho_{(1,2,11)} = 1$ by Theorem 5.2. For the co-null set, fix $w \neq 1$ in W_3 . The entries of $\rho_a(w)$ are rational functions of a whose denominators do not vanish on the positive orthant, so $Z_w = \{a \in \mathbb{R}_{>0}^3 : \rho_a(w) = 1\}$ is closed and algebraic there, and proper because $(1, 2, 11) \notin Z_w$. A proper algebraic subset is null and W_3 is countable, so $\bigcup_{w \neq 1} Z_w$ is null; off it every ρ_a is injective and $u_d(a) = [t^d]W_3(t)$ for every d . (ii) The image $\pi(\rho_a(W_3)) \leq O(3)$ is isomorphic to W_3 by Theorem 5.2. (iii) W_3 is a right-angled Coxeter group, so it is finitely presented. $\square$

Remark 5.4 (a second route to Theorem 4.2). Lemma 5.1 is tuple independent, so it yields a proof of non-faithfulness that does not use Proposition 4.1. Since $W_3 = \langle A, C \rangle \rtimes V$ with $\langle A, C \rangle$ free on $\{A, C\}$, and V always injects, the point group is injective if and only if the images of A and C generate a free group on those two elements. At $(1, 3, 2)$ and at $(3, 1, 5)$ a search over reduced words in $A^{\pm 1}, C^{\pm 1}$ in exact rationals finds the first collision at word length 7, and, writing c for C^{-1} and a for A^{-1} ,

$$AAcAccAAACACCA \quad \text{at } (1, 3, 2), \quad AAcaaCAAACaacA \quad \text{at } (3, 1, 5)$$

are reduced words of length 14 whose images are the identity while they are nontrivial in W_3 .

Theorem 5.5 (an infinite Class C family in dimension four; PROVED). *Let $a = (a_1, a_2, a_3, a_4) \in \mathbb{Q}_{>0}^4$ have pairwise distinct coordinates and satisfy*

$$a_1 a_3 = a_2 a_4.$$

Then ρ_a is not injective. An element of $\ker \rho_a$ of length 54 is exhibited, uniformly across the family.

Proof. Take the two fixed words of length eight

$$w_1 = R_1 R_2 R_1 R_0 R_3 R_2 R_1 R_4, \quad w_2 = R_0 R_3 R_2 R_1 R_4 R_3 R_2 R_3,$$

and put $u = w_1 w_2^{-1}$, of length 16. These do not depend on a .

Computing $\pi(\rho_a(w_1)) - \pi(\rho_a(w_2))$ symbolically in $\mathbb{Q}(a_1, a_2, a_3, a_4)$, every one of its fourteen nonzero entries carries the factor

$$(a_1 a_3 - a_2 a_4)(a_1 a_3 + a_2 a_4).$$

The coordinates are positive, so the second factor never vanishes, and therefore $\pi(\rho_a(u)) = I$ if and only if $a_1 a_3 = a_2 a_4$. Both directions are confirmed directly: substituting $a_4 = a_1 a_3 / a_2$ makes the difference identically zero, while at $(1, 2, 5, 7)$, off the locus, it is nonzero.

So on the locus $\rho_a(u)$ is a translation, and $u \neq 1$ in W_4 by the Tits representation. Proposition 4.1 places $c = [u, R_0 u R_0]$ in $\ker \rho_a$, and $c \neq 1$ in W_4 , again by the Tits representation; reducing c gives a geodesic of length 54. Since w_1, w_2 and hence c are independent of a , the same element works for every member of the family. $\square$

Remark 5.6 (the family is genuinely infinite, and the criterion is sharp). The equation $a_1a_3 = a_2a_4$ has infinitely many pairwise-distinct positive integer solutions, for instance (k, km, kmn, kn) for distinct k, m, n scaled to be primitive, so Theorem 5.5 exhibits infinitely many Class C tuples at which ρ_a fails. This is a stronger statement than the dimension-three refutation, which produced only a finite list.

The criterion was found by scanning the 828 primitive pairwise-distinct tuples with entries at most 8, of which 24 fail by depth 12; the sixteen that fail at depth exactly 8 are precisely those with $a_1a_3 = a_2a_4$. It was then frozen and tested out of sample on ten tuples outside that box, all of which deviate at depth 8, against four controls with $a_1a_3 \neq a_2a_4$, none of which deviates by depth 12. The proof above supersedes that evidence: the colliding pair is a single pair of words, the same for every tuple, and the locus on which its linear parts agree is cut out exactly by $a_1a_3 = a_2a_4$. The remaining eight failures, at depth 12, form a separate family not covered by this criterion; their leg ratios are (1:2, 3:4) and (1:3, 4:3) up to reversal, and we do not characterise them.

Remark 5.7 (aligned valuations do not suffice). For an amalgam argument in dimension four one wants a tuple at which the splitting $W_4 = \langle R_0, R_2, R_4 \rangle *_{\langle R_0, R_4 \rangle} \langle R_0, R_1, R_3, R_4 \rangle$, which exists because $\{0, 4\}$ is a separating clique of Γ_4 , is matched by a p -adic valuation, so that one factor is integral and only the extra generator has a pole. At $(1, 2, 3, 4)$ this holds exactly, with $|m_1|^2 = 5$, $|m_2|^2 = 13$, $|m_3|^2 = 25$: at $p = 13$ every generator but R_2 is integral, and $v_{13}(R_2) = -1$. Nevertheless the point group at that tuple collapses at depth 12, and ρ_a is not injective there by Theorem 4.2. So aligning a valuation with the splitting is not sufficient for faithfulness, and an amalgam argument in dimension four needs an input that this alignment does not provide.

Remark 5.8 (the picture in dimension three). Theorems 4.2 and 5.2 are complementary and together settle the three-dimensional case. Faithfulness holds generically, and fails on a nonempty set of rational Class C tuples. So Class C is not the right condition: it neither implies faithfulness, by Theorem 4.2, nor is needed for it. The exceptional set is where the point group collapses, and by Corollary 4.4 that is exactly where ρ_a can fail.

Nothing here decides $n \geq 4$. The argument used that W_3 is virtually free, and that already fails at $n = 4$, though not for the reason one might expect. A triangle in Γ_n is no obstruction: it contributes a *finite* parabolic $(\mathbb{Z}/2)^3$, compatible with virtual freeness. The obstruction is an induced four-cycle. In Γ_4 the set $\{0, 3, 1, 4\}$ induces the cycle $0 - 3 - 1 - 4 - 0$, its non-edges being $\{0, 1\}$ and $\{3, 4\}$; a four-cycle is the join of two non-adjacent pairs, so the corresponding parabolic is $D_\infty \times D_\infty$, which contains $\mathbb{Z}^2$. Hence W_4 is not virtually free and Lemma 5.1 has no analogue there.

Remark 5.9 (what the reduction does not buy). It does not decide Problem 10.7 for any $n \geq 4$, the case $n = 3$ being settled affirmatively by Corollary 5.3, and by itself it does not touch Problem 10.5: an abstract embedding of W_n into $O(n)$ need not be by reflections, and the faithfulness of the particular map ρ_a is a separate question. The two standard invariants remain undecisive. Finite subgroups do not separate: every finite subgroup of W_n is conjugate into an elementary abelian 2-group on a clique of Γ_n , of rank at most $\lceil (n+1)/2 \rceil \leq n$, while $O(n)$ contains $(\mathbb{Z}/2)^n$. Free subgroups do not separate either, both groups containing free subgroups of rank two for $n \geq 3$.

6 The planar case

Let $F = C_2 * C_2 * C_2$ and $\rho_\tau: F \rightarrow G_\tau$ the generic triangle group.

Theorem 6.1 (Census; PROVED, computational input VERIFIED). *There are 33 pairs of reduced words of length 11, distinct in F , with $\rho_\tau(w) = \rho_\tau(w')$ for every τ . Outside a proper Zariski-closed*

subset of shape space these are the only coincidences in the ball of radius 11, and there are 132 further pairs at radius 12. The generic sphere sizes are

$$1, 3, 6, 12, 24, 48, 96, 192, 384, 768, 1536, 3039, 6012, 11925, 23570.$$

The 33 identities are polynomial identities in $\mathbb{Z}[p, q]$; the lower bounds come from counts in a finite field chosen to contain the needed roots of unity and $\sqrt{-1}$, and match the upper bounds from the identities, so the counts are exact.

Theorem 6.2 (Right-triangle structure; PROVED). *For $W = \langle R_0, R_1, R_2 \mid R_i^2 = (R_0 R_1)^2 = 1 \rangle$ the rotation subgroup is of index 4 and isomorphic to $\mathbb{Z} \wr \mathbb{Z}$, with translation lattice $\mathcal{T} = 2(t-1)\mathbb{Z}[t^{\pm 1}]$ obtained from Crowell's exact sequence and the Bass–Serre tree of the free product. The universal sequence is shape-independent through depth 32, and the shape $(1, 2)$ deviates at depth 33. If $c_T = N(a - ib) > 2d$ then no deviation occurs at depth d .*

Remark 6.3 (status of the metric). Word length equals relaxed length plus twice the connectivity defect, $\ell_T = \ell_R + 2c$. The upper bound is PROVED. The lower bound is open and is stated as a conjecture; the closed form c_{pred} computed by the explicit transducer agrees with the true defect with 0 exceptions on the 275,823 elements of the ball of radius 24, held out at depths 19, 21, 23, 24, which is HEURISTIC support and not a proof. The junction site cost is $\max(|d_L - 1|, |d_R|)$, which depends on the sign of d_L and not on its magnitude alone; the form $\max(|d_L| - 1, |d_R|)$ is false on every cell with $d_L \leq -2$ and $|d_R| \leq |d_L|$. This is a term inside the site-cost chain of the strand model, not an additive correction to the length formula.

7 The growth rate β_2

With $q = x^2$ and Σ_1 the odd k -recursion sum, $1 - \Sigma_1(q)$ is a Hahn–Exton q -cosine, and $q^* = 0.449453630558948046\dots$ is its smallest positive zero.

Theorem 7.1 (PROVED). *The growth rate of the planar right-triangle group is $\beta_2 = 1/\sqrt{q^*} = 1.4916177871\dots$, and $\beta_2 < 3/2$.*

The proof squeezes the true count between the pure-travel and relaxed counts, which share the exponential rate, and locates q^* by a contour argument with exact rational winding number. Irrationality of β_2 is open (Remark 1.4).

8 Transcendence: the unconditional part

Theorem 8.1 (PROVED). *Σ_1 and S_1 are transcendental over $\mathbb{Q}(q)$ and have $|q| = 1$ as a natural boundary.*

Theorem 8.2 (Travel resolvent; PROVED). *$\Sigma_0/(1 - \Sigma_1)$ is transcendental over $\mathbb{Q}(x)$, unconditionally.*

Transcendence of V is conditional on (M) and $(R - J)$; transcendence of U is conditional on (M) , $(R - J)$ and (T) . Neither is claimed here.

9 The q -cosine increment

Theorem 9.1 (PROVED). *For every algebraic $q \in (0, 1)$ the difference Galois group of $G(q, z) = {}_1\phi_1(0; q; q^2, z)$ over $\mathbb{C}(z)$ contains SL_2 , so G is hypertranscendental. The connection constant is $C_1 = 1/(q; q)_\infty$, whence $\prod_{k \geq 1} z_k(q) q^{2(k-1)} = (q; q^2)_\infty$.*

The general hypertranscendence statement, for all q not a root of unity and over $\bigcup_j \mathbb{C}(z^{1/j})$, is due to Adamczewski, Dreyfus and Hardouin and is stronger. The increment is the removal of a transcendence hypothesis on q in the SL_2 statement.

10 The five directions

Problem 10.1 (Prove $(R-J)$). Show $\Pi_y(q_m) \neq 0$ for infinitely many travel poles. This would make V , and with (T) also U , transcendental over $\mathbb{Q}(x)$ unconditionally, modulo (M) . What exists is an upper bound $|P_{12}|w^3 < 2$; what is needed is a lower bound $|P_{12}(q_m)| \geq c w_m^{-3}$.

Remark 10.2 (where the chain terminates). The measured laws for the two sides of $(S2)$ reduce, on unification, to the single scale $w = \sqrt{2/\tau}$ at a travel pole, together with the quantisation $w_m = (m - \frac{1}{2})\pi$. That quantisation is not an independent statement. Writing $S_e = \cos W - T_2$, the size of T_2 is $(\sqrt{2}/36)\sqrt{\tau} \sin X + O(\tau)$ uniformly in the phase, so the zeros of S_e are those of $\cos W$ displaced by $O(\sqrt{\tau})$, which is exactly $w_m = (m - \frac{1}{2})\pi + O(\sqrt{\tau})$.

So the chain reads: $(R-J)$ follows from $(S1)$ and $(S2)$; $(S2)$ is $|C_\lambda|/(xB) = w/2$; both sides are measured; and the scale w itself is fixed by the sharp size of T_2 . The last of these is a known HEURISTIC of the companion paper, with the obstruction already identified: the steepest-descent argument would conclude by invoking a standard remainder estimate for a simple non-degenerate saddle, but the saddle here is flat, and the bounded variation of the amplitude is available only on a bounded piece of the contour. The constant and the remainder exponent are certain to eight digits over $m \leq 1024$, and no theorem of that paper consumes them.

The chain therefore does not end in an unexamined gap. It ends at a named statement whose proof obstruction is diagnosed, and closing $(R-J)$ by this route requires the flat-saddle estimate rather than further measurement.

Remark 10.3 (two observations, and a route that does not work). First, the obstruction is a rank count, not a sign. The junction matrix $q^{\max(|a_L|-1, |a_R|)}$ has rank 4 on the tested block, and 6 after sign symmetrisation, so Π_y is a sum of several products rather than the single product that a rank-one kernel would give, and the terms can in principle cancel. At the same time the positivity does not degrade as quickly as one might expect from the interlacing of the zeros of S_e with those of $1 - \Sigma_1$: at the first six travel poles the dressed bulk values B_V and B_U are positive and increasing, 3.71, 493, 1979, $1.17 \cdot 10^4$, $2.31 \cdot 10^4$, $6.67 \cdot 10^4$ for B_V , and the gate signs $b_0 > 0$, $\mathfrak{g}_V t_1 < 1$ hold at every one.

Second, the sign of the travel null vector is a red herring, and this appears to be missed in the source. Since $L_s = R_s/e_s$, both factors of $\Pi_y = \langle \lambda, R \rangle \langle L, \mu \rangle$ are linear in R , so Π_y is a *quadratic form* in R . The vectors λ and μ are built from the bulk and do not involve R , so the overall sign and normalisation of R cancel in the product. That sign is precisely what flips when q_m crosses a zero of S_e : the values $\Sigma_0(q_m) = \sum_s R_s$ alternate, being 1.156, -4.545 , 7.752, -10.923 , 14.080, -17.232 , 20.381, -23.528 , 26.6, at the first nine poles, with $|\Sigma_0(q_m)|$ increasing by 3.389, 3.207, 3.171, 3.157, 3.152, 3.149, 3.147, 3.145, so $\Sigma_0(q_m) \sim (-1)^{m+1} \pi m$. The alternation is real, and it does not reach Π_y . This observation is independent of how λ and μ are normalised, since it uses only that neither depends on R .

Numerically $\Pi_y(q_m) > 0$ at each of the first nine travel poles, the two factors sharing the sign of R throughout. The travel null vector is computed from the forward recursion $R_s = e_s(q^s A_s + B_s)$ with $A_0 = 0$, $B_0 = 1$, whose consistency condition $B_\infty = 0$ is exactly $\Sigma_1(q) = 1$; the bulk solve is the gap-marked bulk system, whose gap term $q^{a+b} \mathbf{g}$ is rank one and so is handled by a Sherman–Morrison update. Both are validated against the published certificate: $\sum_s R_s$ reproduces $\Sigma_0(q_m)$ to ten digits, and B_V, B_U reproduce 3.7105027, 3.3241016, 492.95283, 233.43575, 1978.5952, 897.12122, 11663.55, 5110.5716 exactly. This is evidence, not proof: nine poles are not infinitely many, and the identification of the far-junction vector λ used here agrees with μ , which reproduces the proved positivity at the dominant pole but is not independently derived.

Third, the mechanism behind that positivity is a Sturm oscillation. At the m -th travel pole the null vector R has exactly $m-1$ sign changes, for $m = 1, \dots, 5$ observed to be 0, 1, 2, 3, 4 with the last change at $s = -, 3, 27, 75, 149$, so R is the m -th eigenvector of the travel operator and carries $m-1$ nodes. The portion beyond the last node dominates: the partial sums split as head $\approx -\Sigma_0(q_m)$ and tail $\approx 2\Sigma_0(q_m)$, so both weighted sums $\langle \lambda, R \rangle$ and $\langle R, \mu/e \rangle$ inherit the sign of the tail, which is the sign of $\Sigma_0(q_m)$. That is why the two factors share a sign and the product is positive. Making this a proof requires a quantitative tail-domination estimate uniform in m , which we do not have. A cheaper route is unavailable: the weight vectors are not positive, μ_s taking negative values at 2, 30, 33 and 130 of its entries for $m = 2, 3, 4, 5$, so no bare positivity argument applies.

Fourth, this reduces $(R-J)$ to two explicit inequalities. Using $\sum_s q^s R_s = \Sigma_1(q_m) = 1$, the source's own decomposition gives

$$\langle \lambda, R \rangle = xB + C_\lambda, \quad C_\lambda = \sum_\sigma \frac{B_\sigma}{2} (\Delta_\sigma^+ + \Delta_\sigma^-), \quad \Delta_\sigma^\pm = \sum_{2s+1 < \sigma \mp 1} (x^{\sigma \mp 1} - xq^s) R_s,$$

and we verified this identity numerically against the directly computed pairing at the first four poles. Every weight in C_λ is negative, since $2s+1 < \sigma \mp 1$ forces $x^{\sigma \mp 1} < x^{2s+1}$, so C_λ is a negatively weighted sum over the *head* of R , the part before its last node. Writing $\varepsilon_m = \text{sign } \Sigma_0(q_m)$, the observed picture is that the head of R carries sign $-\varepsilon_m$, hence C_λ carries sign ε_m , and $|C_\lambda| > xB$, so $\langle \lambda, R \rangle$ carries sign ε_m ; the same for $\langle L, \mu \rangle$, and the product is positive. Consequently $(R-J)$ follows from the two statements

(S1) the head of R carries sign $-\varepsilon_m$, for infinitely many m ;

(S2) $|C_\lambda| > xB$ and the corresponding inequality for μ , for infinitely many m .

Both are inequalities about explicitly given finite sums, and neither is proved here. This is where we leave the problem: not at “show the pairing is non-zero” but at (S1) and (S2).

Both hold at every pole computed, and with sharp laws rather than marginally. For (S1) the ratio head/ $(-\Sigma_0(q_m))$ takes the values 0.99634, 1.000475, 0.99999, 1.000044, 0.9999991, 1.000012, 1.000006, 1.000000 at $m = 2, \dots, 12$, so not merely the sign but the value is pinned: head $= -\Sigma_0(q_m)$ asymptotically, to seven digits by $m = 6$, equivalently the part of R beyond its last node sums to $2\Sigma_0(q_m)$. For (S2) the ratio $|C_\lambda|/(xB)$ takes the values 2.813, 3.352, 5.970, 6.533, 9.120, 9.687, 12.266, 12.835, 15.410, 15.980, 18.554 over the same range, increasing by alternately about 2.58 and 0.57, hence by about π every two steps, so $|C_\lambda|/(xB) \sim (\pi/2)m$ and the margin grows linearly in m .

At $m = 1$ both fail, head being empty and $|C_\lambda|/(xB) = 0.3995$, which is precisely the pole where positivity is available and $\Pi_y(q_1) > 0$ is already proved. The two regimes are therefore complementary: the dominant pole by Pringsheim positivity, and every computed pole above it by (S1) and (S2). What remains is to prove the two asymptotic laws, and it is worth saying exactly where that lands. The travel null vector is generated by $A_{s+1} - A_s = 2q(q^{2s} A_s + q^s B_s)$ and

$B_{s+1} - B_s = -2q(q^{3s}A_s + q^{2s}B_s)$ with $A_0 = 0$, $B_0 = 1$. Writing $\tau = -\log q$, $t = \tau s$ and $\varepsilon = \tau/2$, the continuum limit is the singularly perturbed system $\varepsilon Y' = M(t)Y$ with

$$M(t) = \begin{pmatrix} e^{-2t} & e^{-t} \\ -e^{-3t} & -e^{-2t} \end{pmatrix}, \quad \text{tr } M = 0, \quad \det M = 0, \quad M^2 = 0.$$

So M is nilpotent for every t , both eigenvalues vanishing identically. That is not an obstruction here; it is what makes the system solvable. Substituting $u = e^{-t}$ turns $\varepsilon Y' = MY$ into $\varepsilon Y' = -P(u)Y$ with $P = \begin{pmatrix} u & 1 \\ -u^2 & -u \end{pmatrix}$, so with $Y = (A, B)$ one has $\varepsilon A' = -(uA + B)$ and $\varepsilon B' = u(uA + B) = -u\varepsilon A'$. Eliminating B leaves

$$\varepsilon A'' + A = 0, \quad \varepsilon = \tau/2, \quad w := \varepsilon^{-1/2} = \sqrt{2/\tau}. \quad (2)$$

The boundary conditions are $A(1) = A_0 = 0$ and, from $B(0) = B_\infty = 0$, $A'(0) = 0$. Hence $A(u) = C \cos(wu)$, and $A(1) = 0$ forces the quantisation $\cos w = 0$, that is $w_m = (m - \frac{1}{2})\pi$, which is the recorded law $w_m \sim m\pi$; the relative error against $\sqrt{2/\tau_m}$ at the computed poles is $6.8 \cdot 10^{-3}$, $2.3 \cdot 10^{-3}$, $8.8 \cdot 10^{-4}$, $\dots$, $6.2 \cdot 10^{-5}$ for $m = 1, \dots, 10$. Since $A(0) = \Sigma_0(q_m)$ we get $C = \Sigma_0(q_m)$ and

$$A(u) = \Sigma_0(q_m) \cos(wu), \quad R_s \propto u \sin(wu), \quad u = q^s.$$

This settles (S1) at leading order. The nodes of R sit at $u = k\pi/w$ with $1 \leq k \leq m-1$, so R has exactly $m-1$ of them, which is the observed Sturm count at every computed pole; the last node in s is the smallest such u , namely π/w ; and therefore $\text{head} = A(\pi/w) = \Sigma_0(q_m) \cos \pi = -\Sigma_0(q_m)$, which is exactly the observed law. Against the exact null vector, A_s matches $\Sigma_0 \cos(wq^s)$ with maximal relative error 0.020, 0.011, 0.0069, $\dots$, 0.0024 for $m = 1, \dots, 8$, decaying like $1/m$.

The bulk does *not* linearise the same way, and we record the failed attempt. Its recursion carries a source, and the same elimination formally gives $\varepsilon A'' + A = -\mathfrak{g}\beta$, the oscillator with constant forcing. That reduction does not survive checking. Imposing $A(1) = 0$, $A'(0) = -1/\varepsilon$ and the self-consistency $\beta = B(1)$ yields $\mathfrak{g}\beta = (-1)^m w$, which at the second pole predicts +4.70 against an actual $\mathfrak{g}\beta = -81.75$; and testing $\varepsilon A'' + A$ directly on the discrete bulk gives 80.1, 115.2, 113.1 at $u = 0.3, 0.5, 0.7$ rather than a constant. So the bulk boundary analysis above is wrong somewhere, and (S2) is not derived. The likely culprit is that $\mathfrak{g}_V = q/(1-q)$ grows like $w^2/2$, so the forcing is not uniformly small and a boundary layer near $u = 1$ is not negligible; we have not resolved it.

What is therefore established is (S1) at leading order, with the closed form verified above, and not (S2). Uniform control of the discrete-to-continuum error is also missing.

There is, however, a reason to expect the continuum route to be the wrong one, and it is structural. Both chains are exactly solvable as discrete transfer products, and passing to a continuum destroys that. Writing the travel recursion as $(A_{s+1}, B_{s+1})^\top = M_s(A_s, B_s)^\top$ gives

$$M_s = \begin{pmatrix} 1 + 2q^{1+2s} & 2q^{1+s} \\ -2q^{1+3s} & 1 - 2q^{1+2s} \end{pmatrix}, \quad \det M_s = 1,$$

so the transfer lies in SL_2 . Moreover $M_s = I + c_s N_s$ with $c_s = 2q(q^2)^s$ and $N_s^2 = 0$, and $N_s = D_s^{-1} N D_s$ for the *fixed* nilpotent $N = \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}$ and $D_s = \text{diag}(q^{s/2}, q^{-s/2})$, while $D_s D_{s+1}^{-1} = \text{diag}(q^{-1/2}, q^{1/2})$ is constant. The ordered product therefore telescopes into an alternating product of a fixed unipotent family and a fixed diagonal, with purely geometric coefficients. The bulk chain has the same shape, $M_b = I + c_b N_b$ with $c_b = 2(q^2)^b$ and again $\det M_b = 1$, differing from the travel chain only in the prefactor $2q$ against 2 and in carrying a source.

Two consequences. First, geometric coefficients in SL_2 is precisely the q -hypergeometric setting in which the Hahn–Exton q -cosine lives, so the appearance of that function is structural rather than incidental. Second, because $\det M = 1$ the discrete Wronskian is conserved, so the inhomogeneous bulk chain is the homogeneous one plus a particular solution obtained by variation of parameters, with an exact discrete Green’s function and no continuum approximation anywhere. That is the form in which (S2) should be attacked.

The two chains are moreover the same chain. Writing $q = r^2$ and taking $a = q^s$ for the travel argument,

$$D M^{\text{travel}}(r^2, a) = M^{\text{bulk}}(ar) D, \quad D = \text{diag}(1, r),$$

an identity in the polynomial ring, with no hypotheses, no inverse and no fourth root. It is VERIFIED in `lean/mathlib/TransferChain.lean` as `half_shift`, together with $\det = 1$ for both chains and the nilpotence of their increments. An earlier form of this statement carried the hypotheses $r^2 = q$ and $rr^{-1} = 1$ and a conjugator involving $q^{1/4}$; none of that is needed, and the simplification was found in the course of formalising. This is the precise form of the statement that the zeros of S_e and of $1 - \Sigma_1$ are those of one q -cosine sampled a half q -step apart: the two chains differ by a half-step in the index and a constant conjugation, nothing else. In particular the homogeneous solutions of the bulk chain are those of the travel chain at shifted index, so the inhomogeneous bulk problem is closed by the conserved Wronskian alone.

Consistently with this, the bulk partial sums do fit the same oscillatory closed form. A least squares fit of A_b to $C_1 \cos(Wu) + C_2 \sin(Wu) + K$ with $u = q^b$ has relative residual $1.3 \cdot 10^{-7}$, $8.2 \cdot 10^{-4}$, $1.9 \cdot 10^{-3}$, $2.0 \cdot 10^{-3}$, $5.0 \cdot 10^{-4}$ at the first five poles, with $W/\sqrt{2/\tau} = 1.2, 1.05, 1.015, 1.01, 1.005$. The bulk is an oscillator with the same limiting frequency; the earlier failures came from imposing the travel frequency and the travel boundary data on it, not from any absence of the closed form.

One gap in the above should be recorded rather than passed over. Deriving the travel equation from $dA/ds = 2q^{1+s}(q^s A + B)$ and $du/ds = -\tau u$ gives $\varepsilon = \tau/2q$ and hence $w = \sqrt{2q/\tau}$, whereas the quantisation is matched by $\sqrt{2/\tau}$, and by a margin: at the tenth pole the two candidates miss $(m - \frac{1}{2})\pi$ by $1.2 \cdot 10^{-3}$ and $6.2 \cdot 10^{-5}$ respectively. A factor of q is unaccounted for, and until it is explained the closed form of (2) is supported by its verification and not by its derivation.

We record the exact structure, and not any consequence for (S2), which we have not derived.

Fifth, a Blaschke argument cannot be used, and this is worth recording so that it is not attempted. If Π_y vanished at all but finitely many q_m , one would like to conclude that the numerator vanishes identically. That needs the zeros to violate the Blaschke condition. They do not: writing $\tau_m = -\log q_m$, the products $m^2 \tau_m$ are 0.800, 0.362, 0.292, 0.265, 0.250, 0.241, 0.235, 0.231, 0.227, 0.225, 0.222, 0.220, for $m = 1, \dots, 14$, so $1 - q_m \asymp m^{-2}$ and $\sum_m (1 - q_m)$ converges, to about 0.72. A function of bounded characteristic on the disc may therefore vanish at every q_m without vanishing identically, and no contradiction follows.

Remark 10.4 (the arithmetic structure at $(1, 2, 3)$). For the generic form of Problem 10.5 a single faithful tuple suffices, and by Corollary 4.4 it is enough to find one at which the *point group* is faithful. The tuple $(1, 2, 3)$, where the point group agrees with the envelope through depth 22, carries an arithmetic structure that the others do not, and we record it because it is what any attempt should use.

The two denominators are $|m_1|^2 = a_1^2 + a_2^2 = 5$ and $|m_2|^2 = a_2^2 + a_3^2 = 13$, both prime. So R_0, R_2, R_3 lie in $GL_3(\mathbb{Z}_5)$ while $v_5(R_1) = -1$, and R_0, R_1, R_3 lie in $GL_3(\mathbb{Z}_{13})$ while $v_{13}(R_2) = -1$. The generators are orthogonal for the standard form, and $x^2 + y^2 + z^2$ is isotropic over $\mathbb{Q}_5$ and over $\mathbb{Q}_{13}$, so SO_3 is split at both places and its Bruhat–Tits building there is a *tree* rather than a rank-two building. Finally the graph splits: with $\Gamma_1 = \{0, 2, 3\}$ and $\Gamma_2 = \{0, 1, 3\}$ one has

$\Gamma_1 \cap \Gamma_2 = \{0, 3\}$, and 1 and 2 are non-adjacent in Γ_3 , so

$$W_3 = (\mathbb{Z}/2 \times D_\infty) *_{\mathbb{Z}/2 \times \mathbb{Z}/2} (D_\infty \times \mathbb{Z}/2),$$

the first factor $\langle R_0, R_2, R_3 \rangle$ being 5-integral and the second $\langle R_0, R_1, R_3 \rangle$ being 13-integral, with the amalgamated $\langle R_0, R_3 \rangle$ integral at both. Both factors are faithfully represented: each contains a D_∞ acting as rotations by an angle with rational cosine $3/5$ respectively $-5/13$, which has infinite order by Niven's theorem, and in each the extra involution reverses an axis that the D_∞ fixes. What is missing is the step from these data to injectivity of the amalgam.

The obstacle is that this is a ping-pong on *two* trees, one at each place, and a group acting on a product of trees is outside Bass–Serre theory. That can be removed. Solving $a_1^2 + a_2^2 = 5^j$ and $a_2^2 + a_3^2 = 5^k$ in distinct positive integers gives $(1, 2, 11)$, where $|m_1|^2 = 5$ and $|m_2|^2 = 125$, so the whole point group lies in $O_3(\mathbb{Z}[1/5])$ and acts on the *single* tree at $p = 5$, with R_0, R_3 integral and $v_5(R_1) = -1$, $v_5(R_2) = -3$. The point group there agrees with the envelope through depth 21 at both primes. So $(1, 2, 11)$, not $(1, 2, 3)$, is the tuple at which the argument should be attempted: one tree, and Bass–Serre theory applies to it. We have not carried the argument out, and record only that the arithmetic preconditions hold.

Problem 10.5 (The alternative in dimension $n \geq 4$). Is $N_{\text{gen}} = \bigcap_a \ker \rho_a$ trivial, equivalently is ρ_a injective for generic a ? A single faithful tuple answers this affirmatively. The arithmetic form is FALSE (Theorem 4.2), which does not decide the generic form.

Remark 10.6 (Class C faithfulness). The conjecture that ρ_a is injective for every $a \in \mathbb{Q}_{>0}^n$ with $e(a) = t(a) = 0$ and $n \geq 3$ is FALSE, by Theorem 4.2. Only the generic form (a) survives.

Problem 10.7 (Ambient embedding). For $n \geq 3$, does W_n embed in $O(n)$? By Proposition 3.2 this is equivalent to the ambient embedding question for $\text{Isom}(\mathbb{R}^n)$, and by Remark 3.5 it asks whether W_n has a faithful orthogonal representation of degree n .

Problem 10.8 (Finite presentation). For $n \geq 3$, is W_n/N_{gen} finitely presented? A negative answer refutes Problem 10.5(a). At $n = 2$ the answer is no.

Problem 10.9 (Transcendence of the zeros). Are the zeros $z_k(q)$ of the Hahn–Exton q -cosine transcendental at algebraic $q \in (0, 1)$, and is β_2 transcendental? This is the q -analogue of Siegel's theorem on the zeros of Bessel functions. Even irrationality of β_2 is open.