

Universality for orthoscheme reflection groups: the right-angled Coxeter envelope and the collision depth

Vico Bonfioli*

August 10, 2026

Abstract

The $n + 1$ facet reflections of an n -dimensional right-corner orthoscheme $O(a_1, \dots, a_n)$ generate a group $W(a) \leq \text{Isom}(\mathbb{R}^n)$, and $W(a)$ is a quotient of the right-angled Coxeter group W_n on the orthogonality graph of the facets. This paper determines what the quotient can do to the growth, and states precisely where that determination stops.

Four statements are proved unconditionally. First, a dimension-free dichotomy: if $Q = G/N$ is a quotient of a marked group and $K(N)$ is the length of a shortest nontrivial element of N , then the sphere sizes of Q and of G agree in every degree below $\lceil K(N)/2 \rceil$ and Q is strictly smaller in that degree (Theorem 5). Second, the growth series of W_n is rational, equal to $(1+t)^{\lfloor n/2 \rfloor + 1} / K_{n+1}(t)$ for an explicit integer polynomial K_{n+1} , with growth rate $r_n = 1 + 2 \cos(2\pi/(n+3))$ (Theorem 11). Third, for $n \geq 3$ and every positive leg tuple, every element of $\ker \rho_a$ of word length 6 is $(R_i R_{i+1})^{\pm 3}$ for a pair of consecutive facets meeting at angle $\pi/3$ (Theorem 9). Fourth, a complete classification of the rank-two relations: $\ker(W_n \rightarrow W(a))$ meets a rank-two standard parabolic nontrivially if and only if the leg sequence has two equal legs at an endpoint of the staircase or three consecutive equal legs (Theorem 19). The Diophantine input to the fourth is that three quartics in the flanking legs have only trivial rational points, their Jacobians being the rank-zero curves 24a1, 72a2 and $y^2 = x^3 - x$.

Together these determine the collision depth $\text{cd}_n(a) = \inf\{d : u_d(a) < [t^d] W_n(t)\}$, which equals $K(\ker \rho_a)/2$ and is at least 3 for every positive leg tuple, from the two positional statistics (e, t) of the leg sequence: $\text{cd}_n = 3$ when three consecutive legs are equal, $\text{cd}_n = 4$ when only an endpoint pair is equal, and on the remaining stratum $e = t = 0$, $\text{cd}_n = \infty$ exactly when ρ_a is injective and $\text{cd}_n \geq 4$ otherwise (Theorem 25). The first two cases are unconditional.

Whether ρ_a is injective at all is decided here in dimension two only, where it is not. That boundary is declared, not incidental. The planar proof is an amenability argument, and Theorem 30 shows that it cannot be run in any dimension $n \geq 3$: $O(n)$ contains a free subgroup of rank two for $n \geq 3$, so $\text{Isom}(\mathbb{R}^n)$ is not amenable as an abstract group, and any argument of that shape would have to prove the strictly stronger statement that W_n admits no embedding at all into $\text{Isom}(\mathbb{R}^n)$. Section 8 states the three problems that remain, with what is known about each. We also record that nonvanishing of the Schur-complement determinant $\det Q_n = -\prod_i a_i^2$ (Lemma 21) does not imply faithfulness, contrary to an argument used in an earlier version of this material (Remark 22).

1. Introduction

An n -dimensional right-corner orthoscheme, or path simplex, $O = O(a_1, \dots, a_n) \subset \mathbb{R}^n$ is the simplex with vertices

$$V_0 = 0, \quad V_k = V_{k-1} + a_k e_k \quad (1 \leq k \leq n),$$

*Independent researcher, Italy. vicobonfioli@gmail.com. <https://elvecio.github.io/home/posts/>.

where $e_1, \dots, e_n$ is the standard basis and $a = (a_1, \dots, a_n)$ is a tuple of positive reals, the *legs*. Its $n + 1$ facets carry reflections $\widehat{R}_0, \dots, \widehat{R}_n \in \text{Isom}(\mathbb{R}^n)$, and we write $W(a) = \langle \widehat{R}_0, \dots, \widehat{R}_n \rangle$ for the group they generate.

Two facets $\widehat{R}_i, \widehat{R}_j$ with $|i - j| \geq 2$ are perpendicular for every leg tuple (Lemma 1), while consecutive facets meet at an angle that depends on the legs and is generically not a rational multiple of π . The largest group compatible with the leg-independent relations is therefore the right-angled Coxeter group

$$W_n = \langle R_0, \dots, R_n \mid R_i^2 = 1, (R_i R_j)^2 = 1 \ (|i - j| \geq 2) \rangle, \quad (1)$$

on the *orthogonality graph* Γ_n (vertices $0, \dots, n$, an edge whenever $|i - j| \geq 2$), and there is a surjection $\rho_a: W_n \twoheadrightarrow W(a)$ for every a . Distinguishing W_n from $W(a)$ is essential: writing W_n for the geometric group would make “ ρ_a is faithful” vacuous, and it is exactly the possible failure of injectivity that this paper is about.

1.1 Results

The sphere sizes $u_d(a) = \#\{w \in W(a) : \ell(w) = d\}$ are bounded above by the coefficients of the growth series $W_n(t)$ of the envelope (1), and the whole question is where and by how much the bound fails to be sharp. Four unconditional theorems answer the “where”.

- **The dichotomy** (Theorem 5). For any group G with a finite symmetric generating set and any $N \trianglelefteq G$ with $N \neq \{1\}$, put $K(N) = \min\{\ell_G(g) : g \in N \setminus \{1\}\}$. Then the sphere sizes of G/N agree with those of G in every degree $d < \lceil K(N)/2 \rceil$, and are strictly smaller in degree $\lceil K(N)/2 \rceil$. So a family of quotients of a fixed G has a universal initial segment, and the horizon of universality is half the shortest added relator.
- **The envelope** (Theorem 11). $W_n(t)$ is rational, and is $(1+t)^{\lfloor n/2 \rfloor + 1} / K_{n+1}(t)$ for the explicit integer polynomials K_m of (2). Its growth rate is $r_n = 1 + 2 \cos(2\pi/(n+3))$.
- **The rank-two classification** (Theorem 19). For $a \in \mathbb{Q}_{>0}^n$, the kernel of ρ_a meets a rank-two standard parabolic $\langle R_i, R_{i+1} \rangle$ nontrivially if and only if $e(a) + t(a) \geq 1$, where $e(a)$ counts equal legs at the two endpoints of the staircase and $t(a)$ counts three consecutive equal legs. The shortest such element has length 6 when $t(a) \geq 1$ and length 8 when $t(a) = 0 < e(a)$.
- **Length six is dihedral** (Theorem 9). For $n \geq 3$ and every $a \in U$, an element of $\ker \rho_a$ of word length 6 is $(R_i R_{i+1})^{\pm 3}$ for a consecutive pair of facets meeting at angle $\pi/3$. There is no length-six relation outside the rank-two standard parabolics, in any dimension $n \geq 3$ and at any positive leg tuple.

The first two combine into the statement that gives the paper its shape. Define the *collision depth*

$$\text{cd}_n(a) = \inf\{d \geq 0 : u_d(a) < [t^d] W_n(t)\} \in \{1, 2, \dots, \infty\}.$$

By the dichotomy, $\text{cd}_n(a) = \lceil K(\ker \rho_a)/2 \rceil$, and since a relator of a reflection group has even length this is $K(\ker \rho_a)/2$ (Corollary 7). A short geometric argument then shows $K(\ker \rho_a) \geq 6$ for every positive leg tuple (Lemma 8), so $\text{cd}_n \geq 3$ unconditionally. The last two theorems pin the value: $K(a) = 6$ if and only if three consecutive legs are equal (Corollary 24), so $\text{cd}_n = 3$ exactly on that stratum, $\text{cd}_n = 4$ when only an endpoint pair is equal, and $\text{cd}_n \geq 4$ everywhere else (Theorem 25). The remaining case $e = t = 0$ is $\text{cd}_n = \infty$ precisely when ρ_a is injective.

1.2 Scope

The four theorems above are unconditional; they hold at every positive leg tuple apart from the arithmetic hypotheses of Theorem 19, and they are dimension-free except that Theorem 9

requires $n \geq 3$. The question they do not answer is whether ρ_a is injective at all when $e = t = 0$, and this paper answers that question in dimension two only, where the answer is no (Proposition 28). That is the declared boundary of the paper, and it is a boundary of the available method rather than of the write-up.

Theorem 30 makes that last sentence precise. The planar proof is an amenability argument: $\text{Isom}(\mathbb{R}^2)$ is virtually solvable, hence amenable, and so is every subgroup of it, whereas W_2 contains a free subgroup of rank two and is not. For $n \geq 3$ that argument is not merely unproved but unavailable, since $SO(3) \leq SO(n)$ contains a free subgroup of rank two and $\text{Isom}(\mathbb{R}^n)$ is therefore not amenable as an abstract group. Theorem 30 also identifies what any argument of that shape would have to establish, namely that W_n admits no embedding whatsoever into $\text{Isom}(\mathbb{R}^n)$, a statement strictly stronger than non-injectivity of the particular map ρ_a .

Section 8 states the three problems that remain: faithfulness on the $e = t = 0$ stratum (Conjecture 20, with the generic and arithmetic forms of Problem 32), the embedding question just described (Problem 33), and finite presentation of the generic n -orthoscheme group (Problem 34). Each is stated with what is known about it. Nothing outside those three is left implicit: Table 3 assigns a status to every numbered statement in the paper.

1.3 What is and is not new

The dichotomy is elementary. It is recorded here with a proof because it is what turns the two halves of this paper into statements about each other, and because its deviation half needs a case distinction on the parity of $K(N)$ that is easy to omit; but it is not a difficult argument, and the phenomenon it describes, that a family of quotients of a common group shares an initial segment of its growth, is a general one of which the orthoscheme family is an instance. The present paper claims the instance, not the principle.

Both ingredients of r_n are classical and are used as such. Steinberg's formula for the Poincaré series of a Coxeter group is standard [21, 13, 3], growth series of right-angled Coxeter groups are treated systematically by Kellerhals and Perren [15], and the independence polynomials of paths, their Binet formula, and the location of their roots are classical facts about the Fibonacci–Pell family. What is asserted is the identification: that the growth rate of the n -dimensional right-corner orthoscheme envelope is the largest of those roots at $m = n + 3$. We have not located that identification in the literature and make no claim of novelty for the ingredients. A reader who knows the Coxeter growth-series literature should read Theorem 11 as an assembly of known pieces.

The Diophantine part of Theorem 19 rests on three rank-zero Mordell–Weil computations that are in the standard tables [9, 16, 17]; what is done here is the reduction of the dihedral condition to those three curves.

Two points of contact should be recorded. The all-equal tuple $a_1 = \dots = a_n$ is not a generic point of this family: there the orthoscheme group is the affine Weyl group $\tilde{C}_n$, whose growth is classical and whose Poincaré series is given by Bott's formula, and which reproduces the $\text{cd}_n = 3$ rows of Table 2. In the planar case $n = 2$, Isola and Piergallini [14] prove that the generic triangle reflection group is not finitely presented, which already implies that ρ_a is not injective there; that is a second reason, besides the amenability argument of Proposition 28, why the plane is not a guide to $n \geq 3$.

1.4 Companion papers

The planar case $n = 2$ is treated separately [2]: there the metric has a lamplighter-type closed form, the deviation from the envelope has been computed past its onset, and the resulting growth series (A396406, growth rate $\beta_2 = 1.4916\dots$) is transcendental over $\mathbb{Q}(x)$. Nothing

below is quoted from that work except the two numerical facts recorded in Corollary 29, and the transcendence questions, which are special to $n = 2$, are not attempted here. Non-right-corner orthoschemes are outside the scope of this paper.

2. Orthoschemes and their Coxeter envelope

Throughout, $n \geq 2$, and

$$U = \{a = (a_1, \dots, a_n) \in \mathbb{R}^n : a_i > 0 \text{ for all } i\}$$

is the leg space. Everything below is invariant under simultaneous scaling of the legs, so one may normalise if convenient; we do not.

2.1 Facet normals

Lemma 1 (Facet normals). *With $V_0 = 0$ and $V_k = \sum_{i \leq k} a_i e_i$, the facet of $O(a)$ opposite V_j has normal*

$$m_0 = e_1, \quad m_j = a_{j+1}e_j - a_j e_{j+1} \quad (1 \leq j \leq n-1), \quad m_n = e_n.$$

Consequently $\text{supp}(m_0) = \{1\}$, $\text{supp}(m_j) = \{j, j+1\}$ and $\text{supp}(m_n) = \{n\}$, so $m_i \cdot m_j = 0$ whenever $|i-j| \geq 2$, while $m_i \cdot m_{i+1} \neq 0$ for every i and every $a \in U$.

Proof. The facet opposite V_j is the affine hull of $\{V_i : i \neq j\}$. For $j = 0$ the difference vectors $V_{k+1} - V_k = a_{k+1}e_{k+1}$, $1 \leq k \leq n-1$, span $\langle e_2, \dots, e_n \rangle$, whence $m_0 = e_1$. For $j = n$ the vertices $V_0, \dots, V_{n-1}$ span $\langle e_1, \dots, e_{n-1} \rangle$, whence $m_n = e_n$. For $1 \leq j \leq n-1$ the difference vectors are $a_k e_k$ for $k \leq j-1$, then $V_{j+1} - V_{j-1} = a_j e_j + a_{j+1} e_{j+1}$, then $a_k e_k$ for $k \geq j+2$. Orthogonality to the first and third families confines m_j to coordinates $\{j, j+1\}$, and orthogonality to the middle vector forces $m_j \propto a_{j+1}e_j - a_j e_{j+1}$. Two supports of the listed form are disjoint exactly when $|i-j| \geq 2$. For adjacent indices, $m_0 \cdot m_1 = a_2$, $m_j \cdot m_{j+1} = -a_j a_{j+2}$ for $1 \leq j \leq n-2$, and $m_{n-1} \cdot m_n = -a_{n-1}$, all nonzero because every $a_i > 0$. $\square$

Lemma 1 holds for all legs, with no genericity hypothesis and no case check. Hence $\widehat{R}_i^2 = 1$ always and $\widehat{R}_i \widehat{R}_j = \widehat{R}_j \widehat{R}_i$ whenever $|i-j| \geq 2$, so the assignment $R_i \mapsto \widehat{R}_i$ defines a surjection

$$\rho_a : W_n \twoheadrightarrow W(a) \leq \text{Isom}(\mathbb{R}^n)$$

from the right-angled Coxeter group (1) on the orthogonality graph Γ_n [10]. The abstract group W_n does not depend on a .

We write H_j for the facet hyperplane carrying $\widehat{R}_j$ and $\theta_{i,j} \in (0, \frac{\pi}{2}]$ for the angle between H_i and H_j , so that

$$\cos \theta_{i,j} = \frac{|m_i \cdot m_j|}{|m_i| |m_j|},$$

and $\widehat{R}_i \widehat{R}_j$ is a rotation through $2\theta_{i,j}$ about $H_i \cap H_j$.

The $n+1$ normals live in $\mathbb{R}^n$, so they carry one linear relation. It is unique and involves all of them, which is the source of every independence statement used below.

Lemma 2 (The unique relation among the normals). *Let $a \in U$. The vectors $m_0, \dots, m_n$ span $\mathbb{R}^n$, and the space of linear relations $\sum_j c_j m_j = 0$ is one-dimensional, spanned by the relation determined by*

$$c_0 = -a_2 c_1, \quad c_k = \frac{a_{k-1}}{a_{k+1}} c_{k-1} \quad (2 \leq k \leq n-1), \quad c_n = a_{n-1} c_{n-1},$$

in which every coefficient is nonzero. Consequently any at most n of the vectors $m_0, \dots, m_n$ are linearly independent.

Proof. The matrix with rows $m_1, \dots, m_n$ is upper triangular in the basis $e_1, \dots, e_n$: row j has entries a_{j+1} in column j and $-a_j$ in column $j+1$ for $1 \leq j \leq n-1$, and row n is e_n . Its determinant is $a_2 a_3 \cdots a_n \neq 0$, so $m_1, \dots, m_n$ is a basis and the relation space of the $n+1$ vectors is one-dimensional. Only m_0 and m_1 involve e_1 , so the e_1 -coefficient of $\sum_j c_j m_j$ is $c_0 + a_2 c_1$; for $2 \leq k \leq n-1$ only m_{k-1} and m_k involve e_k , giving $a_{k+1} c_k - a_{k-1} c_{k-1}$; and the e_n -coefficient is $c_n - a_{n-1} c_{n-1}$. Setting these to zero gives the displayed recursion, and all a_i are positive, so $c_1 \neq 0$ forces every $c_j \neq 0$. A dependent subset of size at most n would give a nonzero relation with at least one vanishing coefficient, contradicting uniqueness up to scale. $\square$

2.2 Counting conventions

For a group H with a marked finite generating set we write ℓ_H for the word length and $s_d(H) = \#\{h \in H : \ell_H(h) = d\}$ for the sphere sizes. We abbreviate

$$u_d(a) = s_d(W(a)), \quad W_n(t) = \sum_{d \geq 0} s_d(W_n) t^d,$$

the generators being $\hat{R}_0, \dots, \hat{R}_n$ and $R_0, \dots, R_n$ respectively. Since $O(a)$ is a fundamental domain for the action of $W(a)$, $u_d(a)$ is also the number of distinct chambers $g \cdot O(a)$ at word distance d , which is what the breadth-first search of §10 enumerates. Being the sphere sizes of a quotient of W_n , they satisfy $u_d(a) \leq [t^d] W_n(t)$ for every d .

Finally, set

$$K(a) = \min\{\ell_{W_n}(w) : w \in \ker \rho_a \setminus \{1\}\} \in \{1, 2, \dots\} \cup \{\infty\},$$

with $K(a) = \infty$ when ρ_a is injective.

2.3 Generic rigidity

Before computing anything we record that “the generic orthoscheme group” is a well-defined object: it does not depend on which generic shape one picks. Write $N_{\text{gen}} = \bigcap_{a \in U} \ker \rho_a \trianglelefteq W_n$.

Theorem 3 (Generic rigidity). *For every $n \geq 2$ there is a co-null subset $U_{\text{gen}} \subseteq U$, the complement of a countable union of proper real-algebraic subvarieties, such that $\ker \rho_a = N_{\text{gen}}$, hence $W(a) \cong W_n/N_{\text{gen}}$, for every $a \in U_{\text{gen}}$. Consequently $u_d(a)$ is constant on U_{gen} , equal to $s_d(W_n/N_{\text{gen}})$, and this generic value is the maximum of u_d over all of U . Moreover U_{gen} contains every a whose coordinates are algebraically independent over $\mathbb{Q}$; in particular it is dense.*

Proof. Each facet normal $m_j(a)$ is a fixed polynomial vector in a by Lemma 1, so the reflection $\hat{R}_j = I - 2m_j m_j^\top / (m_j \cdot m_j)$, together with its translation part, has entries in $\mathbb{Q}(a)$ with denominator the nowhere-vanishing polynomial $m_j \cdot m_j$. Hence for each $w \in W_n$, choosing any word representative, every entry of $\rho_a(w)$ lies in $\mathbb{Q}(a)$, independently of the representative because ρ_a is well defined. The locus $V_w = \{a \in U : \rho_a(w) = I\}$ is the simultaneous vanishing of finitely many rational functions cleared of their nonvanishing denominators, hence Zariski-closed in the irreducible variety U . If $w \notin N_{\text{gen}}$ then $\rho_a(w) \neq I$ for some a , so V_w is a proper closed subvariety, of measure zero. As W_n is countable, $B := \bigcup_{w \in W_n \setminus N_{\text{gen}}} V_w$ is a countable union of proper subvarieties, and $U_{\text{gen}} := U \setminus B$ is co-null and dense.

For $a \in U_{\text{gen}}$ one has $N_{\text{gen}} \subseteq \ker \rho_a$ by definition, and $\ker \rho_a \subseteq N_{\text{gen}}$ because any $w \in \ker \rho_a \setminus N_{\text{gen}}$ would place a in $V_w \subseteq B$. Isomorphic groups with marked generators have identical sphere sizes, giving the constant value; for arbitrary a , $\ker \rho_a \supseteq N_{\text{gen}}$ makes $W(a)$ a quotient of W_n/N_{gen} , so $u_d(a) \leq s_d(W_n/N_{\text{gen}})$. Finally each V_w is defined over $\mathbb{Q}$; if the coordinates of a are algebraically independent over $\mathbb{Q}$ then a lies on no proper $\mathbb{Q}$ -subvariety, so $a \notin B$. Such a exist since $\dim U = n \geq 2$. $\square$

Corollary 4 (One faithful shape suffices). *If ρ_a is injective for a single $a \in U$, then $N_{\text{gen}} = \{1\}$ and ρ_b is injective for every $b \in U_{\text{gen}}$. Conversely, if ρ_b fails to be injective for some $b \in U_{\text{gen}}$, then ρ_a fails to be injective for every $a \in U$.*

Proof. $N_{\text{gen}} \subseteq \ker \rho_a = \{1\}$, and $\ker \rho_b = N_{\text{gen}}$ for $b \in U_{\text{gen}}$ by Theorem 3. The converse is the contrapositive, using $N_{\text{gen}} \subseteq \ker \rho_a$ for every a . $\square$

Corollary 4 is what allows the generic question and the arithmetic question of §8 to be treated as two forms of one problem in one direction. It gives nothing in the other direction: a specific rational tuple need not lie in U_{gen} , and U_{gen} contains no tuple with algebraically dependent coordinates, hence no rational tuple at all.

3. The universality–deviation dichotomy

Theorem 5 (Universality–deviation dichotomy). *Let $G = \langle S \rangle$ be a group with finite symmetric generating set S , let $N \triangleleft G$ with $N \neq \{1\}$, write $Q = G/N$ and $K = K(N) = \min\{\ell_G(g) : g \in N \setminus \{1\}\}$, and put $h = \lceil K/2 \rceil$. Then $s_d(Q) = s_d(G)$ for every $d < h$, and $s_h(Q) < s_h(G)$.*

Proof. Write $\pi : G \rightarrow Q$.

Agreement below h . Let $d < h$. If $d \geq 1$ then $2d \leq 2h - 2 \leq K - 1 < K$, since $2h \leq K + 1$. Suppose $g, g' \in B_d(G)$ satisfy $\pi g = \pi g'$. Then $g(g')^{-1} \in N$ and $\ell_G(g(g')^{-1}) \leq 2d < K$, so $g = g'$; thus π is injective on $B_d(G)$. It is also length-preserving there: if $\ell_G(g) = d$ and $\ell_Q(\pi g) = e < d$, pick g' with $\ell_G(g') = e$ and $\pi g' = \pi g$; then $\ell_G(g(g')^{-1}) \leq d + e < 2d < K$ forces $g = g'$, contradicting $e < d$. Finally π maps the sphere of radius d in G onto the sphere of radius d in Q , because $\ell_Q(q) = \min\{\ell_G(g) : \pi g = q\}$ realises every q with $\ell_Q(q) = d$ by some g of length exactly d . Hence $s_d(Q) = s_d(G)$.

Strict drop at h . Choose $g_0 \in N \setminus \{1\}$ with $\ell_G(g_0) = K$ and split a geodesic word for it as $g_0 = w_1 w_2$ with $\ell_G(w_1) = h$ and $\ell_G(w_2) = K - h = \lfloor K/2 \rfloor$; both factors are geodesic as written. From $w_1 w_2 \in N$ we get $\pi(w_1) = \pi(w_2^{-1})$, and $w_1 \neq w_2^{-1}$ since otherwise $g_0 = 1$. Surjectivity of π from the radius- h sphere of G onto that of Q , established in the previous paragraph for $d < h$, holds verbatim for $d = h$.

If K is even then $h = K/2 = \lfloor K/2 \rfloor$, so w_1 and w_2^{-1} are two distinct elements of the sphere of radius h in G with the same image, and $s_h(Q) \leq s_h(G) - 1$.

If K is odd then $h = (K + 1)/2$ and $\ell_G(w_2^{-1}) = h - 1$, so $\ell_Q(\pi w_1) \leq h - 1 < h$ and the image of w_1 does not lie on the sphere of radius h in Q . That sphere is therefore covered by π applied to the radius- h sphere of G with w_1 removed, and again $s_h(Q) \leq s_h(G) - 1$. $\square$

The parity split at the end is not cosmetic. When K is odd the two halves of a shortest relator lie on spheres of different radii, so no two points of the upstairs sphere need collide, and the count drops for the other reason. Both cases are machine-checked (§10).

Applied to ρ_a , Theorem 5 gives an exact alternative with no genericity and no dimension hypothesis.

Corollary 6 (The envelope alternative). *Exactly one of the following holds.*

- (i) ρ_a is injective. Then $W(a) \cong W_n$ and $u_d(a) = [t^d] W_n(t)$ for every d ; in particular the growth series of $W(a)$ is the rational function of Theorem 11, with growth rate r_n .
- (ii) ρ_a is not injective. Then $K(a) < \infty$, $u_d(a) = [t^d] W_n(t)$ for every $d < \lceil K(a)/2 \rceil$, and $u_d(a) < [t^d] W_n(t)$ at $d = \lceil K(a)/2 \rceil$.

Proof. If $\ker \rho_a = \{1\}$ then ρ_a is an isomorphism of marked groups and (i) is immediate. Otherwise $K(a)$ is finite and Theorem 5 with $G = W_n$, $N = \ker \rho_a$ gives (ii). $\square$

Corollary 7 (The collision depth is half the shortest relator). *For every $a \in U$, $\text{cd}_n(a) = \lceil K(a)/2 \rceil$, and $K(a)$ is even, so $\text{cd}_n(a) = K(a)/2$.*

Proof. The first equality is Corollary 6: below $\lceil K(a)/2 \rceil$ the two sequences agree and at $\lceil K(a)/2 \rceil$ they differ. For parity, let $\varepsilon : W_n \rightarrow \{\pm 1\}$ be the sign character, $\varepsilon(R_i) = -1$, which is well defined because all defining relators of (1) have even length; then $\varepsilon(w) = (-1)^{\ell(w)}$. On the other side, $\det \circ \rho_a$ is a homomorphism $W_n \rightarrow \{\pm 1\}$ sending each R_i to -1 , so $\det \rho_a(w) = \varepsilon(w)$. If $w \in \ker \rho_a$ then $\det \rho_a(w) = 1$, hence $\ell(w)$ is even. $\square$

Lemma 8 (No relator of length four). *For every $n \geq 2$ and every $a \in U$, $K(a) \geq 6$. Consequently $\text{cd}_n(a) \geq 3$.*

Proof. By Corollary 7, $K(a)$ is even, so it suffices to exclude $K(a) \in \{2, 4\}$.

Write H_i for the facet hyperplane carrying $\hat{R}_i$. The H_i are the $n + 1$ facet hyperplanes of a nondegenerate simplex, so they are pairwise distinct and pairwise non-parallel, and $H_i \cap H_j$ is the affine hull of $\{V_m : m \neq i, j\}$, of dimension $n - 2$. Distinct index pairs give distinct intersections, because an $(n - 2)$ -flat contains exactly $n - 1$ of the $n + 1$ affinely independent vertices.

$K(a) = 2$ would give a geodesic word $R_i R_j$ with $i \neq j$ and $\hat{R}_i = \hat{R}_j$, impossible since $H_i \neq H_j$.

Suppose $K(a) = 4$, realised by a geodesic word $R_i R_j R_k R_l$ with $i \neq j$, $j \neq k$, $k \neq l$ and $\hat{R}_i \hat{R}_j \hat{R}_k \hat{R}_l = 1$, that is $\hat{R}_i \hat{R}_j = \hat{R}_l \hat{R}_k$. For $H_i \neq H_j$ non-parallel, the isometry $\hat{R}_i \hat{R}_j$ is a rotation about $H_i \cap H_j$ through twice the angle between the hyperplanes, an angle in $(0, 2\pi)$; its fixed-point set is therefore exactly $H_i \cap H_j$. The same applies to $\hat{R}_l \hat{R}_k$, so $H_i \cap H_j = H_k \cap H_l$ and hence $\{i, j\} = \{k, l\}$ by the previous paragraph. Since $j \neq k$ this forces $i = k$ and $j = l$, so the relation reads $\hat{R}_i \hat{R}_j = \hat{R}_j \hat{R}_i$, that is $H_i \perp H_j$. For $|i - j| \geq 2$ that relation already holds in W_n and the word $R_i R_j R_i R_j$ is not geodesic; for $|i - j| = 1$ perpendicularity contradicts $m_i \cdot m_{i+1} \neq 0$ in Lemma 1. Both branches are impossible, so $K(a) \neq 4$. $\square$

3.1 Length six is dihedral

The next value, $K(a) = 6$, is the first that occurs. The point of this subsection is that when it occurs it occurs for the obvious reason: a length-six kernel element is always a dihedral relator inside a rank-two standard parabolic, in every dimension $n \geq 3$ and at every positive leg tuple. Nothing arithmetic enters; the proof is the rank calculus of products of reflections together with the affine geometry of the one facet hyperplane that misses the origin.

We use the following standard fact, in the form given by Carter [5, Lemma 2]. Let V be a Euclidean space and $g = s_{\alpha_1} \cdots s_{\alpha_k}$ a product of reflections in the hyperplanes $\alpha_i^\perp$. Then $\text{im}(g - 1) \subseteq \langle \alpha_1, \dots, \alpha_k \rangle$ always, and if $\alpha_1, \dots, \alpha_k$ are linearly independent then $\text{rank}(g - 1) = k$, so that $\text{im}(g - 1)$ is the whole of $\langle \alpha_1, \dots, \alpha_k \rangle$ and $\text{Fix}(g) = \bigcap_i \alpha_i^\perp$. We also use twice that $\text{im}(g - 1) = \text{Fix}(g)^\perp = \text{Fix}(g^{-1})^\perp = \text{im}(g^{-1} - 1)$ for g orthogonal.

Theorem 9 (Length-six kernel elements are dihedral). *Let $n \geq 3$ and $a \in U$. Every $w \in \ker \rho_a$ with $\ell_{W_n}(w) = 6$ lies in a rank-two standard parabolic. Explicitly, $w = (R_i R_{i+1})^{\pm 3}$ for an index i with $\theta_{i, i+1} = \pi/3$.*

Proof. Write $L : \text{Isom}(\mathbb{R}^n) \rightarrow O(n)$ for the linear part and $s_j = L(\hat{R}_j)$ for the reflection in $m_j^\perp$. The vertex $V_0 = 0$ lies on every facet except the one opposite it, so $H_1, \dots, H_n$ contain the origin while $H_0 = \{x_1 = a_1\}$ does not; accordingly $\hat{R}_j(x) = s_j x$ for $1 \leq j \leq n$ and $\hat{R}_0(x) = s_0 x + 2a_1 e_1$.

Fix a reduced word $w = R_{i_1} \cdots R_{i_6}$ representing an element of $\ker \rho_a$.

(0) *Rotations, and the reduction to large support.* For a factorisation $w = xy$ the element $yx = x^{-1}wx$ lies in $\ker \rho_a$ and is nontrivial in W_n , since $yx = 1$ would give $y = x^{-1}$ and $w = 1$. Hence $\ell(yx) \geq K(a) \geq 6$ by Lemma 8, while $\ell(yx) \leq 6$; so every cyclic rotation of the word is again a reduced word of length 6 representing an element of $\ker \rho_a$, with the same support. We may therefore rotate freely. If $|\text{supp}(w)| \leq 2$ then w lies in a standard parabolic of rank at most two, which is the assertion; so assume $|\text{supp}(w)| \geq 3$ and derive a contradiction.

(1) *A repeated letter at distance two forces the dihedral case.* Suppose $i_k = i_{k+2}$ for some k , indices read cyclically. Rotate so that $k = 1$ and put $a' = i_1 = i_3$, $b = i_2$. Reducedness forces $|a' - b| = 1$, since $R_{a'}R_bR_{a'} = R_b$ when the two generators commute. Let $u = R_{a'}R_bR_{a'}$ and $v = R_{i_4}R_{i_5}R_{i_6}$. Then $L\rho_a(u)$ is the reflection in $s_{a'}(m_b)^\perp$, so $\text{rank}(L\rho_a(u) - 1) = 1$; since $L\rho_a(v) = L\rho_a(u)^{-1}$ is orthogonal, $\text{rank}(L\rho_a(v) - 1) = 1$ too. Were i_4, i_5, i_6 pairwise distinct, their normals would be independent by Lemma 2 (here $3 \leq n$ is used) and Carter's lemma would give rank 3. Hence $i_4 = i_6$; write $c = i_4$, $d = i_5$, so $|c - d| = 1$ as before.

Now $\rho_a(u)$ and $\rho_a(v)$ are the reflections in the affine hyperplanes $\widehat{R}_{a'}(H_b)$ and $\widehat{R}_c(H_d)$, and $\rho_a(u)\rho_a(v) = 1$ with both factors involutions forces the two hyperplanes to coincide; call the common hyperplane P . As $\widehat{R}_{a'}$ fixes $H_{a'}$ pointwise, $P \supseteq H_{a'} \cap H_b$, and likewise $P \supseteq H_c \cap H_d$. Using $H_i \cap H_j = \text{aff}\{V_m : m \neq i, j\}$ as in Lemma 8, P contains every vertex V_m with $m \notin \{a', b\} \cap \{c, d\}$. If that intersection is empty, P contains all $n + 1$ vertices, which affinely span $\mathbb{R}^n$ and lie in no hyperplane. If it is a single index j , then P contains the n affinely independent vertices V_m , $m \neq j$, so $P = H_j$; the case $j = a'$ gives $H_b = \widehat{R}_{a'}(H_{a'}) = H_{a'}$, excluded, and the case $j = b$ gives $\widehat{R}_{a'}(H_b) = H_b$, which makes m_b an eigenvector of $s_{a'}$ and hence, the normals being pairwise non-proportional by Lemma 2, forces $m_{a'} \perp m_b$, contradicting $|a' - b| = 1$ and Lemma 1. Therefore $\{a', b\} = \{c, d\}$. The ordered pairs differ, since $(c, d) = (a', b)$ would make $w = (R_{a'}R_bR_{a'})^2 = 1$, so $(c, d) = (b, a')$ and $w = (R_{a'}R_b)^3$, of support $\{a', b\}$, contradicting $|\text{supp}(w)| \geq 3$.

So every three cyclically consecutive letters of w are pairwise distinct.

(2) *Dimension at least four.* Put $u = R_{i_1}R_{i_2}R_{i_3}$ and $v = R_{i_4}R_{i_5}R_{i_6}$. By (1) each triple has distinct entries, so by Lemma 2 and Carter's lemma $\text{im}(L\rho_a(u) - 1) = \langle m_{i_1}, m_{i_2}, m_{i_3} \rangle$ and $\text{im}(L\rho_a(v) - 1) = \langle m_{i_4}, m_{i_5}, m_{i_6} \rangle$; and these two images agree because $L\rho_a(v) = L\rho_a(u)^{-1}$. For $n \geq 4$ two different three-element subsets of $\{m_0, \dots, m_n\}$ have different spans, since an index in one and not the other would produce a linear dependence among four normals, excluded by Lemma 2. Hence $\{i_1, i_2, i_3\} = \{i_4, i_5, i_6\}$. Combined with (1): $i_4 \notin \{i_2, i_3\}$ gives $i_4 = i_1$, then $i_5 \notin \{i_3, i_4\}$ gives $i_5 = i_2$, and $i_6 = i_3$. Thus $w = (R_pR_qR_r)^2$ with p, q, r distinct.

Let $E = \langle m_p, m_q, m_r \rangle$, of dimension 3, and $g = s_p s_q s_r$. Each of the three reflections is the identity on $E^\perp$ and preserves E , and $g^2 = 1$ because $\rho_a(w) = 1$. Carter's lemma gives $\text{rank}(g - 1) = 3$, so $g - 1$ vanishes on $E^\perp$ and is invertible on E ; an involution of E with no nonzero fixed vector is $-\text{id}_E$, so $s_p s_q s_r = -\text{id}$ on E , that is $s_q s_r = -s_p$ on E . On E the right-hand side is the rotation by π about the line $\mathbb{R}m_p$, and the left-hand side is the rotation about $\langle m_q, m_r \rangle^\perp \cap E$ through twice the angle between m_q and m_r . Equality of the axes gives $m_p \perp m_q$ and $m_p \perp m_r$, equality of the angles gives $m_q \perp m_r$. By Lemma 1 the indices p, q, r are then pairwise at distance at least 2, so R_p, R_q, R_r commute pairwise in W_n and $(R_pR_qR_r)^2 = 1$ there, contradicting $\ell(w) = 6$.

(3) *Dimension three.* Here every three of the four normals span $\mathbb{R}^3$, so the span argument of (2) is unavailable and the affine part is used instead. Let k be the number of occurrences of R_0 in w . By (1) two occurrences of one letter are at cyclic distance at least 3, and a cycle of length 6 admits at most two positions pairwise at cyclic distance at least 3; hence $k \leq 2$. The translation part of $\rho_a(w)$ is $2a_1 \sum_{j: i_j=0} P_{j-1}e_1$, where $P_m = s_{i_1} \cdots s_{i_m} \in O(3)$ and $P_0 = 1$, and it must vanish. Each summand is a unit vector and $a_1 > 0$, so $k \neq 1$.

If $k = 0$ then $\text{supp}(w) = \{1, 2, 3\}$, and by (1) six letters drawn from a three-element set with

every three cyclically consecutive ones distinct repeat with period three, so $w = (R_p R_q R_r)^2$ with $\{p, q, r\} = \{1, 2, 3\}$. The second paragraph of (2) applies verbatim with $E = \mathbb{R}^3$ and yields $m_1 \perp m_2$, contradicting Lemma 1.

If $k = 2$, rotate so that $i_1 = i_4 = 0$. Vanishing of the translation part reads $e_1 + s_0 s_{i_2} s_{i_3} e_1 = 0$; applying s_0 , which negates e_1 , gives $s_{i_2} s_{i_3} e_1 = e_1$. The distinct indices i_2, i_3 lie in $\{1, 2, 3\}$, so their normals are independent and Carter's lemma gives $\text{Fix}(s_{i_2} s_{i_3}) = m_{i_2}^\perp \cap m_{i_3}^\perp$. Thus $e_1 = m_0$ is orthogonal to m_{i_2} and m_{i_3} , so $\{i_2, i_3\} = \{2, 3\}$ by Lemma 1. But then R_0 commutes with both in W_3 , and

$$w = R_0 R_{i_2} R_{i_3} R_0 R_{i_5} R_{i_6} = R_{i_2} R_{i_3} R_0 R_0 R_{i_5} R_{i_6} = R_{i_2} R_{i_3} R_{i_5} R_{i_6}$$

has length at most 4, contradicting $\ell(w) = 6$.

Every branch is impossible, so $|\text{supp}(w)| = 2$ and w lies in a standard parabolic $\langle R_i, R_j \rangle$, $i \neq j$. If $|i - j| \geq 2$ that parabolic is $(\mathbb{Z}/2)^2$ and has no element of length 6, so $|i - j| = 1$ and the parabolic is infinite dihedral; its two elements of length 6 are $(R_i R_{i+1})^3$ and its inverse. Then $\rho_a(R_i R_{i+1})$ has order dividing 3, and is not the identity because $H_i \neq H_j$, so it has order exactly 3 and $\theta_{i,i+1} = \pi/3$. $\square$

Remark 10 (Where each hypothesis is used). Positivity of the legs enters twice: through Lemma 2, and in step (3) through $a_1 > 0$, which is what makes a single occurrence of R_0 leave a nonzero translation. The hypothesis $n \geq 3$ enters through the independence of any three normals, and it cannot be dropped: at $n = 2$ the three normals are dependent, and indeed the plane carries kernel elements of full support, though of length 20 rather than 6 (Corollary 29). No rationality is assumed anywhere in Theorem 9.

4. The envelope in closed form

Theorem 11 (The envelope is rational). *Let $c_{\Gamma_n}(x) = \sum_C \text{clique } x^{|C|}$ be the clique polynomial of Γ_n . The growth series of W_n is rational and satisfies*

$$\frac{1}{W_n(t)} = c_{\Gamma_n}\left(\frac{-t}{1+t}\right).$$

Equivalently $W_n(t) = (1+t)^{n+1}/J_{n+1}(t)$, where

$$J_m = (1+t)(J_{m-1} - tJ_{m-2}), \quad J_0 = J_1 = 1.$$

One has $J_m = (1+t)^{\lfloor m/2 \rfloor} K_m$ with

$$K_0 = K_1 = 1, \quad K_{2j} = K_{2j-1} - tK_{2j-2}, \quad K_{2j+1} = (1+t)K_{2j} - tK_{2j-1}, \quad (2)$$

so that

$$W_n(t) = \frac{(1+t)^{\lfloor n/2 \rfloor + 1}}{K_{n+1}(t)}.$$

The reciprocals of the roots of K_{n+1} in $(1/3, 1)$ are the numbers $1 + 2 \cos \frac{2k\pi}{n+3}$, and the growth rate is the largest of them,

$$r_n = 1 + 2 \cos \frac{2\pi}{n+3}.$$

Proof. W_n is a right-angled Coxeter group, so its finite standard parabolics are exactly the cliques of Γ_n , each a group $(\mathbb{Z}/2)^{|C|}$ with growth polynomial $(1+t)^{|C|}$ [10]. Steinberg's formula [21, 3, 13] gives

$$\frac{1}{W_n(t^{-1})} = \sum_{C \text{ clique}} \frac{(-1)^{|C|}}{(1+t)^{|C|}} = c_{\Gamma_n}\left(\frac{-1}{1+t}\right).$$

Substituting $t \mapsto t^{-1}$ and simplifying $-1/(1+t^{-1}) = -t/(1+t)$ yields the displayed identity for $1/W_n(t)$. The substitution must be carried inside the clique polynomial; retaining the argument $-1/(1+t)$ gives a different rational function, already at $n = 2$.

The cliques of Γ_n are the independent sets of its complement, the path P_{n+1} , so $c_{\Gamma_n} = I_{n+1}$, the independence polynomial of the path on $n+1$ vertices, which obeys $I_m = I_{m-1} + xI_{m-2}$, $I_0 = 1$, $I_1 = 1+x$, and has the coefficient form $I_m(x) = \sum_k \binom{m+1-k}{k} x^k$. Put $J_m(t) = (1+t)^m I_m(-t/(1+t))$. Multiplying the recurrence by $(1+t)^m$ gives $J_m = (1+t)J_{m-1} - t(1+t)J_{m-2}$, which is the stated recurrence, with $J_0 = 1$ and $J_1 = (1+t)(1-t/(1+t)) = 1$. By construction $1/W_n(t) = I_{n+1}(-t/(1+t)) = J_{n+1}(t)/(1+t)^{n+1}$. The factorisation $J_m = (1+t)^{\lfloor m/2 \rfloor} K_m$ and the two-step recurrences (2) follow by induction on m , splitting on the parity of m ; cancelling $(1+t)^{\lfloor (n+1)/2 \rfloor}$ against $(1+t)^{n+1}$ leaves the exponent $\lceil (n+1)/2 \rceil = \lfloor n/2 \rfloor + 1$.

For the roots, fix $t > 0$ and solve the linear recurrence. Its characteristic equation is $\lambda^2 - (1+t)\lambda + t(1+t) = 0$, of discriminant $(1+t)(1-3t)$.

If $0 < t < 1/3$ both roots are real with $0 < \lambda_- < \lambda_+ < 1$, their sum being $1+t \leq 4/3$ and their product $t(1+t) < 4/9$. Writing $J_m = \alpha\lambda_+^m + \beta\lambda_-^m$, the initial conditions give $\alpha = (1-\lambda_-)/(\lambda_+ - \lambda_-) > 0$ and $\beta = (\lambda_+ - 1)/(\lambda_+ - \lambda_-) < 0$ with $\alpha + \beta = 1$. Then $J_m = \lambda_-^m(\alpha(\lambda_+/\lambda_-)^m + \beta)$, and the bracket is increasing in m with value 1 at $m = 0$, so $J_m > 0$. At $t = 1/3$ the root is double and $J_m = (1+m/2)(2/3)^m > 0$. Hence J_m has no root in $(0, 1/3]$.

If $t > 1/3$ the roots are $\lambda_{\pm} = \varrho e^{\pm i\theta}$ with $\varrho = \sqrt{t(1+t)}$ and $\cos \theta = \sqrt{(1+t)/(4t)}$, so θ ranges over $(0, \pi/3)$ and is strictly increasing in t , because $(1+t)/(4t) = \frac{1}{4} + \frac{1}{4t}$ is strictly decreasing. Then $J_m = \varrho^m(\cos m\theta - c \sin m\theta)$, where $J_0 = 1$ is automatic and $J_1 = 1$ forces $c\varrho \sin \theta = (t-1)/2$, that is $c = (t-1)/\sqrt{(1+t)(3t-1)}$. Since $\sin^2 \theta = (3t-1)/(4t)$, a direct computation gives $\cot 2\theta = (1-t)/\sqrt{(1+t)(3t-1)} = -c$. Therefore

$$J_m = 0 \iff \cot m\theta = c = -\cot 2\theta \iff \frac{\sin((m+2)\theta)}{\sin m\theta \sin 2\theta} = 0 \iff (m+2)\theta \in \pi\mathbb{Z}.$$

With $m = n+1$ the roots are exactly $\theta = k\pi/(n+3)$, $1 \leq k < (n+3)/3$. Inverting $\cos^2 \theta = (1+t)/(4t)$ gives $t = 1/(4\cos^2 \theta - 1) = 1/(1+2\cos 2\theta)$, so the corresponding t -values are $1/(1+2\cos \frac{2k\pi}{n+3})$. As θ increases with t , the smallest positive root is $k = 1$; it exceeds $1/3$ because $1+2\cos \frac{2\pi}{n+3} < 3$, and $1+t > 0$ there, so it is a root of K_{n+1} and not of the stripped factor. Hence $r_n = 1+2\cos \frac{2\pi}{n+3}$. $\square$

Remark 12 (Not a path spectral radius). The number $2\cos \frac{2\pi}{n+3}$ is not the spectral radius of the path P_{n+1} , which is $2\cos \frac{\pi}{n+2}$; the two disagree for every $n \geq 2$ (at $n = 2$ they are $0.618\dots$ and $1.414\dots$). The angles produced by the proof are $k\pi/(n+3)$, and the growth rate uses $\cos 2\theta$ at $\theta = \pi/(n+3)$, not $\cos \theta$.

Remark 13 (The clique polynomial is a Fibonacci–Pell polynomial). The coefficient form $I_{n+1}(x) = \sum_k \binom{n+2-k}{k} x^k$ used above is the statement that the clique count of Γ_n in size k is $\binom{n+2-k}{k}$, and it identifies c_{Γ_n} with the Fibonacci–Pell polynomial P_{n+3} , where $P_0 = 0$, $P_1 = 1$ and $P_m = P_{m-1} + yP_{m-2}$: the two families satisfy the same three-term recursion, by Pascal’s identity $\binom{n-k+2}{k} = \binom{n-k+1}{k} + \binom{n-k+1}{k-1}$, and agree at the bottom. This gives a second reading of the roots. The characteristic equation $x^2 = x + y$ has roots $\alpha, \beta = (1 \pm \sqrt{1+4y})/2$, the Binet formula yields $P_m(y) = (\alpha^m - \beta^m)/(\alpha - \beta)$, so $P_m(y) = 0$ iff $(\alpha/\beta)^m = 1$; setting $\alpha/\beta = e^{2\pi i j/m}$ gives $y_j = -1/(4\cos^2(\pi j/m))$, and translating back through $y = -1/(1+t)$ with $4\cos^2 \vartheta - 1 = 1+2\cos 2\vartheta$ returns the same values $1+2\cos \frac{2\pi j}{n+3}$. The Pascal step, the Binet formula and the translation identity are machine-checked (§10); the induction that assembles the Pascal step into the identification $c_{\Gamma_n} = P_{n+3}$ is a paper proof.

Corollary 14 (Asymptotic regime). $r_n \rightarrow 3$ as $n \rightarrow \infty$, with

$$3 - r_n = 4 \sin^2\left(\frac{\pi}{n+3}\right) \sim \frac{4\pi^2}{(n+3)^2}.$$

Since $2\cos(2\pi/m)$ has degree $\phi(m)/2$ over $\mathbb{Q}$, where ϕ denotes Euler's totient, the rate r_n is rational or a quadratic surd exactly when $\phi(n+3) \leq 4$, that is $n+3 \in \{5, 6, 8, 10, 12\}$; see Table 1, where φ denotes the golden ratio.

n	$W_n(t)$	r_n	algebraic form
2	$(1+t)^2/(1-t-t^2)$	1.6180...	$\varphi = (1+\sqrt{5})/2$ (golden)
3	$(1+t)^2/(1-2t)$	2.0000	exactly 2
4	$(1+t)^3/(1-2t-t^2+t^3)$	2.2470...	$1+2\cos(2\pi/7)$, cubic
5	$(1+t)^3/(1-3t+t^2+t^3)$	2.4142...	$1+\sqrt{2}$ (silver)
6	$(1+t)^4/(1-3t+3t^3)$	2.5321...	$1+2\cos(2\pi/9)$, cubic
7	$(1+t)^4/(1-4t+3t^2+2t^3-t^4)$	2.6180...	$\varphi^2 = (3+\sqrt{5})/2$
8	$(1+t)^5/(1-4t+2t^2+5t^3-2t^4-t^5)$	2.6825...	$1+2\cos(2\pi/11)$, degree 5
9	$(1+t)^5/(1-5t+6t^2+2t^3-4t^4)$	2.7321...	$1+\sqrt{3}$
10	$(1+t)^6/(1-5t+5t^2+6t^3-7t^4-2t^5+t^6)$	2.7709...	$1+2\cos(2\pi/13)$, degree 6

Table 1: The envelope $W_n(t) = (1+t)^{\lfloor n/2 \rfloor + 1} / K_{n+1}(t)$ and its growth rate $r_n = 1 + 2\cos(2\pi/(n+3))$. The recurrence (2), its agreement with the Steinberg expansion, the exactness of the division by $(1+t)^{\lfloor (n+1)/2 \rfloor}$, and the identification of the smallest positive root of K_{n+1} with $1/r_n$ are certified for $2 \leq n \leq 30$ by the program of §10.

Remark 15 (The orthoscheme sequences in the OEIS). For pairwise distinct legs the layer counts $u_d(a)$ agree with the envelope as far as they have been computed, and the resulting sequences are recorded in the On-Line Encyclopedia of Integer Sequences: A397439 ($n=3$), A397437 ($n=4$), A396927 ($n=5$) and A397438 ($n=6$), with the $n=7$ entry in preparation. Each of these is the coefficient sequence of the rational function $W_n(t)$ of Table 1, with dominant pole $1/r_n$. The planar entry A396406 is of a different nature: it records the deviated planar sequence, which agrees with $W_2(t)$ only up to degree 9 and whose generating function is transcendental over $\mathbb{Q}(x)$ [2]; see Corollary 29.

5. Rank-two relations

This section determines exactly when $\ker \rho_a$ meets a rank-two standard parabolic $\langle R_i, R_{i+1} \rangle$. Throughout it, the legs are positive rationals. By Lemma 1 the only pairs that can carry a nontrivial dihedral relation are the consecutive ones, and their cosines are

$$\cos \theta_{0,1} = \frac{a_2}{\sqrt{a_1^2 + a_2^2}}, \quad \cos \theta_{n-1,n} = \frac{a_{n-1}}{\sqrt{a_{n-1}^2 + a_n^2}}, \quad \cos \theta_{i,i+1} = \frac{a_i a_{i+2}}{\sqrt{(a_i^2 + a_{i+1}^2)(a_{i+1}^2 + a_{i+2}^2)}} \quad (3)$$

for $1 \leq i \leq n-2$. In each case $\cos^2 \theta \in \mathbb{Q}$. By the extended Niven theorem, a consequence of Conway–Jones [7] (if $\theta \in \mathbb{Q}\pi$ and $\cos^2 \theta \in \mathbb{Q}$ then $\cos \theta \in \{0, \pm \frac{1}{2}, \pm \frac{\sqrt{2}}{2}, \pm \frac{\sqrt{3}}{2}, \pm 1\}$), a relation $(R_i R_{i+1})^m = 1$ in $W(a)$ forces

$$\cos^2 \theta_{i,i+1} \in \{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}.$$

5.1 The two positional statistics

For $a \in \mathbb{Q}_{>0}^n$ set

$$e(a) = \#\{i \in \{1, n-1\} : a_i = a_{i+1}\} \in \{0, 1, 2\},$$

$$t(a) = \#\{i : 1 \leq i \leq n-2, a_i = a_{i+1} = a_{i+2}\}.$$

Thus e counts equal pairs at the two endpoint positions of the staircase and t counts interior triples of three consecutive equal legs. An equal pair sitting strictly inside the staircase, such as $a_2 = a_3$ in $(1, 2, 2, 3)$ at $n=4$, contributes to neither.

5.2 Boundary pairs

Lemma 16 (Boundary pairs). *A boundary pair carries an admissible value of $\cos^2 \theta$ if and only if the two legs involved are equal, in which case $\cos^2 \theta = \frac{1}{2}$ and $\theta = \pi/4$.*

Proof. Take the pair $(0, 1)$, so $\cos^2 \theta = a_2^2/(a_1^2 + a_2^2)$ by (3); the pair $(n-1, n)$ is symmetric. The value 0 forces $a_2 = 0$ and the value 1 forces $a_1 = 0$, both excluded by positivity. The value $\frac{1}{4}$ gives $4a_2^2 = a_1^2 + a_2^2$, that is $(a_1/a_2)^2 = 3$, and the value $\frac{3}{4}$ gives $(a_2/a_1)^2 = 3$; neither is possible over $\mathbb{Q}$. The value $\frac{1}{2}$ gives $a_1^2 = a_2^2$, that is $a_1 = a_2$, and conversely $a_1 = a_2$ gives $\cos \theta = 1/\sqrt{2}$. $\square$

All five cases of Lemma 16 are formalised in Lean 4 (§10).

5.3 Interior pairs: three Diophantine quartics

At an interior pair write $x = a_i$, $y = a_{i+1}$, $z = a_{i+2}$, so that by (3)

$$\cos^2 \theta_{i,i+1} = \frac{x^2 z^2}{(x^2 + y^2)(y^2 + z^2)}.$$

Requiring this to equal a constant c and setting $u = y^2$ gives the quadratic

$$c u^2 + c(x^2 + z^2)u + (c-1)x^2 z^2 = 0, \quad (4)$$

which has a rational root only if its discriminant $\Delta = c^2(x^2 + z^2)^2 - 4c(c-1)x^2 z^2$ is the square of a rational. Substituting the three interior values of c and clearing the constant factors gives, in the two flanking legs $x = a_i$ and $z = a_{i+2}$, the Diophantine *quartics*

$$\mathcal{V}_{1/4} : x^4 + 14x^2 z^2 + z^4 = k^2, \quad \mathcal{V}_{1/2} : x^4 + 6x^2 z^2 + z^4 = k^2, \quad \mathcal{V}_{3/4} : 9x^4 + 30x^2 z^2 + 9z^4 = k^2.$$

It matters that these are quartics in the legs and not conics in $X = x^2$, $Z = z^2$. Written in X and Z the three equations define conics with infinitely many rational points, among them $(X, Z) = (3, 2)$ on $X^2 + 6XZ + Z^2 = k^2$ with $k = 7$, and $(X, Z) = (2, 5)$ and $(5, 9)$ on the other two. Those points are irrelevant here precisely because X and Z must themselves be squares of legs.

Lemma 17. *Each of $\mathcal{V}_{1/4}$, $\mathcal{V}_{1/2}$, $\mathcal{V}_{3/4}$ has only trivial solutions over $\mathbb{Q}_{>0}^2$ in the legs (x, z) : every rational point has $xz = 0$ or $x = z$.*

Proof. $\mathcal{V}_{1/2}$. The quartic $x^4 + 6x^2 z^2 + z^4 = k^2$ is birationally equivalent to the elliptic curve $E_{1/2} : y^2 = x^3 - x$ (Mordell [19], §17). $E_{1/2}$ has Mordell–Weil rank 0 over $\mathbb{Q}$ by Fermat’s classical descent (Cassels [6], Ch. II), and its full rational-point set is the 2-torsion $\{\infty, (0, 0), (\pm 1, 0)\}$, pulling back to $xz = 0$ or $x = z$.

$\mathcal{V}_{1/4}$. The Jacobian elliptic curve is $E_{1/4} : y^2 = x(x-12)(x-16)$, which is Cremona 24a1 (conductor 24, $j = 35\,152/9$, Mordell–Weil rank 0 over $\mathbb{Q}$, torsion $\mathbb{Z}/4 \oplus \mathbb{Z}/2$); see [9] and the LMFDB record [16]. Each of its eight rational points pulls back to $xz = 0$ or $x = z$.

$\mathcal{V}_{3/4}$. The quartic admits a standard conic-to-Weierstrass substitution and reduces to $E_{3/4} : y^2 = x(x-300)(x-1200)$, Cremona 72a2 (conductor 72, the same j -invariant $35\,152/9$ as $E_{1/4}$, of which it is a quadratic twist, also rank 0, torsion $\mathbb{Z}/2 \oplus \mathbb{Z}/2$); see [9, 17]. Each torsion point again pulls back to a trivial quartic solution. $\square$

Lemma 17 says nothing about the diagonal $x = z$, which is where the interior analysis has to be completed separately.

Lemma 18 (Interior pairs with equal flanking legs). *Suppose $a_i = a_{i+2}$. Then the interior pair $(i, i+1)$ carries an admissible value of $\cos^2 \theta$ if and only if $a_i = a_{i+1} = a_{i+2}$, in which case $\cos^2 \theta = \frac{1}{4}$ and $\theta = \pi/3$.*

Proof. With $x = z$ the cosine is $\cos \theta = x^2/(x^2 + y^2)$, a positive rational. The values 0 and 1 of $\cos^2 \theta$ are excluded by positivity. The value $\frac{1}{2}$ requires $\cos \theta = 1/\sqrt{2}$ and the value $\frac{3}{4}$ requires $\cos \theta = \sqrt{3}/2$, neither rational. The value $\frac{1}{4}$ requires $\cos \theta = \frac{1}{2}$, that is $2x^2 = x^2 + y^2$, hence $y = x$; and conversely $x = y = z$ gives $\cos \theta = x^2/(2x^2) = \frac{1}{2}$. $\square$

5.4 The classification

Theorem 19 (Rank-two classification). *Let $n \geq 2$ and $a \in \mathbb{Q}_{>0}^n$. Then $\ker \rho_a$ meets some rank-two standard parabolic $\langle R_i, R_{i+1} \rangle$ nontrivially if and only if $e(a) + t(a) \geq 1$. More precisely:*

- (i) *if $t(a) \geq 1$, the interior pair at a triple has $\theta = \pi/3$ and $(R_i R_{i+1})^3 \in \ker \rho_a$ is an element of length 6;*
- (ii) *if $t(a) = 0$ and $e(a) \geq 1$, the corresponding boundary pair has $\theta = \pi/4$ and $(R_i R_{i+1})^4 \in \ker \rho_a$ is an element of length 8, and no shorter element of any rank-two standard parabolic lies in $\ker \rho_a$;*
- (iii) *if $e(a) = t(a) = 0$, no rank-two standard parabolic meets $\ker \rho_a$ nontrivially.*

Proof. Since $\langle R_i, R_{i+1} \rangle$ is an infinite dihedral standard parabolic of W_n , a word $(R_i R_{i+1})^m$ is geodesic of length $2m$, and $\ker \rho_a$ meets that parabolic nontrivially exactly when $\theta_{i,i+1} \in \pi\mathbb{Q}$, in which case the shortest element is $(R_i R_{i+1})^m$ with $\theta_{i,i+1} = \pi r/m$ in lowest terms. Non-consecutive pairs generate finite parabolics of W_n itself and contribute nothing.

Suppose $\theta_{i,i+1} \in \pi\mathbb{Q}$ for some i . By the extended Niven theorem $\cos^2 \theta_{i,i+1} \in \{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}$. If the pair is a boundary pair, Lemma 16 gives $a_i = a_{i+1}$ with the pair at an endpoint, that is $e(a) \geq 1$, and $\theta = \pi/4$. If the pair is interior with flanking legs $x = a_i \neq a_{i+2} = z$, then (4) has the rational root $u = a_{i+1}^2$, so its discriminant is a rational square and (x, z) is a nontrivial point of one of $\mathcal{V}_{1/4}, \mathcal{V}_{1/2}, \mathcal{V}_{3/4}$, contradicting Lemma 17. If the pair is interior with $a_i = a_{i+2}$, Lemma 18 gives $a_i = a_{i+1} = a_{i+2}$, that is $t(a) \geq 1$, and $\theta = \pi/3$. This proves (iii) and the “only if” direction.

Conversely, if $t(a) \geq 1$ then some interior pair has $\theta = \pi/3$ by Lemma 18, the dihedral image has order 6, and the shortest relator is $(R_i R_{i+1})^3$ of length 6: (i). If $t(a) = 0$ and $e(a) \geq 1$ then some boundary pair has $\theta = \pi/4$ by Lemma 16, the dihedral image has order 8, and the shortest relator in that parabolic is $(R_i R_{i+1})^4$ of length 8; by the previous paragraph no other consecutive pair contributes an element of $\ker \rho_a$, since a boundary pair would force $e \geq 1$ with $\theta = \pi/4$ again and an interior pair would force $t \geq 1$: (ii). $\square$

Following the terminology of the three-dimensional analysis, call a leg tuple *Class C* when its coordinates are pairwise distinct. Class C tuples have $e = t = 0$, so Theorem 19(iii) applies, but so do tuples with an interior equal pair such as $(1, 2, 2, 3)$ and tuples with a non-adjacent repetition such as $(1, 2, 1, 3)$; the latter is precisely the case that Lemma 18, and not Lemma 17, has to handle.

Conjecture 20 (Class C faithfulness). *For every $n \geq 3$ and every $a \in \mathbb{Q}_{>0}^n$ with pairwise distinct coordinates, ρ_a is injective; equivalently $u_d(a) = [t^d] W_n(t)$ for every $d \geq 0$.*

Theorem 19(iii) is the rank-two half of Conjecture 20 and is proved. What is missing is that a nontrivial kernel element must lie in a rank-two parabolic. Theorem 9 supplies exactly that in length 6, which is the shortest length available by Lemma 8; in unbounded length the statement is false in general, the planar case $n = 2$ being a counterexample. There the two consecutive

pairs are both boundary pairs, so for $a_1 \neq a_2$ Theorem 19(iii) applies and no rank-two parabolic meets $\ker \rho_a$; yet $\ker \rho_a \neq \{1\}$ by Proposition 28, and its shortest elements have length 20, arising as $w_1 w_2^{-1}$ from eight pairs of distinct geodesic words w_1, w_2 of length 10 using all three generators [2]. Conjecture 20 is verified computationally at $n = 3$ with legs $(1, 2, 3)$ through depth 10 and at $n = 4$ with legs $(1, 2, 3, 5)$ through depth 8 (§10).

5.5 A determinant identity, and what it does not give

The Lorentzian Gram matrix of the orthoscheme in staircase coordinates, with the standard $(n, 1)$ endpoint correction, is the tridiagonal $(n + 1) \times (n + 1)$ matrix Q_n with

$$\begin{aligned} Q_n[0, 0] &= 1 - a_1^2, & Q_n[i, i] &= a_i^2 + a_{i+1}^2 \quad (1 \leq i \leq n - 1), & Q_n[n, n] &= 1, \\ Q_n[0, 1] &= a_2, & Q_n[i, i + 1] &= a_i a_{i+2} \quad (1 \leq i \leq n - 2), & Q_n[n - 1, n] &= a_{n-1}. \end{aligned}$$

Lemma 21 (Uniform Schur-complement determinant identity). *For every $n \geq 3$ and every positive tuple $(a_1, \dots, a_n)$,*

$$\det Q_n = - \prod_{i=1}^n a_i^2.$$

Proof. Let $D_k = \det Q_n[:k+1, :k+1]$ be the leading principal minor of order $k + 1$. Tridiagonal expansion along the last row gives

$$D_k = Q_n[k, k] D_{k-1} - Q_n[k - 1, k]^2 D_{k-2} \quad (1 \leq k \leq n), \quad (\dagger)$$

with $D_{-1} = 1$. We claim

$$D_k = \left(\prod_{i=1}^k a_i^2 \right) \left(1 - \sum_{i=1}^{k+1} a_i^2 \right) \quad \text{for } 1 \leq k \leq n - 1. \quad (\star)$$

Base. $D_0 = 1 - a_1^2$ and $D_1 = (1 - a_1^2)(a_1^2 + a_2^2) - a_2^2 = a_1^2(1 - a_1^2 - a_2^2)$.

Interior step. For $2 \leq k \leq n - 1$ the recurrence $(\dagger)$ reads $D_k = (a_k^2 + a_{k+1}^2)D_{k-1} - (a_{k-1}a_{k+1})^2 D_{k-2}$. Introducing $S_{k-1} = \sum_{i=1}^{k-1} a_i^2$ and $P_{k-2} = \prod_{i=1}^{k-2} a_i^2$ and substituting $(\star)$ reduces the step to the polynomial identity

$$\begin{aligned} P_{k-2} a_{k-1}^2 a_k^2 (1 - S_{k-1} - a_k^2 - a_{k+1}^2) &= (a_k^2 + a_{k+1}^2) P_{k-2} a_{k-1}^2 (1 - S_{k-1} - a_k^2) \\ &\quad - (a_{k-1}a_{k+1})^2 P_{k-2} (1 - S_{k-1}) \end{aligned}$$

in the five independent symbols $(a_{k-1}, a_k, a_{k+1}, S_{k-1}, P_{k-2})$. It is independent of k and of n .

Final step. At $k = n$ the recurrence uses $Q_n[n, n] = 1$ and $Q_n[n - 1, n]^2 = a_{n-1}^2$, giving $D_n = D_{n-1} - a_{n-1}^2 D_{n-2}$. With $S_{n-1} = \sum_{i=1}^{n-1} a_i^2$ and $P_{n-2} = \prod_{i=1}^{n-2} a_i^2$,

$$D_{n-1} - a_{n-1}^2 D_{n-2} = P_{n-2} a_{n-1}^2 [(1 - S_{n-1} - a_n^2) - (1 - S_{n-1})] = -P_{n-2} a_{n-1}^2 a_n^2 = - \prod_{i=1}^n a_i^2.$$

The three identities close the induction for every $n \geq 3$, with no per- n verification. All of them, the induction that assembles them, and the boundary step are machine-checked in Lean 4 over an arbitrary commutative ring; no positivity, ordering or characteristic hypothesis is used (§10). $\square$

Remark 22 (Why nonvanishing of $\det Q_n$ does not give faithfulness). An earlier version of this material argued from Lemma 21 that $\det Q_n \neq 0$, hence the form is nondegenerate, hence ρ_a is

faithful by Tits' theorem. That argument is a non sequitur, in three independent ways, and the deduction is withdrawn.

First, it proves too much. The hypothesis of Lemma 21 is that the legs are positive and nothing more, so the conclusion $\det Q_n \neq 0$ holds at every positive tuple, including $(1, 1, \dots, 1)$. Were the inference valid, ρ_a would be faithful there too. It is not: breadth-first search at $n = 3$ with legs $(1, 1, 1)$ gives layer counts 1, 4, 9, 17, 28, 42 against the envelope 1, 4, 9, 18, 36, 72, a deficit already at depth 3, in agreement with Theorem 25, which assigns $\text{cd}_3 = 3$ to that tuple. The same computation at $n = 4$ with legs $(1, 1, 1, 1)$ gives 1, 5, 14, 31, 59, 101 against 1, 5, 14, 33, 75, 169.

Second, Tits' theorem concerns the canonical geometric representation of an abstract Coxeter system, on the space spanned by the simple roots with the form $B(\alpha_i, \alpha_j) = -\cos(\pi/m_{ij})$ [3]. It is not a statement about ρ_a , which is a different representation, on $\mathbb{R}^n$ rather than $\mathbb{R}^{n+1}$ and affine rather than linear. Nor is Q_n a Coxeter Gram matrix: its diagonal is not identically 1, and $Q_n[0, 0] = 1 - a_1^2$ is negative whenever $a_1 > 1$.

Third, run at $n = 2$ the argument returns a false answer. There $\det Q_2 \neq 0$ just as well, so it would give $u_d(a) = [t^d]W_2(t) = F_{d+3}$ for all d ; in fact the two agree only up to $d = 9$ and differ at $d = 10$, where the orbit count is 225 against the envelope's 233, a deficit of 8 accounted for by eight coincidences between distinct words of length 10 [2].

6. The collision depth

Definition 23. For $a \in \mathbb{Q}_{>0}^n$ the *collision depth* is

$$\text{cd}_n(a) = \inf\{d \geq 0 : u_d(a) < [t^d] W_n(t)\} \in \{1, 2, \dots\} \cup \{\infty\}.$$

By Corollary 7, $\text{cd}_n(a) = K(a)/2$, and by Lemma 8, $K(a) \geq 6$. Conjecture 20 says exactly that $\text{cd}_n = \infty$ for every Class C tuple. The value $K(a) = 6$ is decided by combining the two preceding sections.

Corollary 24 (When $K(a) = 6$). *Let $n \geq 3$ and $a \in \mathbb{Q}_{>0}^n$. Then $K(a) = 6$ if and only if $t(a) \geq 1$. In particular $K(a) \geq 8$, hence $\text{cd}_n(a) \geq 4$, at every tuple with $t(a) = 0$.*

Proof. If $t(a) \geq 1$ then Theorem 19(i) exhibits an element of $\ker \rho_a$ of length 6, and $K(a) \geq 6$ by Lemma 8, so $K(a) = 6$.

Conversely let $K(a) = 6$. By Theorem 9 the witnessing element is $(R_i R_{i+1})^{\pm 3}$ with $\theta_{i,i+1} = \pi/3$, so $\cos^2 \theta_{i,i+1} = \frac{1}{4}$. If $(i, i+1)$ is a boundary pair, Lemma 16 says the only value in the Niven list attained there is $\frac{1}{2}$, so this is impossible. If it is an interior pair with $a_i \neq a_{i+2}$, the proof of Theorem 19 places (a_i, a_{i+2}) on one of the three quartics of Lemma 17, which have no such point. If it is an interior pair with $a_i = a_{i+2}$, Lemma 18 gives $a_i = a_{i+1} = a_{i+2}$, that is $t(a) \geq 1$. The last statement follows since $K(a)$ is even and at least 6. $\square$

The next theorem describes cd_n for every leg multiplicity pattern. Although the multiplicity partition of $(a_1, \dots, a_n)$ has $p(n)$ possible shapes, the collision depth takes only three values, determined by the two local positional statistics (e, t) .

Theorem 25 (Collision depth, all $n \geq 3$). *Let $n \geq 3$ and $a \in \mathbb{Q}_{>0}^n$. Then $\text{cd}_n(a) \geq 3$, and*

$$\text{cd}_n(a) = \begin{cases} 3 & \text{if } t \geq 1, \\ 4 & \text{if } e \geq 1 \text{ and } t = 0, \\ \infty & \text{if } e = t = 0 \text{ and } \rho_a \text{ is injective.} \end{cases}$$

The first two cases are unconditional. In the third case, if ρ_a is not injective then $\text{cd}_n(a) \geq 4$.

Proof. $\text{cd}_n(a) = K(a)/2$ with $K(a)$ even and at least 6, by Corollary 7 and Lemma 8; this gives $\text{cd}_n(a) \geq 3$ in all cases.

If $t \geq 1$ then $K(a) = 6$ by Corollary 24, so $\text{cd}_n(a) = 3$.

If $e \geq 1$ and $t = 0$ then Theorem 19(ii) exhibits an element of $\ker \rho_a$ of length 8, so $K(a) \leq 8$, while Corollary 24 gives $K(a) \geq 8$. Hence $K(a) = 8$ and $\text{cd}_n(a) = 4$.

If $e = t = 0$ then $\text{cd}_n(a) = \infty$ if and only if $\ker \rho_a = \{1\}$, by Corollary 6; and if $\ker \rho_a \neq \{1\}$ then $K(a) \geq 8$ by Corollary 24, so $\text{cd}_n(a) \geq 4$. $\square$

Remark 26 (What is and is not decided by (e, t)). Theorem 25 determines cd_n outright on the two strata $t \geq 1$ and $t = 0 < e$, with no residual ambiguity in the value. On the third stratum $e = t = 0$ it reduces the question to a single one, whether ρ_a is injective, and gives $\text{cd}_n \geq 4$ if it is not. What Theorem 9 supplies is a lower bound on $K(a)$ off the triple stratum; it supplies no upper bound anywhere. The rank-two analysis of §5 controls the elements of $\ker \rho_a$ lying in a dihedral standard parabolic, Theorem 9 controls all elements of length 6, and beyond length 6 nothing in this paper controls the elements outside those parabolics.

Remark 27 (The run-length rule fails in $n \geq 4$). A natural but incorrect heuristic is to declare cd_n finite as soon as some consecutive equal pair appears, that is as soon as the run length $R = \max_i \#\{j : a_j = a_i\}$ is at least 2. At $n = 3$ this rule coincides with the (e, t) rule, because every consecutive equal pair in a three-term sequence is automatically at an endpoint. At $n \geq 4$ the two diverge: the orthoscheme $(1, 2, 2, 3)$ has an interior equal pair, so $R = 2$ but $(e, t) = (0, 0)$, and Theorem 25 places it in the first case; empirically its layer counts 1, 5, 14, 33, 75, 169, ... are identical to those of the Class C tuple $(1, 2, 3, 5)$ through every computed depth, falsifying the run-length rule.

n	leg sequence	(e, t)	cd_n	first six layer counts
3	(2, 3, 5)	(0, 0)	∞	1, 4, 9, 18, 36, 72
3	(1, 1, 2)	(1, 0)	4	1, 4, 9, 18, 35, 67
3	(1, 1, 1)	(2, 1)	3	1, 4, 9, 17, 28, 42
4	(1, 2, 3, 5)	(0, 0)	∞	1, 5, 14, 33, 75, 169
4	(1, 2, 2, 3)	(0, 0)	∞	1, 5, 14, 33, 75, 169
4	(1, 1, 2, 3)	(1, 0)	4	1, 5, 14, 33, 74, 163
4	(1, 1, 2, 2)	(2, 0)	4	1, 5, 14, 33, 73, 157
4	(1, 1, 1, 2)	(1, 1)	3	1, 5, 14, 32, 67, 135
5	(1, 2, 3, 5, 7)	(0, 0)	∞	1, 6, 20, 54, 136, 334
5	(1, 2, 2, 3, 5)	(0, 0)	∞	1, 6, 20, 54, 136, 334
5	(1, 1, 2, 3, 5)	(1, 0)	4	1, 6, 20, 54, 135, 327
5	(2, 1, 1, 1, 2)	(0, 1)	3	1, 6, 20, 53, 128, 297
6	(1, 1, 2, 3, 2, 2)	(2, 0)	4	1, 7, 27, 82, 224, 581

Table 2: A small atlas of representative orthoschemes in $n = 3, 4, 5, 6$ illustrating each attainable (e, t) class. The layer counts are exact-rational breadth-first search (§10); in every row the observed collision depth is the value predicted by Theorem 25, and the entries ∞ are observed agreement with the envelope as far as computed, not a proof. The two $n = 4$ rows with $(e, t) = (0, 0)$ have identical counts through every computed depth despite differing in run structure.

The rows with $\text{cd}_n = \infty$ saturate the upper bound $[t^d] W_n(t)$ as far as they are computed, as Conjecture 20 predicts; for $n = 3$ this reproduces $u_d = 9 \cdot 2^{d-2}$ for $d \geq 2$.

7. The plane, and why its argument does not extend

Proposition 28 (The plane). *For $n = 2$ and every $a \in U$, ρ_a is not injective, so alternative (ii) of Corollary 6 holds.*

Proof. $W(a) \leq \text{Isom}(\mathbb{R}^2) = \mathbb{R}^2 \rtimes O(2)$. The group $O(2)$ contains the abelian $SO(2)$ with index 2, so $\text{Isom}(\mathbb{R}^2)$ is virtually solvable as an abstract group, hence amenable, and so is every subgroup of it; in particular $W(a)$ is amenable. The right-angled Coxeter group W_2 is not amenable: its defining graph Γ_2 is not complete multipartite, since its complement P_3 is connected and is not a disjoint union of cliques, and a right-angled Coxeter group whose defining graph is not complete multipartite contains a free subgroup of rank 2 [10]. An amenable group is not isomorphic to a non-amenable one, so ρ_a is not injective. $\square$

Corollary 29 (The planar horizon). *For $n = 2$ and every shape admissible in the sense of [2], the collision depth is $\text{cd}_2 = 10$, equivalently $K(a) = 20$: $u_d(a) = F_{d+3} = [t^d]W_2(t)$ for $1 \leq d \leq 9$ (while $u_0 = 1$ and $F_3 = 2$), and at $d = 10$ the count is 225 against the envelope's 233. Past that horizon the sequence is A396406, with growth rate $\beta_2 = 1.4916\dots < \varphi = r_2$, and its generating function is transcendental over $\mathbb{Q}(x)$ [2]. This is the only dimension in which the deviation has been computed past its onset.*

The isosceles right triangle $a_1 = a_2$ is excluded from that statement, and must be: there $e(a) = 1$, so Theorem 19(ii) gives an element of $\ker \rho_a$ of length 8 and hence $\text{cd}_2 \leq 4$.

7.1 The scope of the ambient obstruction

Proposition 28 is an instance of one argument scheme: find a property of abstract groups that passes to subgroups, holds for the ambient group $\text{Isom}(\mathbb{R}^n)$, and fails for W_n . Call such a property an *ambient obstruction* in dimension n . The scheme decides the alternative of Corollary 6 in the plane. The next theorem determines exactly when it can decide anything at all, and shows that in dimension $n \geq 3$ the specific property used in the plane is not merely unproved but false of the ambient group.

Theorem 30 (Ambient obstructions exist only in the plane). *For $n \geq 2$ consider the statement*

$$(E_n) : \quad W_n \text{ is isomorphic to a subgroup of } \text{Isom}(\mathbb{R}^n).$$

- (i) *If (E_n) fails, then ρ_a is not injective for every $a \in U$.*
- (ii) *If (E_n) holds, then every property of abstract groups that passes to subgroups and holds for $\text{Isom}(\mathbb{R}^n)$ also holds for W_n . No ambient obstruction exists in dimension n , and no argument of that scheme can decide Problem 32 there.*
- (iii) *$\text{Isom}(\mathbb{R}^n)$ is amenable as an abstract group if and only if $n \leq 2$, while W_n contains a free subgroup of rank 2 for every $n \geq 2$. Hence amenability is an ambient obstruction in dimension 2 and in no other; for $n \geq 3$ both groups contain free subgroups of rank 2.*

Proof. (i) If ρ_a were injective then $W_n \cong W(a) \leq \text{Isom}(\mathbb{R}^n)$, which is (E_n) .

(ii) Immediate from the definition: a subgroup-closed property of $\text{Isom}(\mathbb{R}^n)$ holds for all its subgroups, and under (E_n) the group W_n is one of them.

(iii) Write $\text{Isom}(\mathbb{R}^n) = \mathbb{R}^n \rtimes O(n)$. For $n \leq 2$ the group $SO(n)$ is abelian of index 2 in $O(n)$, so $\mathbb{R}^n \rtimes SO(n)$ is solvable of index 2 in $\text{Isom}(\mathbb{R}^n)$, which is therefore virtually solvable and amenable. For $n \geq 3$ the block embedding $SO(3) \leq SO(n) \leq \text{Isom}(\mathbb{R}^n)$ and Hausdorff's free subgroup of rank 2 in $SO(3)$ [22, Thm. 2.1] make $\text{Isom}(\mathbb{R}^n)$ non-amenable. For W_n : the complement of Γ_n is the path P_{n+1} on $n+1 \geq 3$ vertices, which is connected and is not a clique, so Γ_n is not complete multipartite and W_n contains a free subgroup of rank 2 [10]. $\square$

Remark 31 (Reading of Theorem 30). Three consequences are worth separating.

First, the temptation to be resisted. One is tempted to argue that $\text{Isom}(\mathbb{R}^n)$ is amenable because $O(n)$ is compact. Compactness gives amenability of $O(n)$ as a *topological* group, with respect to Haar measure, and that property does not pass to abstract subgroups. As an abstract group $O(n)$ is non-amenable for every $n \geq 3$, by the Hausdorff free subgroup that is the standard ingredient of the Banach–Tarski paradox. At $n = 2$ the argument survives only because $O(2)$ is virtually abelian.

Second, the amenability route is not repairable by replacing amenability with a different invariant. By Theorem 30(ii) the whole scheme is available in dimension n exactly when (E_n) fails, so any successful instance of it proves $\neg(E_n)$, whatever invariant it is dressed in.

Third, the scheme is strictly stronger than what Problem 32 asks. By Theorem 30(i), $\neg(E_n)$ gives non-injectivity of ρ_a at *every* leg tuple simultaneously, whereas Problem 32(a) asks only about the generic one. So an affirmative answer to Problem 33 below would close the scheme without saying anything about ρ_a , and a negative answer would settle Problem 32 outright. Either way the question is not a detail of the write-up; it is the exact content of the barrier.

8. What is proved, and the three open problems

8.1 What is proved

Table 3 lists the status of every numbered statement. The following are proved unconditionally: the dichotomy (Theorem 5); generic rigidity (Theorem 3), which makes “the generic n -orthoscheme growth” a well-defined object realised by a co-null set of shapes; rationality of the envelope with rate r_n (Theorem 11); the exclusion of length-six kernel elements outside the rank-two standard parabolics, at every positive leg tuple in every dimension $n \geq 3$ (Theorem 9); and the rank-two classification (Theorem 19). Together the last two determine the collision depth from the positional statistics (e, t) on the two strata $t \geq 1$ and $t = 0 < e$, and give $\text{cd}_n \geq 3$ everywhere and $\text{cd}_n \geq 4$ wherever $t = 0$ (Theorem 25).

What is not determined is cd_n on the third stratum $e = t = 0$, and there the question is exactly whether ρ_a is injective. That is settled here in the plane only, by Proposition 28, where the answer is no and the horizon is $\text{cd}_2 = 10$. For $3 \leq n \leq 7$ the geometric growth has been computed to depth 13 to 16 and agrees with the envelope throughout that range. That agreement is consistent with both alternatives of Corollary 6: under (i) it would continue forever, and under (ii) it would end at a horizon beyond the computed range. Nothing here decides between them.

8.2 The three open problems

The three problems below are the whole of what this paper leaves open. Each is stated in full, with what is known about it. Only the first is stated as a conjecture, and the evidence on both sides of it is recorded below; for the other two we state a question and decline to guess a direction.

Problem 32 (The alternative in dimension $n \geq 3$). *Fix $n \geq 3$.*

- (a) *Generic form. Is $N_{\text{gen}} = \bigcap_{a \in U} \ker \rho_a$ trivial? Equivalently, by Theorem 3, is ρ_a injective for every a in the co-null set U_{gen} ; equivalently, by Corollary 6, is the growth series of the generic n -orthoscheme group the rational envelope $W_n(t)$ in every degree?*
- (b) *Arithmetic form. Is ρ_a injective for every Class C tuple $a \in \mathbb{Q}_{>0}^n$? This is Conjecture 20.*

What is known. The two forms are related in one direction only. By Corollary 4, a single faithful tuple, rational or not, answers (a) affirmatively; so (b) implies (a). The converse is not automatic, because U_{gen} contains no tuple with algebraically dependent coordinates, hence

no rational tuple: an affirmative answer to (a) says nothing directly about any given rational a . The rank-two half of (b) is proved, Theorem 19(iii). No soft upgrade of that half to the whole is available: faithfulness would follow if every nontrivial kernel element lay in a rank-two parabolic, and that implication is false, the plane being a counterexample, where for $a_1 \neq a_2$ no rank-two parabolic meets $\ker \rho_a$ while $\ker \rho_a$ is nontrivial with shortest elements of length 20 using all three generators [2]. The same counterexample is a warning about (b) itself: the $n = 2$ analogue of Conjecture 20 is false. The positive evidence is confined to $n \geq 3$ and is computational, the agreement to depths 13 to 16 recorded above, together with Theorem 9, which removes the shortest length at which a relation could have appeared. If the answer to (a) is affirmative for all $n \geq 3$, the plane is the unique dimension with non-rational orbit growth. If instead $\ker \rho_a$ is nontrivial in some dimension $n \geq 3$, the growth deviates below the envelope at $\text{cd}_n(a) = K(a)/2 \geq 4$, and the arithmetic of the deviated series past that horizon is open even in the cases where the horizon can be located.

Problem 33 (The ambient embedding). *Fix $n \geq 3$. Is W_n isomorphic to a subgroup of $\text{Isom}(\mathbb{R}^n)$; that is, does statement (E_n) of Theorem 30 hold?*

What is known. (E_2) is false, by amenability, and that is the entire content of Proposition 28. For $n \geq 3$ the amenability route to $\neg(E_n)$ is closed: $\text{Isom}(\mathbb{R}^n)$ contains a free subgroup of rank 2 and so does W_n (Theorem 30(iii)). The obvious finite-subgroup invariant does not separate them either: every finite subgroup of W_n is conjugate into one of the elementary abelian 2-groups on the cliques of Γ_n [10], hence has rank at most $\lceil (n+1)/2 \rceil \leq n$, while $O(n)$ contains $(\mathbb{Z}/2)^n$. By Theorem 30(ii) this single question governs the entire class of ambient obstructions: if (E_n) holds, no such obstruction exists in dimension n ; if it fails, Problem 32 is answered negatively at every leg tuple at once. We do not conjecture a direction. Note that a positive answer would not touch Problem 32: an abstract embedding of W_n need not be by reflections in the facets of an orthoscheme, or by reflections at all.

Problem 34 (Finite presentation). *Fix $n \geq 3$. Is the generic n -orthoscheme reflection group W_n/N_{gen} finitely presented?*

What is known. At $n = 2$ the answer is no: Isola and Piergallini [14] show that the generic triangle reflection group is not finitely presented, which gives non-injectivity in the plane independently of the amenability argument. Since W_n is finitely presented, an answer either way decides Problem 32(a): a negative answer refutes it, and hence also refutes Conjecture 20 by Corollary 4, while an affirmative answer is consistent with it and does not prove it. Problems 32 and 34 are therefore not independent, and the direction in which the plane answers Problem 34 is the direction that would refute Conjecture 20. We know of no argument in either direction for $n \geq 3$.

8.3 Further questions

Three further questions are open and are stated for the record.

- (Q1) *Other diagrams.* Does an analogue of Theorem 25 hold for other families of simplicial reflection chambers in $\mathbb{R}^n$, for instance Coxeter orthoschemes with bent paths or simplices from non-path-shaped diagrams? The analysis here uses the right-angled-with-one-nontrivial-bond structure of the path diagram throughout.
- (Q2) *Intermediate rates.* In every dimension $n \geq 3$, multiplicity patterns that are neither Class C nor saturated by triples produce growth rates strictly between two consecutive values of r_n . Whether these intermediate rates are algebraic over $\mathbb{Q}$ at all is open.
- (Q3) *Beyond (e, t) .* Theorem 25 shows that the collision depth is determined by the positional statistics (e, t) . Is there a finer positional invariant that determines the entire sequence $u_d(a)$, not only its first deviation?

Statement	Status	Note
Lemma 1 (facet normals)	proved (Lean)	OrthoschemeNormals
Theorem 3 (generic rigidity)	proved	
Corollary 4	proved	
Theorem 5 (dichotomy)	proved (Lean)	EnvelopeDichotomy
Corollaries 6, 7	proved	
Lemma 8 ($K \geq 6$)	proved	
Lemma 2 (unique relation)	proved	
Theorem 9 (length six is dihedral)	proved	all $n \geq 3$, all positive legs
Theorem 11 (envelope)	proved (Lean, partly)	EnvelopeDichotomy , NdimRate
Corollary 14	proved (Lean)	NdimRate
Lemma 16 (boundary pairs)	proved (Lean)	RankTwoExclusion
Lemma 17 (three quartics)	proved	rank-0 descents, not formalised
Lemma 18 (equal flanking legs)	proved	
Theorem 19 (rank-two classification)	proved	interior half uses Lemma 17
Lemma 21 ($\det Q_n$)	proved (Lean)	OrthoschemeDet
Corollary 24 ($K = 6 \iff t \geq 1$)	proved	
Theorem 25, case $t \geq 1$	proved	$cd_n = 3$
Theorem 25, case $e \geq 1, t = 0$	proved	$cd_n = 4$, unconditional
Theorem 25, case $e = t = 0$	conditional	equivalent to Conj. 20
Conjecture 20 (Class C)	conjecture	checked to depth 10 ($n = 3$), 8 ($n = 4$)
Proposition 28 (the plane)	proved	
Theorem 30 (ambient obstruction)	proved	
Problem 32	open in both directions	
Problem 33 (ambient embedding)	open in both directions	false at $n = 2$
Problem 34 (finite presentation)	open in both directions	answered “no” at $n = 2$ [14]

Table 3: Status of the numbered statements. “Lean” means that the indicated Lean 4 file, which builds against Mathlib with no `sorry`, covers the statement; “partly” means that it covers some steps of the proof and not others. The scope of each file is described in §10. The three rows at the bottom are the whole of what the paper leaves open.

9. Related work

Path Coxeter groups and growth-rate calculus. The closed form $r_n = 1 + 2 \cos(2\pi/(n+3))$ is the signature of a linear, path-shaped Coxeter group whose Poincaré-series denominator reduces to a tridiagonal Chebyshev-type characteristic polynomial. Cannon and Wagreich [4], Floyd and Plotnick [12] and Parry [20] developed the growth-rate calculus of such groups, and Kellerhals and Perren [15] treat right-angled Coxeter growth systematically. Theorem 11 fits into that framework with parameter $m = n + 3$.

A Salem-type family on the surface-dynamics side. The family $\{1 + 2 \cos(2\pi/m)\}_{m \geq 3}$ to which r_n belongs also arises in birational surface dynamics. Diller and Favre [11] introduced the dynamical-degree framework for bimeromorphic maps of surfaces; McMullen [18] showed that this family arises as dynamical degrees of automorphisms of blowups of $\mathbb{P}^2$ associated with the E_n T-shaped Coxeter graph, which is not the path graph occurring here; and Bedford and Kim [1] realised specific members through linear-fractional recurrences. The rate r_n lies in this family, but the right-corner orthoscheme reflection group is not known to be realised as a McMullen involution product on a rational surface, and the coincidence is at the level of the algebraic-number family only. A birational-geometry construction realising r_n as a dynamical degree on a $\mathbb{P}^2$ -blowup would be of independent interest.

10. Certification

Orbit counts. The layer counts quoted above, the envelope coefficients and the Diophantine searches are produced by `code/zeta_probe/tools/ortho_cd` (Rust, exact arithmetic in $\mathbb{F}_p$ at two primes). It builds the $n + 1$ facet reflections directly from the legs, enumerates the group by breadth-first search, and compares layer by layer against $(1 + t)^{n+1}/J_{n+1}(t)$. In particular it produces the counts of Remark 22 at $(1, 1, 1)$ and $(1, 1, 1, 1)$ and at $n = 2$, confirms the (e, t) prediction on every tuple of Table 2 including the non-adjacent repetition $(1, 2, 1, 3)$, and searches the three quartics of Lemma 17 for nontrivial solutions with legs up to 300, finding none, while exhibiting rational points of the corresponding conics in (X, Z) to show that the restriction to squares is not cosmetic.

Short kernel elements. Theorem 9 is a proof, not a computation, but it was first tested by exhaustive search and the search is worth recording, since it also tests Lemma 8 and the (e, t) rule of Corollary 24. The program `code/zeta_probe/tools/ortho_len6` (Rust, exact integer arithmetic in homogeneous coordinates, no floating point) enumerates the reduced words of length at most 3 in W_n , computes their images under ρ_a , and inspects every collision; a nontrivial element of $\ker \rho_a$ of length at most 6 exists exactly when a collision does, and its length and support are read off by the deletion condition for graph products. Every positive rational tuple scales to a primitive integer tuple and $\rho_a(w) = 1$ is scale invariant, so integer boxes cover the rational case. The boxes run are

$$n = 3, \text{ legs } \leq 200; \quad n = 4, \leq 40; \quad n = 5, \leq 14; \quad n = 6, \leq 8; \quad n = 7, \leq 5; \quad n = 8, \leq 4,$$

comprising 9 954 322 primitive tuples in total. In every one of them: no element of $\ker \rho_a$ has length below 6; no element of length 6 has support of size 3 or more; and an element of length 6 occurs if and only if $t(a) \geq 1$, in which case it is $(R_i R_{i+1})^{\pm 3}$ at a triple of equal consecutive legs. The per-dimension counts are in the tool's `README.md`.

The envelope. Theorem 11 is certified by `code/zeta_probe/tools/racg_envelope` (Rust, exact integer polynomial arithmetic), which for $2 \leq n \leq 30$ checks that the recurrence for J_m and the Steinberg expansion agree coefficient by coefficient, performs the division by $(1 + t)^{\lfloor (n+1)/2 \rfloor}$ exactly, and matches the smallest positive root of K_{n+1} against $1/(1 + 2 \cos \frac{2\pi}{n+3})$. Independently, for $2 \leq n \leq 6$ it enumerates W_n by breadth-first search in the canonical geometric representation, whose matrices are integral here because the bilinear form takes only the values 1, 0, -1 and which is faithful by Tits' theorem [3]; the sphere sizes agree with the series coefficients to depths 9 to 14. The same program records two checks on the shape of the answer: the variant substitution $-1/(1 + t)$ in place of $-t/(1 + t)$ gives a different rational function already at $n = 2$, and the polynomial $1 - 2t - 3t^2 + 4t^3 - t^4$ is not the $n = 7$ denominator, being the negated reciprocal of $1 - 4t + 3t^2 + 2t^3 - t^4$ and predicting 6 elements at depth 1 in a group with 8 generators.

Lean 4. Six files in `lean/with_mathlib/` build against Mathlib with no sorry.

Paper result	Lean declaration
<i>OrthoschemeNormals.lean</i> (7 theorems)	
Lemma 1, support pattern	supp_disjoint
Lemma 1, orthogonality on disjoint supports	mul_eq_zero_of_disjoint, sum_eq_zero_of_disjoint
Lemma 1, the three adjacent products	dot_zero_one, dot_interior, dot_last
Lemma 1, combined	orthogonality_pattern
<i>EnvelopeDichotomy.lean</i> (12 theorems)	
Thm. 5, horizon arithmetic	two_mul_lt_of_lt_horizon, split_lengths
Thm. 5, even case	card_image_lt_of_collision
Thm. 5, odd case	card_lt_of_escapes
Thm. 11, three-term recurrence	cos_three_term, sin_three_term, J_recurrence
Thm. 11, root condition	cot_add_cot_eq_zero_iff, J_eq_zero_iff
Thm. 11, recovering t from θ	t_of_theta, rate_of_theta
Remark 12	rate_ne_path_spectral_radius
<i>NdimRate.lean</i> (7 theorems)	
Remark 13, Pascal step	pascal_step (no axioms), coeff_pascal
Remark 13, Binet formula	sq_eq, pell_binet
Remark 13, root translation	root_translation
Corollary 14	rate_residual, three_sub_r
<i>RankTwoExclusion.lean</i> (10 theorems)	
Lemma 16, the five Niven values	niven_zero, niven_one, niven_quarter, niven_three_quarter, niven_half, boundary_no_relation
Lemma 16, 3 is not a rational square	sq_ne_three, not_isSquare_three_int
Boundary cosine nonzero, and $\cos^2 \neq \frac{1}{4}$ there	no_length_two, no_length_three_boundary
<i>OrthoschemeDet.lean</i> (3 theorems)	
Lemma 21, closed form ($\star$)	minor_closed_form
Lemma 21, boundary step	det_Q
Lemma 21, full statement	det_Q_of_recurrence
<i>ReflectionTriple.lean</i> (13 theorems)	
Thm. 9, reflection calculus	inner_self_ne_zero', refl_self, refl_of_inner_eq_zero, refl_add, refl_smul, refl_neg, refl_sub, refl_involutive, refl_injective, inner_refl_left
Thm. 9, cancellation in the ambient group	eq_of_sub_eq_sub, eq_zero_of_sub_eq_self
Thm. 9, step (2): $s_p s_q s_r = -\text{id}$ forces pairwise orthogonality	pairwise_orth_of_comp_eq_neg

NdimRate.lean, *OrthoschemeDet.lean* and *ReflectionTriple.lean* end with `#print axioms`, one line per theorem of those three files. Of the twenty-three declarations audited, nineteen report exactly the standard axioms `propext`, `Classical.choice`, `Quot.sound`, and four report strictly fewer, with `pascal_step` depending on no axioms at all.

What is not formalised. *RankTwoExclusion.lean* covers the boundary half of Theorem 19 only; the interior cases rest on the rank-0 Mordell descents of Lemma 17 and are not formalised. *EnvelopeDichotomy.lean* formalises the arithmetic of Theorem 5 and the trigonometry of the root identity, but no word metric, no Coxeter group and no clique polynomial: the group theory around those steps is on paper. The step identifying the growth rate with the largest root t_1 is a dominant-pole argument on a rational generating function, proved on paper; so is the induction assembling the Pascal step into $c_{\Gamma_n} = P_{n+3}$ in Remark 13. Everything downstream of those two inputs is machine-checked. Of Theorem 9, *ReflectionTriple.lean* formalises the reflection calculus and the rigidity step (2) in an arbitrary real inner product space, with the reflection defined from the inner product and no dimension hypothesis; the word combinatorics in W_n , Carter's rank lemma, and the affine argument of step (3) are paper proofs. Lemma 2, Lemma 8,

Lemma 18, Theorem 3 and Theorem 30 are paper proofs.

Acknowledgments

Computational assistance from large language model assistants was used at multiple stages of this work; all mathematical claims have been verified by the author, who is solely responsible for the content of this paper and for any errors.

References

- [1] E. Bedford, K. Kim, *Dynamics of rational surface automorphisms: linear fractional recurrences*, J. Geom. Anal. **19** (2009), no. 3, 553–583.
- [2] V. Bonfioli, *The universal right-triangle reflection sequence*, companion paper; code and formalization at github.com/ElVec1o/pythagorean-reflection-sequence.
- [3] N. Bourbaki, *Groupes et algèbres de Lie, Chapitres IV–VI*, Hermann, Paris, 1968. (Geometric representation of a Coxeter group and its faithfulness; Steinberg’s identity.)
- [4] J. W. Cannon, P. Wagreich, *Growth functions of surface groups*, Math. Ann. **293** (1992), no. 2, 239–257.
- [5] R. W. Carter, *Conjugacy classes in the Weyl group*, Compositio Math. **25** (1972), 1–59. (Lemma 2: a product of reflections has $\text{rank}(g - 1)$ equal to the number of factors exactly when the roots are independent.)
- [6] J. W. S. Cassels, *Lectures on Elliptic Curves*, London Mathematical Society Student Texts 24, Cambridge University Press, 1991.
- [7] J. H. Conway, A. J. Jones, *Trigonometric Diophantine equations (On vanishing sums of roots of unity)*, Acta Arith. **30** (1976), 229–240.
- [8] H. S. M. Coxeter, *Discrete groups generated by reflections*, Ann. of Math. (2) **35** (1934), 588–621.
- [9] J. E. Cremona, *Algorithms for Modular Elliptic Curves*, 2nd edition, Cambridge University Press, 1997.
- [10] M. W. Davis, *The Geometry and Topology of Coxeter Groups*, London Math. Soc. Monographs **32**, Princeton University Press, 2008. (Right-angled Coxeter groups; finite parabolics as cliques; the complete multipartite criterion.)
- [11] J. Diller, C. Favre, *Dynamics of bimeromorphic maps of surfaces*, Amer. J. Math. **123** (2001), no. 6, 1135–1169.
- [12] W. J. Floyd, S. P. Plotnick, *Growth functions on Fuchsian groups and the Euler characteristic*, Invent. Math. **88** (1987), no. 1, 1–29.
- [13] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics 29, Cambridge University Press, 1990.
- [14] S. Isola, R. Piergallini, *On the generic triangle group*, preprint, arXiv:1405.1881.
- [15] R. Kellerhals, G. Perren, *On the growth of cocompact hyperbolic Coxeter groups*, European J. Combin. **32** (2011), 1299–1316.
- [16] LMFDB Collaboration, elliptic curve 24a1 over $\mathbb{Q}$, <https://www.lmfdb.org/EllipticCurve/Q/24/a/1>.

- [17] LMFDB Collaboration, elliptic curve 72a2 over $\mathbb{Q}$, <https://www.lmfdb.org/EllipticCurve/Q/72/a/2>.
- [18] C. T. McMullen, *Dynamics on blowups of the projective plane*, Publ. Math. Inst. Hautes Études Sci. **105** (2007), 49–89.
- [19] L. J. Mordell, *Diophantine Equations*, Pure and Applied Mathematics 30, Academic Press, 1969.
- [20] W. Parry, *Growth series of some hyperbolic Coxeter groups*, Geom. Dedicata **32** (1989), 75–96.
- [21] R. Steinberg, *Endomorphisms of Linear Algebraic Groups*, Mem. Amer. Math. Soc. **80** (1968). (The Poincaré-series identity for Coxeter groups.)
- [22] S. Wagon, *The Banach–Tarski Paradox*, Cambridge University Press, 1985. (Free subgroups of $SO(3)$.)