

A dimension-independent collision-depth invariant for right-corner orthoscheme reflection groups

Closed-form growth rate $r_n = 1 + 2 \cos(2\pi/(n+3))$ and unconditional Class C faithfulness in every dimension

Vico Bonfioli*

August 1, 2026

Abstract

For every $n \geq 3$ the reflection group W_n generated by the $n+1$ face-reflections of an n -dimensional right-corner orthoscheme admits a positional combinatorial invariant $\text{cd}_n(a_1, \dots, a_n) \in \{\infty, 4, 3\}$, the *collision-depth*, determined entirely by two integer statistics (e, t) of the leg sequence: the number of endpoint-adjacent equal pairs and the number of consecutive equal triples (Theorem 9). The same group has Coxeter–Steinberg growth rate $r_n = 1 + 2 \cos(2\pi/(n+3))$ in closed form (Theorem 2), and its affine representation $\rho_n : W_n \rightarrow \text{Aff}(\mathbb{R}^n)$ is unconditionally faithful on every leg tuple with pairwise distinct positive rational coordinates (Theorem 8). The faithfulness proof rests on a uniform Schur-complement determinant identity $\det Q_n = -\prod_i a_i^2$ (Lemma 6) together with rank-0 Mordell descent on three Diophantine quartics whose Jacobians are Cremona curves 24a1 and 72a2.

1. Setting

Fix $n \geq 2$. An n -dimensional right-corner orthoscheme $O = O(a_1, \dots, a_n)$ is the simplex in $\mathbb{R}^n$ with vertex sequence

$$V_0 = 0, \quad V_k = V_{k-1} + a_k e_k \quad (1 \leq k \leq n),$$

where $e_1, \dots, e_n$ is the standard basis and $(a_1, \dots, a_n) \in \mathbb{Q}_{>0}^n$. The $n+1$ bounding hyperplanes carry reflections $R_0, R_1, \dots, R_n$; we write $W_n = W(O) := \langle R_0, \dots, R_n \rangle \leq \text{Isom}(\mathbb{R}^n)$. The orbit of O under W_n is explored by breadth-first search in the Cayley graph relative to $\{R_0, \dots, R_n\}$, and

$$\mathcal{O}_d(O) = \{g \cdot O : g \in W_n, \text{ minimal word-length } d\}, \quad u_d^n(O) := |\mathcal{O}_d(O)|, \quad u_0 = 1.$$

All distances are computed in exact rational arithmetic.¹

Independence of legs. A symbolic computation in $\mathbb{Q}(a_1, \dots, a_n)$ shows that the perpendicular face-pair structure of O is independent of the leg values: pairs (i, j) with $|i - j| \geq 2$ commute identically, while pairs $(i, i+1)$ have a leg-dependent dihedral angle with $m_{i,i+1} = \infty$ for generic legs. The abstract Coxeter group is therefore the right-angled Coxeter group on the path diagram P_{n+1} :

$$W_n \cong \langle R_0, \dots, R_n \mid R_i^2 = 1, (R_i R_j)^2 = 1 \text{ for } |i - j| \geq 2 \rangle, \quad (1)$$

the same group for every nondegenerate orthoscheme of dimension n . This is the standard right-angled Coxeter system on the commutation graph that is the complement of P_{n+1} .

*Independent researcher, Italy. vicobonfioli@gmail.com. <https://elveclo.github.io/home/posts/>.

¹A companion paper treats the dimension- $n = 2$ case in detail, including a universality theorem and eight length-10 affine relations specific to planar right triangles, none of which generalise to $n \geq 3$. The present paper is self-contained and does not depend on that companion.

Steinberg upper bound. Steinberg’s word-growth formula (see Humphreys [1], §5.12) gives $1/W_n(t^{-1}) = \sum_I (-1)^{|I|}/W_I(t)$, summed over cliques I of the commutation graph G_n (cliques correspond to subsets of pairwise-commuting generators, hence to finite parabolics $W_I \cong (\mathbb{Z}/2)^{|I|}$). The number of k -cliques of G_n is $\binom{n-k+2}{k}$, so substituting $W_I(t) = (1+t)^{|I|}$:

$$\frac{1}{W_n(t^{-1})} = \sum_{k \geq 0} (-1)^k \binom{n-k+2}{k} \frac{1}{(1+t)^k}. \quad (2)$$

For every faithful or non-faithful affine representation of W_n one has $u_d^n \leq [t^d] W_n(t)$. The aim of §§2–4 is (a) to identify $W_n(t)$ in closed form, (b) to certify equality $u_d^n = [t^d] W_n(t)$ unconditionally for distinct positive rational legs, and (c) to read off the depth at which the inequality first becomes an equality (the collision-depth).

Remark 1 (Scope, and relation to the planar case). This is a companion note. The planar case $n = 2$ — where the group is $V_4 * \mathbb{Z}/2$, the metric is the lamplighter-type formula, and the growth series is transcendental — is treated in the main paper and is not reproved here; what follows is the part of the picture that is uniform in the dimension. Three things are established: the closed form for r_n (§2), the uniform Schur-complement determinant identity (§1) which is what makes Class C faithfulness unconditional in every dimension (§1), and the dimension-independent collision-depth invariant (§4). Two things are deliberately *not* attempted: the transcendence questions, which are special to $n = 2$; and any statement about non-right-corner orthoschemes.

2. The closed-form growth rate r_n

Theorem 2 (Closed-form n -dim Coxeter rate). *For every $n \geq 2$, the abstract orbit-growth rate of W_n is*

$$r_n = 1 + 2 \cos\left(\frac{2\pi}{n+3}\right).$$

Proof. Start from Steinberg’s identity (2). Substitute $y = -1/(1+t)$, so that $(1+t)^{-k} = (-y)^k = (-1)^k y^k$ and the two $(-1)^k$ factors cancel:

$$\frac{1}{W_n(t^{-1})} = \sum_{k \geq 0} \binom{n-k+2}{k} y^k =: f_n(y).$$

The polynomial $f_n(y)$ is the standard Fibonacci–Pell polynomial: writing $P_0(y) = 0$, $P_1(y) = 1$, and $P_m(y) = P_{m-1}(y) + y P_{m-2}(y)$ for $m \geq 2$, Pascal’s identity $\binom{n-k+2}{k} = \binom{n-k+1}{k} + \binom{n-k+1}{k-1}$ gives $f_n(y) = P_{n+3}(y)$: the two sides satisfy the same three-term recursion in n , and agree at $n = -2, -1$. The Pascal step is machine-checked (`NdimRate.lean`, `pascal_step` and `coeff_pascal`; `pascal_step` depends on *no* axioms), so the only unformalised part of this paragraph is the induction that assembles it. The characteristic equation $x^2 = x + y$ has roots $\alpha, \beta = (1 \pm \sqrt{1+4y})/2$, and the Binet formula yields $P_m(y) = (\alpha^m - \beta^m)/(\alpha - \beta)$; hence $P_m(y) = 0$ iff $(\alpha/\beta)^m = 1$. Setting $\alpha/\beta = e^{2\pi i j/m}$ and solving for y :

$$y_j = -\frac{1}{4 \cos^2(\pi j/m)}, \quad j = 1, 2, \dots, \lfloor (m-1)/2 \rfloor.$$

Translating back via $y = -1/(1+t)$ and using $4 \cos^2 \theta - 1 = 1 + 2 \cos(2\theta)$:

$$t_j = 4 \cos^2\left(\frac{\pi j}{n+3}\right) - 1 = 1 + 2 \cos\left(\frac{2\pi j}{n+3}\right), \quad 1 \leq j \leq \lfloor (n+2)/2 \rfloor.$$

The asymptotic growth rate is the largest root, attained at $j = 1$. □ □

Remark 3 (Prior art). Both ingredients of Theorem 2 are classical and are used as such. Steinberg’s formula for the Poincaré series of a Coxeter group is standard, and the Fibonacci–Pell polynomials $P_m(y)$, their Binet formula, and the location of their roots at $y_j = -1/(4 \cos^2(\pi j/m))$ are classical facts about that family; the values $1 + 2 \cos(2\pi j/m)$ are likewise the familiar Chebyshev-type quantities. What is asserted here is the *identification*: that the abstract orbit-growth rate of the n -dimensional right-corner orthoscheme group is the largest of those roots at $m = n + 3$, together with the collision-depth invariant of §4. We have not located that identification in the literature, but we make no claim of novelty for the ingredients, and a reader who knows the Coxeter growth-series literature should treat the closed form as an assembly of known pieces rather than a new evaluation.

Remark 4 (The Class C orthoscheme sequences in the OEIS). For pairwise distinct legs the layer counts u_d^n are recorded in the On-Line Encyclopedia of Integer Sequences: the planar base case ($n = 2$) is A396406, and the orthoschemes are A397439 ($n = 3$), A397437 ($n = 4$), A396927 ($n = 5$), and A397438 ($n = 6$), with the $n = 7$ entry in preparation. Each is a rational generating function whose dominant pole is $1/r_n$ for r_n of Theorem 2; for instance $(1+x)^2/(1-2x)$ at $n = 3$, $(1+x)^3/(1-2x-x^2+x^3)$ at $n = 4$, $(1+x)^4/(1-3x+3x^3)$ at $n = 6$, and $(1+x)^4/(1-4x+3x^2+2x^3-x^4)$ at $n = 7$. The planar case $n = 2$ and its lamplighter structure are treated in a companion paper.

Corollary 5 (asymptotic regime). $r_n \rightarrow 3$ as $n \rightarrow \infty$, with the explicit residual

$$3 - r_n = 4 \sin^2\left(\frac{\pi}{n+3}\right) \sim \frac{4\pi^2}{(n+3)^2}.$$

The closed form gives quadratic-surd values exactly when $\varphi(n+3) \leq 4$ (Euler’s totient), namely $n+3 \in \{5, 6, 8, 10, 12\}$; see Table 1.

n	r_n	algebraic form
2	1.6180...	$\varphi = (1 + \sqrt{5})/2$ (golden)
3	2.0000	exactly 2
4	2.2470...	$1 + 2 \cos(2\pi/7)$, cubic
5	2.4142...	$1 + \sqrt{2}$ (silver)
6	2.5321...	$1 + 2 \cos(2\pi/9)$, cubic
7	2.6180...	$\varphi^2 = (3 + \sqrt{5})/2$
8	2.6825...	$1 + 2 \cos(2\pi/11)$, degree 5
9	2.7321...	$1 + \sqrt{3}$
10	2.7709...	$1 + 2 \cos(2\pi/13)$, degree 6

Table 1: The closed-form rate $r_n = 1 + 2 \cos(2\pi/(n+3))$ for the right-corner orthoscheme reflection group, verified to 50-digit precision for n up to 500.

3. Unconditional Class C faithfulness

The Steinberg series (2) is an upper bound: only when the affine representation $\rho_n : W_n \rightarrow \text{Aff}(\mathbb{R}^n)$ is faithful and no additional Coxeter relation closes does the BFS orbit count u_d^n saturate $[t^d] W_n(t)$. Following the terminology of the 3D analysis, we call a leg tuple $(a_1, \dots, a_n)$ *Class C* when its coordinates are pairwise distinct. The main result of this section certifies that the Class C case is unconditional in every dimension.

3.1 The uniform Schur-complement determinant identity

The Lorentzian Gram matrix of the orthoscheme in staircase coordinates — with simple roots $n_0 = e_1$, $n_i = a_i e_{i+1} - a_{i+1} e_i$ ($1 \leq i \leq n-1$), $n_n = e_n$, and the standard $(n, 1)$ endpoint

correction — is the tridiagonal matrix Q_n of size $(n+1) \times (n+1)$ given by

$$\begin{aligned} Q_n[0,0] &= 1 - a_1^2, & Q_n[i,i] &= a_i^2 + a_{i+1}^2 \quad (1 \leq i \leq n-1), & Q_n[n,n] &= 1, \\ Q_n[0,1] &= a_2, & Q_n[i,i+1] &= a_i a_{i+2} \quad (1 \leq i \leq n-2), & Q_n[n-1,n] &= a_{n-1}. \end{aligned}$$

Lemma 6 (Uniform Schur-complement determinant identity). *For every $n \geq 3$ and every positive tuple $(a_1, \dots, a_n)$,*

$$\det Q_n = - \prod_{i=1}^n a_i^2.$$

Proof. Let $D_k = \det Q_n[:k+1, :k+1]$ be the leading principal minor of order $k+1$. Tridiagonal expansion along the last row gives

$$D_k = Q_n[k,k] D_{k-1} - Q_n[k-1,k]^2 D_{k-2} \quad (1 \leq k \leq n), \quad (\dagger)$$

with $D_{-1} = 1$. We claim

$$D_k = P_k := \left(\prod_{i=1}^k a_i^2 \right) \left(1 - \sum_{i=1}^{k+1} a_i^2 \right) \quad \text{for every } 1 \leq k \leq n-1. \quad (\star)$$

Base case. $D_0 = 1 - a_1^2$, and $D_1 = (1 - a_1^2)(a_1^2 + a_2^2) - a_2^2 = a_1^2(1 - a_1^2 - a_2^2) = P_1$.

Interior step (uniform). For $2 \leq k \leq n-1$ the recurrence $(\dagger)$ reads $D_k = (a_k^2 + a_{k+1}^2) D_{k-1} - (a_{k-1} a_{k+1})^2 D_{k-2}$. Introduce the abstract symbols $S_{k-1} := \sum_{i=1}^{k-1} a_i^2$ and $P_{k-2} := \prod_{i=1}^{k-2} a_i^2$; substituting the closed forms $(\star)$ reduces the step to the polynomial identity

$$\begin{aligned} P_{k-2} a_{k-1}^2 a_k^2 (1 - S_{k-1} - a_k^2 - a_{k+1}^2) &= (a_k^2 + a_{k+1}^2) P_{k-2} a_{k-1}^2 (1 - S_{k-1} - a_k^2) \\ &\quad - (a_{k-1} a_{k+1})^2 P_{k-2} (1 - S_{k-1}) \end{aligned}$$

in the five independent symbols $(a_{k-1}, a_k, a_{k+1}, S_{k-1}, P_{k-2})$. This identity is independent of k and of n . It, the two base cases, the induction that assembles them into $(\star)$, and the boundary step yielding $\det Q_n$, are machine-checked in Lean 4 over an arbitrary commutative ring: see `OrthoschemeDet.lean`, theorem `det_Q_of_recurrence` (Section 7). No positivity, ordering, or characteristic hypothesis is used.

Final step (uniform). At $k = n$ the recurrence uses $Q_n[n,n] = 1$ and $Q_n[n-1,n]^2 = a_{n-1}^2$, giving $D_n = D_{n-1} - a_{n-1}^2 D_{n-2}$. With $S_{n-1} = \sum_{i=1}^{n-1} a_i^2$, $P_{n-2} = \prod_{i=1}^{n-2} a_i^2$,

$$D_{n-1} - a_{n-1}^2 D_{n-2} = P_{n-2} a_{n-1}^2 [(1 - S_{n-1} - a_n^2) - (1 - S_{n-1})] = -P_{n-2} a_{n-1}^2 a_n^2 = - \prod_{i=1}^n a_i^2.$$

This final identity is again a single polynomial identity in the four independent symbols $(a_{n-1}, a_n, S_{n-1}, P_{n-2})$, independent of n . The three identities (base, interior, final) jointly close the induction for every $n \geq 3$, with no per- n verification required. $\square$ $\square$

3.2 Diophantine emptiness on three Cremona curves

A non-perpendicular dihedral pair $(i, i+1)$ in the staircase orthoscheme has $\cos^2 \theta_{i,i+1} \in \mathbb{Q}$; by the extended Niven theorem (a consequence of Conway–Jones [4]: if $\theta \in \mathbb{Q}\pi$ and $\cos^2 \theta \in \mathbb{Q}$, then $\cos \theta \in \{0, \pm 1/2, \pm \sqrt{2}/2, \pm \sqrt{3}/2, \pm 1\}$), any non-trivial closing relation $(R_i R_{i+1})^m = 1$ forces $\cos^2 \theta_{i,i+1} \in \{0, 1/4, 1/2, 3/4, 1\}$. The boundary values $\{0, 1\}$ are immediately incompatible with positive distinct legs. The remaining three cut out, in the two legs flanking the interior pair (with $X = a_i^2$, $Z = a_{i+2}^2$), the Diophantine quartics

$$\mathcal{V}_{1/4} : X^2 + 14XZ + Z^2 = m^2, \quad \mathcal{V}_{1/2} : X^2 + 6XZ + Z^2 = k^2, \quad \mathcal{V}_{3/4} : 9X^2 + 30XZ + 9Z^2 = k^2.$$

Lemma 7. *Each of $\mathcal{V}_{1/4}$, $\mathcal{V}_{1/2}$, $\mathcal{V}_{3/4}$ is empty over distinct-coordinate $\mathbb{Q}_{>0}^2$ (i.e., the only rational solutions have $XZ = 0$ or $X = Z$).*

Proof. $\mathcal{V}_{1/2}$. The quartic $X^2 + 6XZ + Z^2 = k^2$ is birationally equivalent to the elliptic curve $E_6 : y^2 = x^3 - x$ (Mordell [5], §17). E_6 has Mordell–Weil rank 0 over $\mathbb{Q}$ by Fermat’s classical descent (Cassels [6], Ch. II), and its full rational-point set is the 2-torsion $\{\infty, (0, 0), (\pm 1, 0)\}$, pulling back to $XZ = 0$ or $X = Z$.

$\mathcal{V}_{1/4}$. The Jacobian elliptic curve is

$$E_{1/4} : y^2 = x(x - 12)(x - 16),$$

which is Cremona 24a1 (conductor 24, $j = 35\,152/9$, Mordell–Weil rank 0 over $\mathbb{Q}$, torsion $\mathbb{Z}/4 \oplus \mathbb{Z}/2$); see [7] and the LMFDB record [8]. Each of its eight rational points pulls back to $XZ = 0$ or $X = Z$.

$\mathcal{V}_{3/4}$. The quartic $9X^2 + 30XZ + 9Z^2 = k^2$ admits a standard conic-to-Weierstrass substitution and reduces to

$$E_{3/4} : y^2 = x(x - 300)(x - 1200),$$

Cremona 72a2 (conductor 72, same j -invariant $35\,152/9$ as $E_{1/4}$ — they are quadratic twists — and also rank 0 with torsion $\mathbb{Z}/2 \oplus \mathbb{Z}/2$); see [7, 9]. Each torsion point again pulls back to a trivial quartic solution. $\square$ $\square$

3.3 Faithfulness theorem

Theorem 8 (Class C faithfulness, all $n \geq 3$, unconditional). *For every $n \geq 3$ and every leg tuple $(a_1, \dots, a_n) \in \mathbb{Q}_{>0}^n$ with pairwise distinct coordinates, the affine representation $\rho_n : W_n \rightarrow \text{Aff}(\mathbb{R}^n)$ is faithful, no Coxeter relation closes beyond the $\binom{n}{2} - n$ perpendicular pairs, and the BFS layer counts saturate the abstract Coxeter sphere size:*

$$u_d^n(O) = [t^d] W_n(t) \quad \text{for every } d \geq 0.$$

Proof. By Lemma 6, $\det Q_n = -\prod_{i=1}^n a_i^2 \neq 0$ for every positive tuple, so the Lorentzian Coxeter system is non-degenerate. Tits’ theorem on faithfulness of the geometric representation (Humphreys [1], Thm. 5.4) then gives faithfulness of ρ_n on every positive-rational leg tuple.

It remains to exclude additional Coxeter relations. Each non-perpendicular pair is of type $(i, i + 1)$, $0 \leq i \leq n - 1$; its cosine satisfies $\cos^2 \theta_{i,i+1} \in \mathbb{Q}$ (boundary pairs $(0, 1)$ and $(n - 1, n)$ involve two legs, interior pairs involve three). By the extended Niven theorem, a closing Coxeter relation requires $\cos^2 \theta_{i,i+1} \in \{0, 1/4, 1/2, 3/4, 1\}$. The boundary values $\{0, 1\}$ contradict positivity of legs. At boundary pairs, the values $1/4, 3/4$ force $a_i = a_{i+1}/\sqrt{3}$ or $a_i = a_{i+1}$, the first irrational and the second excluded by the distinct-coordinate hypothesis; the value $1/2$ forces $a_i = a_{i+1}$, again excluded. At each interior pair, the three values $1/4, 1/2, 3/4$ define exactly the three Diophantine quartics $\mathcal{V}_{1/4}, \mathcal{V}_{1/2}, \mathcal{V}_{3/4}$ in the two flanking legs (a_i, a_{i+2}) (the dihedral analysis is uniform in n). By Lemma 7 each of the three is empty over distinct-coordinate $\mathbb{Q}_{>0}^2$.

Hence outside the symmetry locus $\{a_i = a_{i+1}\}$ for some i , no non-perpendicular dihedral is a rational multiple of π , no closing relation appears, and $u_d^n(O) = [t^d] W_n(t)$. $\square$ $\square$

4. Dimension-independent collision-depth

For a leg sequence $(a_1, \dots, a_n)$ define the *collision-depth* as

$$\text{cd}_n(a_1, \dots, a_n) = \inf \{ d \geq 0 : u_d^n < [t^d] W_n(t) \} \in \{1, 2, 3, \dots, \infty\}.$$

By Theorem 8, $\text{cd}_n = \infty$ for every pairwise-distinct tuple. The next theorem extends the description to every leg multiplicity pattern and shows that, in spite of the multiplicity partition having $p(n)$ possible shapes, the collision-depth takes only three values, determined by two local positional statistics.

For $(a_1, \dots, a_n) \in \mathbb{Q}_{>0}^n$ set

$$\begin{aligned} e(a_1, \dots, a_n) &= \#\{i \in \{1, n-1\} : a_i = a_{i+1}\} \in \{0, 1, 2\}, \\ t(a_1, \dots, a_n) &= \#\{i : 1 \leq i \leq n-2, a_i = a_{i+1} = a_{i+2}\}. \end{aligned}$$

Thus e counts equal pairs at the two *endpoint* (boundary) positions of the staircase, and t counts *interior triples* of three consecutive equal legs. An equal pair sitting strictly in the interior of the staircase — such as $a_2 = a_3$ in $(1, 2, 2, 3)$ at $n = 4$ — contributes to neither e nor t .

Theorem 9 (Dimension-independent collision-depth, all $n \geq 3$). *For every $n \geq 3$ and every leg sequence $(a_1, \dots, a_n) \in \mathbb{Q}_{>0}^n$,*

$$\text{cd}_n(a_1, \dots, a_n) = \begin{cases} \infty & \text{if } e = t = 0, \\ 4 & \text{if } e \geq 1, t = 0, \\ 3 & \text{if } t \geq 1. \end{cases}$$

In particular, $\text{cd}_n \in \{\infty, 4, 3\}$ for every n , and the value depends only on the local positional statistics (e, t) .

Proof. The non-perpendicular dihedral pairs of $O = O(a_1, \dots, a_n)$ are (R_i, R_{i+1}) , $0 \leq i \leq n-1$. Their cosines have the following uniform form:

$$\begin{aligned} \cos \theta_{0,1} &= \frac{a_2}{\sqrt{a_1^2 + a_2^2}}, & \cos \theta_{n-1,n} &= \frac{a_{n-1}}{\sqrt{a_{n-1}^2 + a_n^2}}, \\ \cos \theta_{i,i+1} &= \frac{a_i a_{i+2}}{\sqrt{(a_i^2 + a_{i+1}^2)(a_{i+1}^2 + a_{i+2}^2)}} & & \text{for } 1 \leq i \leq n-2. \end{aligned}$$

Endpoint pair $\Rightarrow \text{cd} = 4$. An endpoint pair $a_1 = a_2$ (or $a_{n-1} = a_n$) makes the boundary cosine $1/\sqrt{2}$, so the dihedral is $\pi/4$ and the Coxeter relation $(R_0 R_1)^4 = 1$ (resp. $(R_{n-1} R_n)^4 = 1$) closes. The parabolic $\langle R_0, R_1 \rangle$ then has order 8 rather than infinite, and the first collision in the orbit BFS appears at length 4, giving $\text{cd}_n = 4$.

Interior triple $\Rightarrow \text{cd} = 3$. An interior triple $a_i = a_{i+1} = a_{i+2}$ ($1 \leq i \leq n-2$) makes the interior cosine $\cos \theta_{i,i+1} = a^2/(2a^2) = 1/2$ where $a := a_i$, so the dihedral is $\pi/3$ and the braid relation $R_i R_{i+1} R_i = R_{i+1} R_i R_{i+1}$ closes at length 3. Since a triple subsumes two pairs, the $\pi/3$ relation appears before any $\pi/4$ relation that the same triple might also induce at a boundary position, giving $\text{cd}_n = 3$.

Interior pair only $\Rightarrow \text{cd} = \infty$. Suppose $(e, t) = (0, 0)$, but the sequence is not Class C; then some interior pair $a_j = a_{j+1}$ exists with $1 \leq j \leq n-1$, neither boundary position nor part of a triple. At each non-perpendicular pair $(i, i+1)$ the cosine depends on the flanking legs, which are by hypothesis not all equal. At the two interior pairs flanking the equal pair — $(j-1, j)$ and $(j+1, j+2)$ (whichever exist) — the cosine takes the $a_i a_{i+2} / \sqrt{(a_i^2 + a_{i+1}^2)(a_{i+1}^2 + a_{i+2}^2)}$ form with (a_i, a_{i+2}) distinct positive rationals (otherwise the equal pair would extend to a triple, contradicting $t = 0$). By the extended Niven theorem and Lemma 7, no closing relation is possible: each of the candidate values $1/4, 1/2, 3/4$ for $\cos^2$ defines one of the three Diophantine quartics, all empty over distinct $\mathbb{Q}_{>0}^2$. At every non-flanking pair the flanking-leg analysis is the same. Hence no Coxeter relation closes, and by Tits faithfulness on the resulting non-degenerate Coxeter system (Lemma 6 plus Humphreys [1], Thm. 5.4) $u_d^n = [t^d] W_n(t)$ for every d , i.e., $\text{cd}_n = \infty$.

Combining the three cases proves the tri-classification. $\square$ $\square$

Remark 10 (the R -rule fails in $n \geq 4$). A natural but incorrect heuristic is to set cd_n to a finite value as soon as some consecutive equal pair appears, i.e., as soon as the run-length $R := \max_i \#\{j : a_j = a_i\}$ is at least 2. At $n = 3$ this R -rule coincides with the (e, t) -rule because every consecutive equal pair in a 3-term sequence is automatically at an endpoint. At $n \geq 4$ the two rules diverge: the orthoscheme $(1, 2, 2, 3)$ has an interior equal pair ($R = 2$, but $(e, t) = (0, 0)$) and Theorem 9 predicts $\text{cd}_4 = \infty$; empirically the BFS sequence $1, 5, 14, 33, 75, 169, \dots$ is bit-identical to the Class C sequence $1, 5, 14, 33, 75, 169$ at $(1, 2, 3, 5)$ through every observed depth, falsifying the R -rule.

5. Atlas: small dimensions and the (e, t) classes

We collect a small atlas of representative leg sequences for $n = 3, 4, 5, 6$, illustrating each attainable (e, t) class. The layer counts are exact-rational breadth-first search; in every case the observed collision depth agrees with the value cd_n predicted by Theorem 9.

n	leg sequence	(e, t)	cd_n	first six BFS layer counts
3	(2, 3, 5)	(0, 0)	∞	1, 4, 9, 18, 36, 72
3	(1, 1, 2)	(1, 0)	4	1, 4, 9, 18, 35, 67
3	(1, 1, 1)	(2, 1)	3	1, 4, 9, 17, 28, 42
4	(1, 2, 3, 5)	(0, 0)	∞	1, 5, 14, 33, 75, 169
4	(1, 2, 2, 3)	(0, 0)	∞	1, 5, 14, 33, 75, 169
4	(1, 1, 2, 3)	(1, 0)	4	1, 5, 14, 33, 74, 163
4	(1, 1, 2, 2)	(2, 0)	4	1, 5, 14, 33, 73, 157
4	(1, 1, 1, 2)	(1, 1)	3	1, 5, 14, 32, 67, 135
5	(1, 2, 3, 5, 7)	(0, 0)	∞	1, 6, 20, 54, 136, $\dots$
5	(1, 2, 2, 3, 5)	(0, 0)	∞	1, 6, 20, 54, 136, $\dots$
5	(1, 1, 2, 3, 5)	(1, 0)	4	1, 6, 20, 54, 135, 327
5	(2, 1, 1, 1, 2)	(0, 1)	3	1, 6, 20, 53, 128, 297
6	(1, 1, 2, 3, 2, 2)	(2, 0)	4	1, 7, 27, 82, 224, 581

Table 2: A small atlas of representative orthoschemes in $n = 3, 4, 5, 6$ illustrating each (e, t) class. The collision-depth value $\text{cd}_n \in \{\infty, 4, 3\}$ is determined uniformly by (e, t) in every dimension, by Theorem 9. The two $n = 4$ sequences with $(e, t) = (0, 0)$ have bit-identical BFS counts through every observed depth despite differing in run structure ($(1, 2, 3, 5)$ has $R = 1$ but $(1, 2, 2, 3)$ has $R = 2$; only (e, t) is the right invariant).

The all-distinct rows of the table saturate the Steinberg upper bound $[t^d] W_n(t)$ by Theorem 8; for $n = 3$ this reproduces the classical sequence $u_d = 9 \cdot 2^{d-2}$ for $d \geq 2$.

6. Related work and open questions

Path Coxeter groups and the Cannon–Floyd–Parry framework. The closed form $r_n = 1 + 2 \cos(2\pi/(n+3))$ is the structural signature of a linear (path-shaped) Coxeter group whose Poincaré-series denominator reduces to a tridiagonal Chebyshev-type characteristic polynomial. Cannon and Wagreich [10] and Floyd and Plotnick [11] developed the growth-rate calculus of such groups; Theorem 2 fits into their framework with parameter $m = n + 3$ and is recovered cleanly through the Fibonacci–Pell substitution of §2.

Salem-type family on the surface-dynamics side. The family of algebraic integers $\{1 + 2\cos(2\pi/m)\}_{m \geq 3}$ to which r_n belongs also arises in birational surface dynamics. Diller and Favre [13] introduced the dynamical-degree framework for bimeromorphic maps of surfaces; McMullen [14] showed that this family arises as dynamical degrees of automorphisms of blowups of $\mathbb{P}^2$ associated with the E_n T-shaped Coxeter graph (different from the path Coxeter graph here); and Bedford–Kim [15] realised specific members through linear-fractional recurrences. The rate r_n lies in this family, but the right-corner orthoscheme reflection group is not known to be realised as a McMullen involution-product on a rational surface; the coincidence is at the level of the algebraic-number family. A birational-geometry construction realising r_n as a dynamical degree on a $\mathbb{P}^2$ -blowup would be of independent interest.

Open questions. (1) *Non-orthoscheme reflection groups.* Does an analogue of Theorem 9 hold for other families of simplicial reflection chambers in $\mathbb{R}^n$ — for example, Coxeter orthoschemes with bent paths, or simplices arising from non-path-shaped Coxeter diagrams? The proof relies on the right-angled-with-one- ∞ structure of the path Coxeter diagram; how robust is the (e, t) -classification under deformation of the diagram?

(2) *Classes of intermediate growth.* In every dimension $n \geq 3$, partial multiplicity patterns (neither distinct nor saturated by triples) produce intermediate growth rates strictly between two consecutive values of r_n . An algebraic characterisation of these intermediate rates — in particular, whether they are algebraic over $\mathbb{Q}$ at all — is open.

(3) *Beyond (e, t) .* Theorem 9 shows that the collision-depth is determined by the maximum multiplicity of *positional* structure (e, t) . Is there a finer positional invariant that captures the entire u_d^n sequence (not only cd_n) in terms of $(a_1, \dots, a_n)$?

Acknowledgments

Computational assistance from large language model assistants was used at multiple stages of this work; all mathematical claims have been verified by the author, who is solely responsible for the content of this paper and for any errors.

7. Machine verification

The algebraic content of Lemma 6 and of Theorem 2 is formalised in Lean 4 with Mathlib. Each file ends with `#print axioms`; every theorem below reports only the standard axioms `propext`, `Classical.choice`, `Quot.sound` (two report strictly fewer), and no file contains `sorry`.

Paper result	Lean declaration	File
Lemma 6, closed form ($\star$)	<code>minor_closed_form</code>	<code>OrthoschemeDet.lean</code>
Lemma 6, boundary step	<code>det_Q</code>	<code>OrthoschemeDet.lean</code>
Lemma 6, full statement	<code>det_Q_of_recurrence</code>	<code>OrthoschemeDet.lean</code>
Thm. 2, root of $x^2 = x + y$	<code>sq_eq</code>	<code>NdimRate.lean</code>
Thm. 2, Binet formula	<code>pell_binet</code>	<code>NdimRate.lean</code>
Thm. 2, root translation	<code>root_translation</code>	<code>NdimRate.lean</code>
Corollary (asymptotic regime)	<code>three_sub_r</code>	<code>NdimRate.lean</code>
Thm. 2, Pascal step	<code>pascal_step</code> (no axioms)	<code>NdimRate.lean</code>
Thm. 2, coefficient form	<code>coeff_pascal</code>	<code>NdimRate.lean</code>

What is not formalised. The step identifying the abstract orbit-growth rate with the largest root t_1 of W_n is a dominant-pole argument on the rational generating function; it is proved on paper only. Likewise the combinatorial identification $f_n(y) = P_{n+3}(y)$ via Pascal’s identity is a paper proof. Everything downstream of those two inputs — the Binet formula, the location of the roots, and the translation to $1 + 2\cos(2\pi j/(n+3))$ — is machine-checked.

References

- [1] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics 29, Cambridge University Press, 1990.
- [2] R. Steinberg, *Endomorphisms of Linear Algebraic Groups*, Memoirs of the American Mathematical Society 80, 1968.
- [3] M. W. Davis, *The Geometry and Topology of Coxeter Groups*, London Mathematical Society Monographs 32, Princeton University Press, 2008.
- [4] J. H. Conway, A. J. Jones, *Trigonometric Diophantine equations (On vanishing sums of roots of unity)*, Acta Arith. 30 (1976), 229–240.
- [5] L. J. Mordell, *Diophantine Equations*, Pure and Applied Mathematics 30, Academic Press, 1969.
- [6] J. W. S. Cassels, *Lectures on Elliptic Curves*, London Mathematical Society Student Texts 24, Cambridge University Press, 1991.
- [7] J. E. Cremona, *Algorithms for Modular Elliptic Curves*, 2nd edition, Cambridge University Press, 1997.
- [8] LMFDB Collaboration, elliptic curve 24a1 over $\mathbb{Q}$, <https://www.lmfdb.org/EllipticCurve/Q/24/a/1>.
- [9] LMFDB Collaboration, elliptic curve 72a2 over $\mathbb{Q}$, <https://www.lmfdb.org/EllipticCurve/Q/72/a/2>.
- [10] J. W. Cannon, P. Wagreich, *Growth functions of surface groups*, Math. Ann. 293 (1992), no. 2, 239–257.
- [11] W. J. Floyd, S. P. Plotnick, *Growth functions on Fuchsian groups and the Euler characteristic*, Invent. Math. 88 (1987), no. 1, 1–29.
- [12] W. Parry, *Growth series of some hyperbolic Coxeter groups*, Geom. Dedicata 32 (1989), 75–96.
- [13] J. Diller, C. Favre, *Dynamics of bimeromorphic maps of surfaces*, Amer. J. Math. 123 (2001), no. 6, 1135–1169.
- [14] C. T. McMullen, *Dynamics on blowups of the projective plane*, Publ. Math. Inst. Hautes Études Sci. 105 (2007), 49–89.
- [15] E. Bedford, K. Kim, *Dynamics of rational surface automorphisms: linear fractional recurrences*, J. Geom. Anal. 19 (2009), no. 3, 553–583.
- [16] H. S. M. Coxeter, *Discrete groups generated by reflections*, Annals of Math. 35 (1934), 588–621.