

Roots of d -matching polynomials and the spectrum of the universal cover: a closed form at first Betti number one, and a refuted conjecture

Vico Bonfioli*

August 2026

Abstract

Hall, Puder and Sawin place the roots of the d -matching polynomial $\mu_{d,G}$ in the interval $[-\rho, \rho]$, ρ the spectral radius of the universal cover T . We conjectured that they lie in $\text{spec}(T)$ itself, a proper subset of that interval whenever T has a spectral gap: no root ever enters a gap. This strengthens their theorem and is the matching-polynomial form of their Question 6.3.

We prove it for every connected G of first Betti number one and every d , by giving $\mu_{d,G}$ in closed form.

In general it is false. The counterexample is due to Chris Hall [12]: a bipartite graph on 41 vertices with $\mu_G = x^{21}(x^4 - 11x^2 + 25)^4(x^2 - 5)(x^2 - 11)$, where $\sqrt{5}$ lies in an internal gap of $\text{spec}(T)$.

The natural repairs, minimum degree three and 3-connectivity, are open. Our contribution is to show that the obvious attack on them cannot work. A separator of size k met by p identical branches forces $A^{p-k} \mid \mu_G$, where A is the branch's matching polynomial (Lemma 7), so it produces roots of μ_G at will; but when the branch is a tree that same configuration is a θ -Aomoto subset in the sense of Banks, Garza-Vargas and Mukherjee, so the root it produces is an *eigenvalue* of the cover (Proposition 10). The condition $p > k$ that makes the divisor nontrivial is literally the Aomoto inequality $|\partial S| < \text{cc}(G[S])$. The mechanism manufactures points of $\text{spec}(T)$, not counterexamples.

This explains the one feature of Hall's family that had resisted explanation, the pendant leaves. They create the root $\sqrt{5}$ and simultaneously double the boundary of the relevant star, carrying it away from the Aomoto inequality; removing them destroys both. A counterexample must give its branches a private boundary, which is a condition on attachment rather than on degrees, and that is where a repair should be sought.

Everything asserted is machine-checked in Lean 4/Mathlib unless labelled otherwise. Supplementary material is in [18]; the biregular case is [17].

2020 Mathematics Subject Classification. Primary 05C31; Secondary 05C50, 05C70, 68V20.

Keywords. matching polynomial, d -matching polynomial, universal cover, spectral gap, graph covering, interlacing families, Lean 4, formalization.

*Lean 4 sources: `RamaLean/` in the `rama-notes` repository. Correspondence: `vicobonfioli@gmail.com`.

Code and data availability. All computations, the Lean 4/Mathlib sources, and the scripts backing every numerical statement are in the `rama-notes` repository. Each numerical claim names the script that produced it; every named script is executed and its output compared against a recorded baseline by `code/cite_check.py`, and every Lean declaration cited is checked for its axiom base by `code/audit_papers.py`.

Status conventions. Claims here are of three kinds, and the text says which each time rather than leaving it to the reader. A result described as *formalized*, or cited by a Lean declaration name, is machine-checked in Lean 4/Mathlib: the sources are in `RamaLean/` of the accompanying repository, carry no `sorry`, and rest only on the standard axioms `propext`, `Classical.choice` and `Quot.sound`. A claim described as *measured*, *verified to* a stated tolerance, or *checked numerically* is computational evidence and is not a proof; the script producing it is named, and every named script is run and its output compared against a recorded baseline (`code/cite_check.py`). A claim called a *conjecture* is unproved, and is called that even where the evidence is strong.

1 Introduction and statement

The expected characteristic polynomial of a random lift is the object at the heart of the interlacing-families method of Marcus–Spielman–Srivastava [11], who used the $r = 2$ (signing) case to prove that bipartite Ramanujan graphs exist in every degree; Hall–Puder–Sawin [2] extended the framework to arbitrary coverings. We evaluate it in closed form for the cycle.

A *permutation r -lift* of a graph G replaces each vertex by r copies and each edge $\{u, v\}$ by the perfect matching $u_i \sim v_{\sigma(i)}$ of a permutation $\sigma \in S_r$ chosen independently and uniformly per edge. For the cycle C_n , gauge-fixing a spanning path reduces the lift to a single permutation $\sigma \in S_r$ on the closing edge, and the lift decomposes into disjoint cycles: each ℓ -cycle of σ contributes a cycle $C_{n\ell}$ (fixed points contribute copies of C_n). Since $\chi_{C_m}(x) = \det(xI - A_{C_m}) = 2T_m(x/2) - 2$, the expected characteristic polynomial is the expected cycle weight

$$\Phi_{n,r}(x) = \mathbb{E}_{\sigma \in S_r} \prod_{\ell \in \text{cycles}(\sigma)} (2T_{n\ell}(x/2) - 2). \quad (1)$$

Theorem 1. *For all $n \geq 1$, $r \geq 1$, with $Y = T_n(x/2)$,*

$$\Phi_{n,r}(x) = U_r(Y) - 2U_{r-1}(Y) + U_{r-2}(Y) = \underbrace{(2T_n(x/2) - 2)}_{\chi_{C_n}(x)} \cdot \underbrace{U_{r-1}(T_n(x/2))}_{\Psi_r(x)}.$$

The factored form identifies the quotient Ψ_r , by the theorem of Hall–Puder–Sawin [2] equating the expected new-eigenvalue characteristic polynomial with the d -matching polynomial ($d = r - 1$), as

$$\mu_{d,C_n}(x) = U_d(T_n(x/2)),$$

which is equivalent (through $\mathcal{P}_d(2t) = U_d(t)$, $\mathcal{C}_n(2t) = 2T_n(t)$ and the composition identity $U_{dn+n-1} = U_d(T_n)U_{n-1}$) to *Hall’s conjecture* $\mathcal{C}_{n,d} = \mathcal{P}_d(\mathcal{C}_n)$, proved combinatorially in [3]. That preprint has remained unrefereed since 2018 and carries no journal reference or DOI as of August 2026. The reason is not mathematical. Chris Hall, who posed the conjecture, has

told us (personal communication, August 2026) that Andrew Herring was his first doctoral student; Hall gave him the conjecture, Herring presented it as a problem at a workshop, and the group of [3] solved it there. They submitted the result, met a couple of rejections, and abandoned the attempt for want of motivation rather than for any defect. Hall regards the proof as correct, and so do we. We record the episode because a reader weighing what is established about this identity should know both that the combinatorial proof stands and that the present derivation is, as far as we can tell, the first refereed or machine-checked one. Our route is different and elementary; we make no use of matching-polynomial combinatorics.

What this paper does

The closed form is the starting point rather than the subject. Read as a statement about where the roots of $\mu_{d,G}$ sit, it says that for the cycle they land exactly in $\text{spec}(T)$, and not merely in the interval $[-\rho, \rho]$ that Hall–Puder–Sawin guarantee. The two differ whenever T has a spectral gap, and whether the stronger statement holds in general is the question here.

§3 proves it for every graph of first Betti number one and every d . §4 states it as a conjecture for all graphs. §5 is the counterexample, due to Chris Hall. §5.3 isolates the algebra behind it, and §5.4 shows that this algebra cannot be turned against the two natural repairs, because the configuration it needs is exactly an Aomoto subset. Supplementary material is in [18]; the biregular case, which survives the refutation, is [17].

2 Proof

Write $f(\ell) = 2T_\ell(Y) - 2$ with $Y = T_n(x/2)$; by the composition identity $T_{n\ell} = T_\ell \circ T_n$ this is the per-cycle factor in (1).

Step 1 (exponential formula). For any commutative ring R and $f : \mathbb{N} \rightarrow R$, the S_r -average $\Phi_r = \mathbb{E}_\sigma \prod_{\ell \in \text{cycles}(\sigma)} f(\ell)$ satisfies

$$r \Phi_r = \sum_{k=1}^r f(k) \Phi_{r-k}, \quad \Phi_0 = 1, \quad (2)$$

the coefficient form of $\sum_r \Phi_r z^r = \exp(\sum_\ell f(\ell) z^\ell / \ell)$. (Proof: group σ by cycle type λ with Cauchy’s count $r!/z_\lambda$, mark a part, and re-index; see the Lean file `Paper2ExpFormula.lean` for the fully detailed version.)

Step 2 (Chebyshev generating function). With $f(\ell) = 2T_\ell(Y) - 2$,

$$\sum_{\ell \geq 1} f(\ell) \frac{z^\ell}{\ell} = -\log(1 - 2Yz + z^2) + 2\log(1 - z) \implies \sum_{r \geq 0} \Phi_r z^r = \frac{(1 - z)^2}{1 - 2Yz + z^2}.$$

Extracting coefficients against $\sum U_r(Y) z^r = (1 - 2Yz + z^2)^{-1}$ gives $\Phi_r = U_r - 2U_{r-1} + U_{r-2}$, and the three-term recurrence $U_r + U_{r-2} = 2YU_{r-1}$ collapses this to $(2Y - 2)U_{r-1}(Y)$. $\square$

In the formalization the analytic Step 2 is replaced by an equivalent finite argument: the closed form is shown to satisfy (2) directly (a second-difference induction), and (2) with $\Phi_0 = 1$ determines Φ over a characteristic-zero domain.

3 Graphs of first Betti number one

The proof above uses cycles in exactly one place: gauge-fixing a spanning path leaves a *single* permutation, because C_n has $|E| - |V| + 1 = 1$. The argument therefore applies verbatim to any connected G with $b_1(G) = |E| - |V| + 1 = 1$, a cycle with trees attached, and we record what it gives.

By [2] the expected characteristic polynomial of a random r -lift of *any* graph factors as $\chi_G \cdot \mu_{r-1,G}$; the content below is therefore not that factorization but a closed form for the factor $\mu_{d,G}$ itself. Question 6.5 of [2] asks which parts of the theory of the matching polynomial $\mu_{1,G}$ generalize to $\mu_{d,G}$; the following answers it for $b_1(G) = 1$.

Let C be the unique cycle of G and let $G - V(C)$ be the forest obtained by deleting its vertices. Write μ_H for the ordinary matching polynomial of a graph H , with $\mu_\emptyset = 1$. Define

$$V_0 = 1, \quad V_1 = \mu_G, \quad V_d = \mu_G V_{d-1} - \mu_{G-V(C)}^2 V_{d-2}. \quad (3)$$

Theorem 2. *For every connected G with $b_1(G) = 1$ and every $d \geq 0$, $\mu_{d,G} = V_d$.*

Theorem 2 is proved below, through the twisted characteristic polynomial; it is not formalized.

For $G = C_n$ the deleted forest is empty, so $\mu_{G-V(C)} = 1$ and (3) is the recurrence for $U_d(\mu_{C_n}/2) = U_d(T_n(x/2))$: Theorem 1 is the case $\mu_{G-V(C)} = 1$. In general the attached trees enter only through $\mu_{G-V(C)}$; for the triangle with one pendant edge, $\mu_G = x^4 - 4x^2 + 1$ and $\mu_{G-V(C)} = x$.

The proof goes through the twisted characteristic polynomial. Let $e = uv$ be the unique non-tree edge, let $A_G(z)$ be the adjacency matrix of G with the (u, v) entry replaced by z and the (v, u) entry by z^{-1} , and set $F(x, z) = \det(xI - A_G(z))$.

Lemma 3. $F(x, z) = \mu_G(x) - \mu_{G-V(C)}(x)(z + z^{-1})$.

Proof. By the Sachs coefficient theorem $\det(xI - A)$ expands over elementary subgraphs, that is disjoint unions of edges and cycles. Since a permutation uses each of the two twisted entries at most once and $A_G(z)^\top = A_G(z^{-1})$, the determinant is of the form $P(x) + Q(x)(z + z^{-1})$. Replacing z by ± 1 and averaging kills exactly the elementary subgraphs whose cycle traverses e ; as G has the single cycle C , those are the subgraphs containing C , so $P = \mu_G$, and the surviving terms contribute $Q = -\mu_{G-V(C)}$, a cycle component carrying the Sachs weight -2 . $\square$

Proof of Theorem 2. Gauge-fix a spanning tree: the r -lift is determined by one permutation $\sigma \in S_r$ on e , and its components are indexed by the cycles of σ , the component of an ℓ -cycle being the ℓ -fold cyclic cover G_ℓ . Decomposing the $\mathbb{Z}/\ell$ action into characters gives the classical factorization $\chi_{G_\ell}(x) = \prod_{j=0}^{\ell-1} F(x, \omega^j)$ with $\omega = e^{2\pi i/\ell}$ [9, 10]. With P, Q as in Lemma 3 and $y = -P/(2Q)$,

$$\chi_{G_\ell} = \prod_{j=0}^{\ell-1} \left(P + 2Q \cos \frac{2\pi j}{\ell} \right) = (-Q)^\ell (2T_\ell(y) - 2),$$

so by the exponential formula (2), with $w = -Qz$,

$$\sum_{r \geq 0} \Phi_{G,r} z^r = \exp \left(\sum_{\ell \geq 1} (2T_\ell(y) - 2) \frac{w^\ell}{\ell} \right) = \frac{(1-w)^2}{1-2yw+w^2},$$

and extracting coefficients as in Section 2 gives $\Phi_{G,r} = \chi_G \cdot (-Q)^{r-1} U_{r-1}(y)$. Comparing with $\Phi_{G,r} = \chi_G \cdot \mu_{r-1,G}$ [2] yields $\mu_{d,G} = (-Q)^d U_d(y)$, which satisfies (3). $\square$

Since $U_d(y) = 2^d \prod_{k=1}^d (y - \cos \frac{k\pi}{d+1})$, the theorem can be restated as a product, $\mu_{d,G}(x) = \prod_{k=1}^d F(x, e^{ik\pi/(d+1)})$, whence:

This is the real-rootedness theorem of [2] for this family, with the roots identified. The localization itself, that a vanishing V_d forces $|\mu_G| \leq 2|\mu_{G-V(C)}|$, so that a root can never sit in a spectral gap, is machine-checked in Lean 4 as `cV.root_mem_band`, resting on `cU.ne_zero_of_one_le_abs`: the Chebyshev polynomial U_d has no root of absolute value at least one.

The roots can be located much more precisely than that. Since $b_1(G) = 1$ the group $\pi_1(G)$ is $\mathbb{Z}$, so the universal cover T of G is the $\mathbb{Z}$ -cover, and T is a periodic decorated chain whose adjacency operator decomposes by Floquet–Bloch into the fibres $A_G(e^{i\theta})$. Hence

$$\text{spec}(T) = \bigcup_{\theta} \text{spec } A_G(e^{i\theta}) = \left\{ x : \exists s \in [-1, 1], \mu_G(x) = 2s \mu_{G-V(C)}(x) \right\},$$

a finite union of bands, together with a flat band at each common zero of μ_G and $\mu_{G-V(C)}$, where $F(x, \cdot)$ vanishes identically.

Corollary 4. *For $b_1(G) = 1$ the roots of $\mu_{d,G}$ are exactly the solutions of $\mu_G(x) = 2 \cos(\frac{k\pi}{d+1}) \mu_{G-V(C)}(x)$ for $1 \leq k \leq d$. In particular every root of every $\mu_{d,G}$ lies in $\text{spec}(T)$, and as $d \rightarrow \infty$ the roots equidistribute on $\text{spec}(T)$ with respect to its density of states.*

Proof. Immediate from Theorem 2 in the form $\mu_{d,G} = \prod_k F(x, e^{ik\pi/(d+1)})$ and $F(x, e^{i\theta}) = \mu_G - 2 \cos \theta \mu_{G-V(C)}$. The equidistribution is the arcsine law for $\cos(k\pi/(d+1))$ transported through the fibre map. $\square$

Hall–Puder–Sawin place the roots of $\mu_{d,G}$ in the Ramanujan *interval* $[-\rho, \rho]$, ρ being the spectral radius of T . Corollary 4 places them in $\text{spec}(T)$, which is a union of bands and is a proper subset of that interval whenever T has a spectral gap: no root of any $\mu_{d,G}$ can ever enter a gap. The two statements coincide exactly when the band structure is trivial, and that is precisely the case of a cycle, where $\mu_{G-V(C)} = 1$ and $\text{spec}(T) = [-2, 2]$ is the whole interval. Attaching even a single pendant edge opens gaps, and they are not small (computed band structures, proportion of $[-\rho, \rho]$ that is a gap):

G	ρ	gap proportion
C_n	2	0%
triangle + pendant	2.170	36.5%
C_4 + pendant	2.136	45.9%
triangle + P_2	2.214	51.3%
C_4 + P_3	2.189	54.1%
triangle + $K_{1,3}$	2.514	65.9%

So the cycle is not merely the easiest case of Theorem 2: it is the degenerate one, the unique member of the family whose spectrum has no gaps and for which the Ramanujan interval is attained. Any cycle with a tree attached exhibits a band structure that no cycle can display.

The boundary is sharp. For $b_1(G) = b$ the lift is determined by b permutations, that is by an action of the free group F_b on $[r]$, whose orbits are the components of the lift. The exponential formula still applies, in the form $N_r = \sum_{\ell} \binom{r-1}{\ell-1} C_{\ell} N_{r-\ell}$ with $N_r = (r!)^b \Phi_{G,r}$ and C_{ℓ} the sum of χ over the transitive b -tuples on $[\ell]$; we have verified this for $b \leq 3$. What fails is the atom. For $b = 1$ every transitive tuple gives the *same* cover G_{ℓ} , so $C_{\ell} = (\ell - 1)! \chi_{G_{\ell}}$: one graph per ℓ , with $\chi_{G_{\ell}}$ Chebyshev in ℓ . For $b \geq 2$ the transitive tuples number $(\ell - 1)!$ times the count of index- ℓ subgroups of F_b and spread over many non-isomorphic covers with distinct spectra, so C_{ℓ} carries no such structure. Accordingly Theorem 2 is not merely unproved for $b \geq 2$: it is false there already at $d = 1$, as we checked for the theta graph, the bowtie, K_4 , $K_{2,3}$ and two triangles joined by an edge.

Theorem 2 was verified by exact integer computation against the definition of $\mu_{d,G}$ as an average of matching polynomials over all d -covers, and against brute-forced lift averages, on thirteen named unicyclic graphs for $r \leq 6$ and on a random sample of thirty-four for $r \leq 5$; Lemma 3 was checked on eleven.

4 A conjecture for all graphs

Corollary 4 is proved by Floquet theory, which is available only because $\pi_1(G) = \mathbb{Z}$. Its *statement*, however, makes sense for every graph, and we know one case of it: at $b_1(G) = 1$ it is Corollary 4, for every d . We therefore propose:

Conjecture 5. *For every finite graph G and every $d \geq 1$, all roots of $\mu_{d,G}$ lie in $\text{spec}(T)$, the spectrum of the universal cover of G .*

Theorem 6 (Hall [12]). *Conjecture 5 is false, already at $d = 1$ and already for simple connected loopless bipartite graphs.*

The counterexample and its certificate are entirely due to Chris Hall. We record it here because it settles the central question of this paper, and because everything else in the note has to be read in its light. §5 states his construction, our independent verification of it, and what we can add.

The biregular case of Conjecture 5 at $d = 1$ is not new. It is Problem 1 of Song, Fan and Miao [5], posed in 2023: they ask whether the matching polynomial of the incidence graph B_H of a (d, r) -regular hypergraph has no root μ with $0 < |\mu| < |\sqrt{d-1} - \sqrt{r-1}|$, and every (d, r) -biregular bipartite graph is such an incidence graph. Their Theorem 1.8 attaches a consequence: an affirmative answer would give every (d, r) -regular hypergraph a left-sided Ramanujan 2-covering. The only bound in that direction we know of is their Lemma 5.5, which goes the other way, $0 < \mu_{\tau}(B_H) \leq \max\{\sqrt{d}, \sqrt{r}\}$. We are not aware of any proof, partial result or counterexample. Conjecture 5 extends their problem in two directions, to arbitrary graphs and to all d .

Question 6.3 of [2] asks the corresponding question for graph eigenvalues, and leaves it open: it distinguishes graphs whose nontrivial spectrum lies in the *interval* $[-\rho, \rho]$ from those whose nontrivial spectrum lies in $\text{spec}(T)$, observes that the two notions separate exactly when T has a gap, and gives the biregular tree as the example. Conjecture 5 is the matching-polynomial form of that distinction.

The conjecture strengthens the root-localization theorem of [2], which places the roots in the interval $[-\rho, \rho]$ with ρ the spectral radius of T ; the conjecture places them in $\text{spec}(T)$

itself. The two agree whenever T has no spectral gap, in particular for regular G , where T is a regular tree and $\text{spec}(T) = [-2\sqrt{q}, 2\sqrt{q}]$ is an interval. They differ exactly when T has a gap, and the smallest examples are biregular: for $K_{2,3}$ the universal cover is the $(3, 2)$ -biregular tree, with spectrum $\{0\} \cup \pm[\sqrt{2}-1, \sqrt{2}+1]$ and hence a gap $(0, \sqrt{2}-1)$ inside the Ramanujan interval. We have tested the conjecture on the biregular family, where the universal cover is an (a, b) -biregular tree with spectrum $\{0\} \cup \pm[|s-t|, s+t]$, $s = \sqrt{a-1}$, $t = \sqrt{b-1}$, so the gap is $(0, |s-t|)$. In each case $\mu_{d,G}$ was computed by averaging matching polynomials over all d -covers. No root ever lies in the gap; we also record the smallest nonzero $|\text{root}|$ as a multiple of the gap edge, which measures how close the bound comes to being attained.

G	b_1	type	gap edge	d	$\min \text{root} /\text{gap edge}$
$K_{2,3}$	2	$(3, 2)$	0.4142	1, 2, 3, 4	2.72, 2.19, 1.94, 1.78
$K_{2,4}$	3	$(4, 2)$	0.7321	1, 2, 3	1.93, 1.65, 1.52
$K_{2,5}$	4	$(5, 2)$	1.0000	1, 2, 3	1.66, 1.46, 1.37
$K_{3,4}$	6	$(4, 3)$	0.3178	1, 2	3.04, 2.31
$K_{3,5}$	8	$(5, 3)$	0.5858	1, 2	2.10, 1.73
subdivision of K_4	3	$(3, 2)$	0.4142	1, 2	1.97, 1.67

All of these have $b_1 \geq 2$, so none is covered by Corollary 4. It is worth recording that even $d = 1$ is not formal. There $\mu_{1,G}$ is the ordinary matching polynomial, whose roots are eigenvalues of the path tree of G (Godsil), and the path tree embeds in T ; but eigenvalues of a finite subtree of T need not lie in $\text{spec}(T)$ when T has a gap. For the $(3, 2)$ -biregular tree, whose gap is $(0, \sqrt{2}-1)$, the path P_8 is a subtree with eigenvalue $2\cos(4\pi/9) = 0.3473$ inside that gap. So the $d = 1$ case does not follow from the path-tree argument. It is nevertheless a theorem for an infinite family.

5 Hall's counterexample

Everything in this section is due to Chris Hall, who constructed the graph, found the certificate, and wrote the verification note [12]. Our contribution is the independent check, the formalization of the algebra. The two-parameter family containing his graph is in [18].

5.1 The graph and its matching polynomial

Let G consist of a central vertex c ; for $1 \leq i \leq 5$ a copy of $K_{2,5}$ with degree-five vertices v_i, w_i and middle vertices $u_{i,1}, \dots, u_{i,5}$; a pendant leaf ℓ_i at each w_i ; and the edges cv_i . It is simple, connected, loopless and bipartite, with 41 vertices, 60 edges and $b_1(G) = 20$.

Writing H for one branch after deleting c , the vertex-deletion recurrence gives $\mu_{H-v} = x^5(x^2 - 6)$ and $\mu_H = x^4(x^4 - 11x^2 + 25)$, and then

$$\mu_G(x) = \mu_H(x)^4(x\mu_H(x) - 5\mu_{H-v}(x)) = x^{21}(x^4 - 11x^2 + 25)^4(x^2 - 5)(x^2 - 11), \quad (4)$$

so $\sqrt{5}$ is a simple root. Identity (4), the simplicity of the root, and the value of the cofactor are machine-checked in `RamaLean/HallCounterexample.lean` as an identity in $\mathbb{Z}[X]$ (`muG_eq`, `muG_root_sqrt5`, `muG_root_simple`).

5.2 The spectral exclusion

Angel, Friedman and Hoory [13] characterise invertibility of $A_T - \lambda I$ by a finite nonzero *ratio system*: an assignment r_e to directed edges with $\lambda = 1/r_e + \sum_{e \rightarrow f} r_f$, whose decay matrix $\mathcal{R}_{e,f} = |r_f|^2$ on non-backtracking pairs has spectral radius below one. Hall exhibits one at $\lambda = \sqrt{5}$ with ratios in $\mathbb{Q}(\sqrt{5}, \sqrt{41})$; its 120-state decay matrix has ten transient states at the leaves, and the recurrent block of 110 reduces by symmetry to a 6×6 orbit quotient K with $x = (20, 121, 47, 33, 28, 16)^\top$ satisfying $Kx < x$ componentwise, so $\rho(K) < 1$ by Perron–Frobenius. Hence $\sqrt{5} \notin \text{spec}(A_T)$, which with (4) proves Theorem 6. We rebuilt every object from the graph rather than copying the displayed matrices (`code/hall_certificate.py`) and machine-checked the algebra (`RamaLean/HallCounterexample.lean`).

5.3 The Divisibility Lemma

The counterexamples are all instances of one algebraic fact, which we state once.

Lemma 7 (Divisibility). *Let G be a finite graph, $S \subseteq V(G)$ with $|S| = k$, and suppose $G - S$ has p components, each inducing a copy of the same graph B . Write $A = \mu_B$. Then*

$$A^{p-k} \mid \mu_G.$$

Proof. Expand $\mu_G = \sum_M (-1)^{|M|} x^{n-2|M|}$ over matchings M of G and group the terms by

$$J(M) = \{i : \text{some edge of } M \text{ joins } S \text{ to the } i\text{-th component}\}.$$

A matching uses each vertex of S at most once, so distinct components in $J(M)$ are entered through distinct vertices of S ; choosing one such vertex per component is an injection $J(M) \hookrightarrow S$, whence $|J(M)| \leq k$. For $i \notin J(M)$ the restriction of M to the i -th component is an arbitrary matching of B , and summing over those choices independently contributes a factor $A^{p-|J(M)|}$. Each term of the resulting sum is therefore divisible by A^{p-k} , and hence so is μ_G . $\square$

At $k = 2$ this is the divisor A^{p-2} of the two-cut identity. The exponent is positive exactly when $p > k$, which is why every construction below needs at least $k + 1$ branches. The counting step and the algebraic step are machine-checked separately (`RamaLean/DivisibilityLemma.lean: card_branches_le_sep, dvd_sum_of_pow_terms, matching_expansion_dvd`); the matching expansion itself is not, Mathlib carrying no matching polynomial.

5.4 Why the branch mechanism cannot produce a counterexample

The natural repairs are local hypotheses on G . Minimum degree two fails and bounded maximum degree fails, so one is led to:

Conjecture 8 (D3). *Every finite graph of minimum degree at least three satisfies $\text{Zeros}(\mu_G) \subseteq \text{spec}(T_G)$.*

Conjecture 9 (C2). *Every finite 3-connected graph satisfies $\text{Zeros}(\mu_G) \subseteq \text{spec}(T_G)$.*

Both are open. We record here why the construction of §5.3, which is the obvious way to attack them, cannot settle either: the configuration that makes Lemma 7 produce a root of μ_G is the same configuration that puts that root *inside* $\text{spec}(T_G)$.

Banks, Garza-Vargas and Mukherjee [14] characterise the eigenvalues of the universal cover: θ is an eigenvalue of T_G if and only if G has a θ -Aomoto subset, that is a set $S \subseteq V(G)$ with $G[S]$ a forest, θ an eigenvalue of every component of $G[S]$, and

$$|\partial_G S| < \text{cc}(G[S]).$$

Li, Magee, Sabri and Thomas [15] prove the corresponding statement for the maximal abelian cover, in the form that θ is an eigenvalue of G^{ab} if and only if θ is a zero of $\mu_{G \setminus \Gamma}$ for every 2-regular subgraph Γ , and Spier [16] proves that T_G and G^{ab} have the same eigenvalues.

Proposition 10 (the branch obstruction). *Let $\theta \in \mathbb{R}$ and let $S \subseteq V(G)$ with $|S| = k$. If at least $k + 1$ of the components of $G - S$ induce forests having θ as an eigenvalue, then θ is an eigenvalue of T_G , and in particular $\theta \in \text{spec}(T_G)$.*

Proof. Let U be the union of $j \geq k + 1$ such components. Then $G[U]$ is a disjoint union of forests, hence a forest, with $\text{cc}(G[U]) = j$ and θ an eigenvalue of every component. Every vertex outside U adjacent to U lies in S , so $|\partial_G U| \leq k < j$. Thus U is a θ -Aomoto subset and [14] applies. $\square$

The components need not be isomorphic to one another, and need not be all of $G - S$; only their number relative to $|S|$ matters. Taking them to be p copies of a single tree B with θ a root of μ_B recovers the case that concerns us:

The hypotheses of Proposition 10 are then exactly those of Lemma 7 together with $p > k$, which is what the lemma needs for the exponent $p - k$ to be positive. So whenever the branch mechanism forces a root of μ_G , it forces that root into $\text{spec}(T_G)$. The two conditions are the same condition:

$$p > k \quad \text{is} \quad |\partial_G S| < \text{cc}(G[S]).$$

A shared separator keeps the boundary at k while the number of components grows with p , and the Aomoto inequality is then automatic.

This is why Hall's construction is subtler than it looks, and it explains the role of the pendant leaves, which [18] records as essential without a mechanism. His branches are not trees, so the obstruction does not apply to them directly; the natural forest inside a branch is the star $K_{1,5}$ at w_i , whose boundary is the pair $\{v_i, \ell_i\}$. With five branches that gives $\text{cc} = 5$ and $|\partial| = 10$, comfortably on the wrong side of the Aomoto inequality. Deleting the pendant leaves brings the boundary down to 5, still not below 5, and it also destroys the root: the leafless graph has $\mu = x^{16}(x^4 - 15x^2 + 45)(x^4 - 10x^2 + 20)^4$, with no factor $x^2 - 5$. The leaves therefore do two jobs at once, creating the root and doubling the boundary of the star, and both are needed (`code/aomoto_obstruction.py`). The general form is checked there on non-isomorphic branches as well, a hub joined to both ends of a P_3 and of a P_7 , where $\sqrt{2}$ is an eigenvalue of each path and the two components already outnumber the single boundary vertex.

The moral for a repair is that it must forbid the *private* boundary structure that Hall uses, not the ambient degrees. Every counterexample must avoid having a θ -Aomoto subset at its offending root, and that is a condition on how branches attach, not on how large their degrees are.

5.5 A caveat on all the negative evidence

The searches reported here and in [18] filter the detected gaps of $\text{spec}(T)$ by a minimum width before looking for roots inside them, and skip a graph when no gap survives. The threshold used throughout is 0.05. Hall’s own counterexample does not survive it: $\text{spec}(T)$ for his graph has a gap

$$(2.220, 2.260), \quad \text{width } 0.040, \quad \sqrt{5} = 2.23607 \dots \in (2.220, 2.260),$$

and $0.040 < 0.05$, so his graph is scored as having no gap and reported clean (`code/gapfilter_blindspot.py`). The detector finds the gap correctly; the threshold throws it away.

Every negative verdict in this development must therefore be read as *no violation was found in a gap wider than 0.05*, which is much weaker than it sounds, and which excludes the one counterexample known to exist. The counterexamples of the two-cut family occupy widths 0.066 to 0.148, which is presumably where the threshold came from; Hall’s is narrower than all of them, and so is the regime the phenomenon actually inhabits. We record this rather than quietly re-running, because the published searches were carried out at the old setting and their stated scale conveys a confidence the instrument did not have.

5.6 What this leaves

Conjecture 5 is false, by Hall’s counterexample. Conjectures 8 and 9 are open, and Proposition 10 says that the branch mechanism cannot settle them in the negative: it manufactures eigenvalues of the cover, not counterexamples.

That is a sharper statement than it may look. Any counterexample must have its offending root avoid every θ -Aomoto subset, so a search should be organised around the Aomoto inequality rather than around degrees or connectivity. Hall’s graph satisfies this by giving each branch a private boundary of two vertices; we know of no systematic way to do that at minimum degree three.

The biregular case, which survives the refutation of Conjecture 5 and is where the phenomenon lives, is [17].

6 Related work and formalization

The d -matching polynomial and the interval $[-\rho, \rho]$ are due to Hall, Puder and Sawin [2], whose Question 6.3 asks for exactly the strengthening considered here; Heilmann and Lieb [6] give the classical interval, Godsil [7, 8] the path-tree method, and Angel, Friedman and Hoory [13] the ratio criterion the certificates use. The cycle identity behind Theorem 1 was proved combinatorially by Cochran, Groothuis, Herring, Rohatgi and Stucky [3]; our route is independent and elementary. Song, Fan and Miao [5] pose the biregular case, which is [17].

The Lean 4/Mathlib sources are in `RamaLean/` of the accompanying repository and carry no `sorry` and no custom axioms: the closed form at first Betti number one, the counterexample’s algebra, and both halves of Lemma 7. The standing gap is that Mathlib has no matching polynomial, so the expansion in Lemma 7 and the Angel–Friedman–Hoory criterion are cited rather than derived.

References

- [1] C. Godsil, I. Gutman, *On the matching polynomial of a graph*, in: Algebraic Methods in Graph Theory, 1981.
- [2] C. Hall, D. Puder, W. Sawin, *Ramanujan coverings of graphs*, Adv. Math. **323** (2018), 367–410.
- [3] G. Cochran, C. Groothuis, A. Herring, R. Rohatgi, E. Stucky, *A new (combinatorial) proof of the commutativity of matching polynomials for cycles*, arXiv:1810.05889 (2018). Preprint; no journal reference or DOI as of August 2026.
- [4] J. Wan, W. Wang, A. Mohammadian, *On the matching polynomials of graphs with small number of cycles*, arXiv:2103.10580 (2021).
- [5] Y.-M. Song, Y.-Z. Fan, Z. Miao, *Hypergraph coverings and Ramanujan hypergraphs*, arXiv:2310.01771 (2023).
- [6] O. J. Heilmann, E. H. Lieb, *Theory of monomer-dimer systems*, Comm. Math. Phys. **25** (1972) 190–232.
- [7] C. D. Godsil, *Matchings and walks in graphs*, J. Graph Theory **5** (1981), 285–297.
- [8] C. D. Godsil, *Algebraic Combinatorics*, Chapman and Hall, New York, 1993, Theorem 6.1.1 and §5.6.
- [9] H. Mizuno, I. Sato, *Characteristic polynomials of some graph coverings*, Discrete Math. **142** (1995) 295–298.
- [10] C. Dalfó, M. A. Fiol, M. Miller, J. Ryan, J. Širáň, *An algebraic approach to lifts of digraphs*, Discrete Appl. Math. **269** (2019) 68–76.
- [11] A. Marcus, D. Spielman, N. Srivastava, *Interlacing families I: Bipartite Ramanujan graphs of all degrees*, Ann. of Math. **182** (2015), 307–325.
- [12] C. Hall, *A simple bipartite counterexample to Conjecture 10: spectral localization of the ordinary matching polynomial*, verification note, personal communication, August 2026.
- [13] O. Angel, J. Friedman, S. Hoory, *The non-backtracking spectrum of the universal cover of a graph*, Trans. Amer. Math. Soc. **367** (2015), 4287–4318; arXiv:0712.0192.
- [14] J. Banks, J. Garza-Vargas, S. Mukherjee, *Point spectrum of the universal cover of a graph*, as cited in [16], Cor. 3.4.
- [15] W. Li, M. Magee, M. Sabri, D. Thomas, *Eigenvalues of maximal abelian covers*, arXiv:2508.17332 (2025).
- [16] T. J. Spier, *Eigenvalues of universal covers and the matching polynomial*, arXiv:2510.05041 (2025).
- [17] V. Bonfiori, *The biregular case of a spectral-inclusion conjecture: the weighted plane class and its obstructions*, companion paper, 2026.

- [18] V. Bonfiori, *Roots of d -matching polynomials: supplementary material*, companion note, 2026.
- [19] Z. Xu, *Note on mixed characteristic polynomials*, personal communication, August 2026.
- [20] S. Mohanty, R. O'Donnell, *X-Ramanujan graphs*, arXiv:1904.03500v2 (2019).