

SUPPORT VECTOR MACHINES





LAGRANGE MULTIPLIERS



EQUALITY CONSTRAINTS

Problem.

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & h_1(x) = 0, \dots, h_l(x) = 0\end{array}$$

Lagrange multipliers.

1. Write down the Lagrangian.

$$L(x, \lambda) = f(x) + \lambda_1 h_1(x) + \dots + \lambda_l h_l(x)$$

2. Solve for critical points x, λ .

$$\nabla_x L(x, \lambda) = 0, \quad h_1(x) = 0, \dots, h_l(x) = 0$$

3. Pick critical point which gives global minimum.



EXAMPLE

$$\begin{array}{ll}\text{minimize} & f(x) = n_1 \log x_1 + \cdots + n_d \log x_d \\ \text{subject to} & h(x) = x_1 + \cdots + x_d - 1 = 0\end{array}$$

Lagrangian

$$L(x, \lambda) = n_1 \log x_1 + \cdots + n_d \log x_d + \lambda(x_1 + \cdots + x_d - 1)$$

Critical points

$$\begin{array}{ll}0 = x_1 + \cdots + x_d - 1 & \\ 0 = n_i/x_i + \lambda & \Rightarrow \quad \begin{array}{l} (-\lambda) = n_1 + \cdots + n_d \\ x_i = n_i/(-\lambda) \end{array}\end{array}$$



TWO-PLAYER GAME

$$L(x, \lambda) = f(x) + \lambda_1 h_1(x) + \cdots + \lambda_l h_l(x)$$

Rules.

- You get to choose the value of x .
Your goal is to minimize $L(x, \lambda)$.
- Your adversary gets to choose the value of λ .
His goal is to maximize $L(x, \lambda)$.



PRIMAL GAME

$$L(x, \lambda) = f(x) + \lambda_1 h_1(x) + \cdots + \lambda_l h_l(x)$$

Primal Game. You go first.

Your Strategy.

- Ensure that $h_1(x) = 0, \dots, h_l(x) = 0$.
- Find x that minimizes $f(x)$.

Final Score. $p^* = \min_x \max_{\lambda} L(x, \lambda)$



DUAL GAME

$$L(x, \lambda) = f(x) + \lambda_1 h_1(x) + \cdots + \lambda_l h_l(x)$$

Dual Game. You go second.

Adversary's Strategy.

- For each λ , compute $\ell(\lambda) = \min_x L(x, \lambda)$
- Find λ that maximizes $\ell(\lambda)$.

Final Score. $d^* = \max_{\lambda} \min_x L(x, \lambda)$



MAX-MIN INEQUALITY

Primal. $p^* = \min_x \max_{\lambda} L(x, \lambda)$

Dual. $d^* = \max_{\lambda} \min_x L(x, \lambda)$

$$\begin{aligned} p^* &= \min_x \max_{\lambda} L(x, \lambda) \\ &\geq \max_{\lambda} \min_x L(x, \lambda) = d^* \end{aligned}$$

Challenge. Can you prove this?



INEQUALITY CONSTRAINTS

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & g_1(x) \leq 0, \dots, g_m(x) \leq 0\end{array}$$

Lagrangian.

$$L(x, \alpha) = f(x) + \alpha_1 g_1(x) + \dots + \alpha_m g_m(x)$$

Solve for x, α satisfying

1. $\nabla_x L(x, \alpha) = 0$
2. $g_1(x) \leq 0, \dots, g_m(x) \leq 0$
3. $\alpha_1 \geq 0, \dots, \alpha_m \geq 0$
4. $\alpha_1 g_1(x) = 0, \dots, \alpha_m g_m(x) = 0$

Karush-Kuhn-Tucker (KKT) Conditions

Complementary Slackness



PRIMAL-DUAL PROBLEMS

Primal Problem.

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & g_1(x) \leq 0, \dots, g_m(x) \leq 0\end{array}$$

Dual Problem.

$$\begin{array}{ll}\text{maximize} & \ell(\alpha) \\ \text{subject to} & \alpha_1 \geq 0, \dots, \alpha_m \geq 0\end{array} \quad \text{where } \ell(\alpha) = \min_{x \in \mathbb{R}^d} L(x, \alpha)$$

Box constraints are
easier to work with!







MAXIMUM MARGINS



COMPUTING THE MARGIN



COMPUTING THE MARGIN



MAXIMUM MARGIN

Our goal is to

maximize $1/\|\theta\|$

subject to $y(\theta \cdot x + \theta_0) \geq 1$ for all data (x, y)

Or equivalently,

minimize $\frac{1}{2} \|\theta\|^2$

subject to $y(\theta \cdot x + \theta_0) \geq 1$ for all data (x, y)



LAGRANGIAN

$$\begin{array}{ll} \text{minimize} & \frac{1}{2} \|\theta\|^2 \\ \text{subject to} & y(\theta \cdot x) \geq 1 \text{ for all data } (x, y) \end{array}$$

Drop θ_0 for now

Lagrangian. $L(\theta, \alpha) = \frac{1}{2} \|\theta\|^2 + \sum_{(x,y)} \alpha_{x,y} (1 - y(\theta \cdot x))$

To find $\ell(\alpha) = \min_{\theta} L(\theta, \alpha)$, we solve

$$0 = \nabla_{\theta} L(\theta, \alpha) = \theta - \sum_{(x,y)} \alpha_{x,y} yx$$

to get $\theta = \sum_{(x,y)} \alpha_{x,y} yx$. Substituting into $L(\theta, \alpha)$ gives

$$\ell(\alpha) = \sum_{(x,y)} \alpha_{x,y} - \frac{1}{2} \sum_{(x,y)} \sum_{(x',y')} \alpha_{x,y} \alpha_{x',y'} y y' (x \cdot x').$$



PRIMAL-DUAL

Primal.

$$\begin{array}{ll}\text{minimize} & \frac{1}{2} \|\theta\|^2 \\ \text{subject to} & y(\theta^\top x) \geq 1 \text{ for all data } (x, y)\end{array}$$

Dual.

$$\begin{array}{ll}\text{maximize} & \sum_{(x,y)} \alpha_{x,y} - \frac{1}{2} \sum_{(x,y)} \sum_{(x',y')} \alpha_{x,y} \alpha_{x',y'} y y' (x \cdot x') \\ \text{subject to} & \alpha_{x,y} \geq 0 \text{ for all } (x, y)\end{array}$$

It can be shown that the primal and dual problems are equivalent (*strong duality*).



SUPPORT VECTORS

Complementary Slackness.

$$\hat{\alpha}_{x,y} > 0: \quad y(\hat{\theta}^\top x) = 1$$


Support Vectors

$$\hat{\alpha}_{x,y} = 0: \quad y(\hat{\theta}^\top x) > 1$$

Non-Support Vectors

Sparsity

Since very few data points are support vectors, **most of the $\hat{\alpha}_{x,y}$ will be zero.**



KERNEL TRICK

Learning.

$$\ell(\alpha) = \sum_{(x,y)} \alpha_{x,y} - \frac{1}{2} \sum_{(x,y)} \sum_{(x',y')} \alpha_{x,y} \alpha_{x',y'} y y' (x \cdot x')$$

Prediction.

$$h(x; \theta) = \text{sign}(\theta \cdot x) = \text{sign} \left(\sum_{(x',y')} \alpha_{x',y'} y' (x \cdot x') \right)$$

For the dual, we don't need the feature vectors x, x' .
Knowing just the dot products $(x \cdot x')$ is enough.





DISCUSSION



SVM WITH OFFSET

Primal.

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|\theta\|^2 \\ &\text{subject to} && y(\theta^\top x + \theta_0) \geq 1 \text{ for all data } (x, y) \end{aligned}$$

Dual.

$$\begin{aligned} &\text{maximize} && \sum_{(x,y)} \alpha_{x,y} - \frac{1}{2} \sum_{(x,y)} \sum_{(x',y')} \alpha_{x,y} \alpha_{x',y'} y y' (x \cdot x') \\ &\text{subject to} && \alpha_{x,y} \geq 0 \text{ for all } (x, y) \\ &&& \sum_{(x,y)} \alpha_{x,y} y = 0 \end{aligned}$$

Parameters.

$$\begin{aligned} \hat{\theta} &= \sum_{(x,y)} \alpha_{x,y} y x \\ \hat{\theta}_0 &= y - \hat{\theta} \cdot x \quad \text{where } (x, y) \text{ is a support vector} \end{aligned}$$



SVM WITH ERRORS

Primal.

$$\begin{array}{ll} \text{minimize} & \frac{\lambda}{2} \|\theta\|^2 + \frac{1}{n} \sum_{(x,y)} \xi_{x,y} \\ \text{subject to} & y(\theta^\top x + \theta_0) \geq 1 - \xi_{x,y} \quad \text{for all data } (x, y) \\ & \xi_{x,y} \geq 0 \quad \text{for all data } (x, y) \end{array}$$

Slack variables allow constraints to be violated for a cost.

Equivalent Primal.

$$\text{minimize} \quad \frac{\lambda}{2} \|\theta\|^2 + \frac{1}{n} \sum_{(x,y)} \text{Loss}_H(y(\theta \cdot x + \theta_0))$$


Hinge-loss classifier with regularization!




SVM WITH ERRORS

Linearly Separable.

$\lambda = 1$ ($C = 1$)




$\lambda = 100$ ($C = 0.01$)




SVM WITH ERRORS

Not Linearly Separable.

$\lambda = 0.1$ ($C = 10$)



$\lambda = 100$ ($C = 0.01$)



SVM WITH ERRORS

Dual.

$$\begin{aligned} &\text{maximize} && \sum_{(x,y)} \alpha_{x,y} - \frac{1}{2} \sum_{(x,y)} \sum_{(x',y')} \alpha_{x,y} \alpha_{x',y'} y y' (x \cdot x') \\ &\text{subject to} && 1/\lambda \geq \alpha_{x,y} \geq 0 \text{ for all } (x,y) \\ &&& \sum_{(x,y)} \alpha_{x,y} y = 0 \end{aligned}$$

Putting limits on what the adversary can do.



SUMMARY

- Lagrange Multipliers
 - Lagrangian
 - Primal-Dual Problems
 - Inequality Constraints
 - Complementary Slackness
- Regularization
 - Slack Variables
 - Regularized Hinge Loss
 - Bounded Multipliers
- Support Vector Machines
 - Maximum Margins
 - Dual Problem
 - Support Vectors
 - Kernel Trick

