easyformat, tests

$$\int_{a}^{b} \mathrm{d}x = b - a$$

Evert Provoost

28th May 2017

Contents

 $1 \quad \text{Main test}$ $2 \quad \textit{Math, } _u_n d_e r_s c_o r_e s \text{ and } \^c^i r^c o^n f^l e^x e^s \dots$ $1 \quad \text{3} \quad \text{Mixing } \textit{styles}$ $1 \quad \text{1} \quad \text{2} \quad \text{3} \quad \text{4} \quad \text$

1 Main test

These two should be identical and definitely have the exact same length:

| Lorem ipsum totalem DA givea zin doram zettim, liefkan. | | Lorem ipsum totalem DA givea zin doram zettim, liefkan. |

2 *Math*, $_{-}u_{n}d_{e}r_{s}c_{o}r_{e}s$ and $_{\hat{-}}c^{i}r^{c}o^{n}f^{l}e^{x}e^{s}...$

This (\uparrow Easy Formatting inside subtitles) is possible without some extra code since v1.1.0 (2017/04/09) and having a $\hat{}$ in mathmode since v1.3.0 (2017/05/28).

$$\int_a^b x \, \mathrm{d}x = \frac{b^2 - a^2}{2}$$

3 Mixing styles

YEAH, yeah?, YEAH!!! \leftarrow notice how we go from SMALLCAPS to *italics* and back to SMALLCAPS? This is possible since v1.2.0 (2017/04/17).