easyformat, *tests*

$$\int_a^b \mathrm{d}x = b - a$$

Evert Provoost

14th July 2017 (v1.5.0)

Contents

1   Main test

These two should be identical and have the—a pixel or two error is possible—same length:

     | *Lorem **ipsum** totalem* DA **givea *zin* doram** *zett*im, liefkan. |

     | *Lorem **ipsum** totalem* DA **givea *zin* doram** *zett*im, liefkan. |

This test has had a consistent and correct result since v1.0.0 (2017/04/07) and has been fixed for TeX Live 2017 in v1.5.0 (2017/06/23).

2   *Math*, $_un d_e r_s c_o r_e s$ and $\hat{c}^i r^c o^n f^l e^x e^s$ . . .

This (↑ easyformat-syntax inside subtitles) is possible without some extra code since v1.1.0 (2017/04/09) and having a ˆ (using `\cir`) in mathmode since v1.3.0 (2017/05/28) a possibility of which the kerning was fixed in v1.4.0 (2017/06/03).

$$\int_a^b x \, \mathrm{d}x = \frac{b^2 - a^2}{2}$$

3   Mixing *styles*

YEAH, *yeah?,* YEAH!!! ← do you notice how we go from SMALLCAPS to *italics* and back to SMALLCAPS? This is possible since v1.2.0 (2017/04/17).