



- [1] Grant, J. A., & Pickup, B. T. (1995). A Gaussian Description of Molecular Shape. *The Journal of Physical Chemistry*, 99(11), 3503–3510. doi:10.1021/j100011a016
- [2] Im, W., Beglov, D., & Roux, B. (1998). Continuum solvation model: Computation of electrostatic forces from numerical solutions to the Poisson-Boltzmann equation. *Computer Physics Communications*, 111(1-3), 59–75. doi:10.1016/S0010-4655(98)00016-2
- [3] Richmond, T. J. (1984). Solvent accessible surface area and excluded volume in proteins. *Journal of Molecular Biology*, 178(1), 63–89. doi:10.1016/0022-2836(84)90231-6
- [4] Can, T., Chen, C.-I., & Wang, Y.-F. (2006). Efficient molecular surface generation using level-set methods. *Journal of Molecular Graphics and Modelling*, 25(4), 442–454. doi:10.1016/j.jmglm.2006.02.012
- [5] Chen, Z., Baker, N. A., & Wei, G. W. (2010). Differential geometry based solvation model I: Eulerian formulation. *Journal of Computational Physics*, 229(22), 8231–8258. doi:10.1016/j.jcp.2010.06.036
- [6] Dzubiella, J., Swanson, J. M. J., & McCammon, J. A. (2006). Coupling nonpolar and polar solvation free energies in implicit solvent models. *The Journal of Chemical Physics*, 124(8), 084905. doi:10.1063/1.2171192