

High Energy Analysis at KamLAND and Application to Dark Matter Search

Michinari Sakai

University of Hawaii, Manoa

michinar@hawaii.edu

July 18, 2015

Overview

Introduction

Neutrino directionality

- Issues

- Idea


- Validation

Track reconstruction and particle discrimination


- Algorithm

- Validation

KamLAND: ν detector in Japan



KamLAND: ν detector in Japan



KamLAND: features

- ▶ Commissioned: 2001
- ▶ Medium: liquid scintillator
 - ▶ Decay constants: $\tau_1 = 4.0 \text{ ns}$, $\tau_2 = 8.6 \text{ ns}$
- ▶ Size: 1 kt
- ▶ Photomultiplier tubes (Hamamatsu):
 - ▶ 1325 17-inch, 7 ns rise-time, 3.5 ns TTS
 - ▶ 779 20-inch, 10 ns rise-time, 5.5 ns TTS
 - ▶ 34 % photo-coverage
- ▶ Analysis: $\sim \text{MeV } \bar{\nu}_e$ (inverse-beta decay)
- ▶ Energy resolution: $7.0 \pm 0.1 \%$
- ▶ Vertex resolution: $13.8 \pm 2.3 \text{ cm} / \sqrt{E(\text{MeV})}$

KamLAND: features


- ▶ Commissioned: 2001
- ▶ Medium: liquid scintillator
 - ▶ Decay constants: $\tau_1 = 4.0 \text{ ns}$, $\tau_2 = 8.6 \text{ ns}$
- ▶ Size: 1 kt
- ▶ Photomultiplier tubes (Hamamatsu):
 - ▶ 1325 17-inch, 7 ns rise-time, 3.5 ns TTS
 - ▶ 779 20-inch, 10 ns rise-time, 5.5 ns TTS
 - ▶ 34 % photo-coverage
- ▶ Analysis: $\sim \text{MeV } \bar{\nu}_e$ (inverse-beta decay)
- ▶ Energy resolution: $7.0 \pm 0.1 \%$
- ▶ Vertex resolution: $13.8 \pm 2.3 \text{ cm} / \sqrt{E(\text{MeV})}$
- ▶ Directional sensitivity: **NONE**

KamLAND: features

- ▶ Commissioned: 2001
- ▶ Medium: liquid scintillator
 - ▶ Decay constants: $\tau_1 = 4.0 \text{ ns}$, $\tau_2 = 8.6 \text{ ns}$
- ▶ Size: 1 kt
- ▶ Photomultiplier tubes (Hamamatsu):
 - ▶ 1325 17-inch, 7 ns rise-time, 3.5 ns TTS
 - ▶ 779 20-inch, 10 ns rise-time, 5.5 ns TTS
 - ▶ 34 % photo-coverage
- ▶ Analysis: $\sim \text{MeV } \bar{\nu}_e$ (inverse-beta decay)
- ▶ Energy resolution: $7.0 \pm 0.1 \%$
- ▶ Vertex resolution: $13.8 \pm 2.3 \text{ cm} / \sqrt{E(\text{MeV})}$
- ▶ Directional sensitivity: NONE
- ▶ No analysis at higher energies

Directionality in water


Super-Kamiokande



- Cherenkov ring

Directionality in water


Super-Kamiokande



- ▶ Cherenkov ring
- ▶ shows charged particle direction


Directionality in water

Super-Kamiokande




- ▶ Cherenkov ring
- ▶ shows charged particle direction
- ▶ Can we do something similar in scintillator?


In scintillator...



In scintillator...




In scintillator...




- Cherenkov is emitted

In scintillator...



- ▶ Cherenkov is emitted
- ▶ Along with isotropic scintillation


In scintillator...



- ▶ Cherenkov is emitted
- ▶ Along with isotropic scintillation
- ▶ \implies Cannot simply use Cherenkov for directionality


Furthermore...

Inverse-beta decay



Furthermore...


Inverse-beta decay



- KamLAND is used to seeing simple kinematics at low energies (MeV)

Furthermore...


Inverse-beta decay




- ▶ KamLAND is used to seeing simple kinematics at low energies (MeV)
- ▶ Single final-state lepton

But at higher energies, the kinematics is not so simple


Resonance production




Deep inelastic scattering




Many photons at high energy in scintillator




Hit time vs angle




Many photons at high energy in scintillator




Hit time vs angle




Many photons at high energy in scintillator




Hit time vs angle



Many photons at high energy in scintillator



Hit time vs angle



- Or we can just use **LAPPDs**!

There are many problems...

- ▶ Light is emitted isotropically

There are many problems...

- ▶ Light is emitted isotropically
- ▶ At high energies:

There are many problems...

- ▶ Light is emitted isotropically
- ▶ At high energies:
 - ▶ complicated kinematics
 - ▶ multiple final-state particles

There are many problems...

- ▶ Light is emitted isotropically
- ▶ At high energies:
 - ▶ complicated kinematics
 - ▶ multiple final-state particles
- ▶ Many photons \implies pre-pulsing

Let's change perspective and
think more simple


Let's change perspective and think more simple

- ▶ There are two pieces of information arriving at PMTs

Let's change perspective and think more simple


- ▶ There are two pieces of information arriving at PMTs
 - ▶ Charge
 - ▶ Time

Fit direction with charge and time




Fit direction with charge and time

- Use center of charge to fit middle of track




Fit direction with charge and time



- ▶ Use center of charge to fit middle of track
- ▶ Use center of time to fit near end of track

Fit direction with charge and time



- ▶ Use **center of charge** to fit middle of track
- ▶ Use **center of time** to fit near end of track
- ▶ And just connect dots to find direction!

Question:

Question:


- ▶ But, what do we use for the weights in the **weighted mean**:

$$\frac{\sum_i w_i x_i}{\sum_i w_i}$$


when calculating center of charge and time?

Let's review some basic physics...

What weight is used for *center of gravity*?




What weight is used for *center of gravity*?



To find center of gravity:

$$\text{net torque} = -(m_1g)l_1 + (m_2g)l_2 = 0$$

What weight is used for *center of gravity*?




To find center of gravity:

$$\text{net torque} = -(m_1g)l_1 + (m_2g)l_2 = 0$$

$$\implies -(m_1)l_1 + (m_2)l_2 = 0$$

What weight is used for *center of gravity*?




To find center of gravity:

$$\text{net torque} = -(m_1g)l_1 + (m_2g)l_2 = 0$$


$$\implies -(m_1)l_1 + (m_2)l_2 = 0$$

\therefore weight is **mass**: $w_i = m_i$

What weight is used for *center of charge*?




What weight is used for *center of charge*?



$$q_1 \propto \frac{1}{l_1^2}, \quad q_2 \propto \frac{1}{l_2^2}$$


What weight is used for *center of charge*?



$$q_1 \propto \frac{1}{l_1^2}, \quad q_2 \propto \frac{1}{l_2^2}$$

$$\Rightarrow \sqrt{q_1} \propto \frac{1}{l_1}, \quad \sqrt{q_2} \propto \frac{1}{l_2}$$

What weight is used for *center of charge*?




$$q_1 \propto \frac{1}{l_1^2}, \quad q_2 \propto \frac{1}{l_2^2}$$

$$\Rightarrow \sqrt{q_1} \propto \frac{1}{l_1}, \quad \sqrt{q_2} \propto \frac{1}{l_2}$$

$$\Rightarrow -(\sqrt{q_1})l_1 + (\sqrt{q_2})l_2 = 0$$

What weight is used for *center of charge*?




$$q_1 \propto \frac{1}{l_1^2}, \quad q_2 \propto \frac{1}{l_2^2}$$

$$\Rightarrow \sqrt{q_1} \propto \frac{1}{l_1}, \quad \sqrt{q_2} \propto \frac{1}{l_2}$$


$$\Rightarrow -(\sqrt{q_1})l_1 + (\sqrt{q_2})l_2 = 0$$

$$\therefore \text{weight is } \sqrt{\text{charge}}: w_i = \sqrt{q_i}$$

What weight is used for *center of time*?




What weight is used for *center of time*?



Let $\Delta t_i \equiv t_i - t_0$

$$\implies \Delta t_1 = \frac{l_1}{c}, \quad \Delta t_2 = \frac{l_2}{c}$$

What weight is used for *center of time*?




Let $\Delta t_i \equiv t_i - t_0$

$$\implies \Delta t_1 = \frac{l_1}{c}, \quad \Delta t_2 = \frac{l_2}{c}$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)\frac{l_1}{c} + \left(\frac{1}{\Delta t_2}\right)\frac{l_2}{c} = 0$$

What weight is used for *center of time*?




Let $\Delta t_i \equiv t_i - t_0$

$$\implies \Delta t_1 = \frac{l_1}{c}, \quad \Delta t_2 = \frac{l_2}{c}$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)\frac{l_1}{c} + \left(\frac{1}{\Delta t_2}\right)\frac{l_2}{c} = 0$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)l_1 + \left(\frac{1}{\Delta t_2}\right)l_2 = 0$$

What weight is used for *center of time*?



Let $\Delta t_i \equiv t_i - t_0$

$$\implies \Delta t_1 = \frac{l_1}{c}, \quad \Delta t_2 = \frac{l_2}{c}$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)\frac{l_1}{c} + \left(\frac{1}{\Delta t_2}\right)\frac{l_2}{c} = 0$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)l_1 + \left(\frac{1}{\Delta t_2}\right)l_2 = 0$$

\therefore weight is **inverse of time**: $w_i = \frac{1}{\Delta t_i}$

Conclusion

- ▶ Use **mass** as weight for *center of gravity*.

Conclusion


- ▶ Use **mass** as weight for *center of gravity*.
- ▶ Use $\sqrt{\text{charge}}$ as weight for *center of charge*.

Conclusion


- ▶ Use **mass** as weight for *center of gravity*.
- ▶ Use $\sqrt{\mathbf{charge}}$ as weight for *center of charge*.
- ▶ Use $\left(\frac{1}{\mathbf{time}}\right)$ as weight for *center of time*.

Test algorithm against μ (Data)

Cosmic ray μ





Agreement with μ -fitter
which uses
entry/exit points



Test algorithm against ν (MC)

$$\nu_e + \text{H} \longrightarrow e^- + \dots$$

$$\nu_e + {}^{12}\text{C} \xrightarrow{\text{CC}} e^- + \dots$$



- ▶ Black line: 1σ of reconstructed angle from ν direction
- ▶ Red line: 1σ of lepton angle from ν direction

Test algorithm against T2K events (Data)


(Selected with spill-time so no backgrounds)

Map



Agreement with J-PARC

direction




Test algorithm against T2K events (Data)

(Selected with spill-time so no backgrounds)

Map



Agreement with MC



Track Reconstruction and Particle ID

Hellgartner's algorithm

(former LENA grad student)

$$h(\vec{x}, t) = \sum_{i=1}^{N_{\text{PMT}}} \Theta(q_i - q_{\text{threshold}}) \sum_{j=1}^{N_{\gamma}} f(t_{ij} - t_i^{\text{TOF}}, t)$$

where N_{PMT} : number of PMTs

N_{γ} : number of photon hits to count per PMT

q_i : charge on i -th PMT, $q_{\text{threshold}}$: minimum charge for analysis


t_{ij} : j -th hit time on i -th PMT

t_i^{TOF} : expected time-of-flight between i -th PMT and \vec{x}

$$f(\Delta t, t) \propto (t - \Delta t) \exp \left[-\frac{(\Delta t - t)^2}{2\sigma_{\text{tts}}} \right]$$


Figure of merit for each test point in space $= \int_{-\infty}^{\infty} |h(\vec{x}, t)|^2 dt$

Test Hellgartner on double 1 GeV muons (MC)






Dominikus Hellgartner

Test Hellgartner on 2 GeV ν_e (MC)



Test Hellgartner on T2K events (Data)



Lepton discrimination algorithm

Explanation is here.

Test lepton discrimination (MC)

Reconstructed Ellipticity

