

High Energy Analysis at KamLAND and Application to Dark Matter Search

Michinari Sakai

University of Hawaii, Manoa

michinar@hawaii.edu

July 17, 2015

Overview

Introduction

Neutrino directionality

- Issues

- Idea


- Validation

Track reconstruction and particle discrimination

- Algorithm

- Validation

KamLAND: ν detector in Japan



KamLAND: features

- ▶ Commissioned: 2001
- ▶ Detector medium: liquid scintillator
- ▶ Size: 1 kt
- ▶ Photomultiplier tubes:
1325 17-inch, 779 20-inch (Hamamatsu), 34 %
photo-coverage
- ▶ Analysis ν energy: \sim MeV
- ▶ Energy resolution: 7.0 ± 0.1 %
- ▶ Vertex resolution: 13.8 ± 2.3 cm/ $\sqrt{E(\text{MeV})}$

KamLAND: features


- ▶ Commissioned: 2001
- ▶ Detector medium: liquid scintillator
- ▶ Size: 1 kt
- ▶ Photomultiplier tubes:
1325 17-inch, 779 20-inch (Hamamatsu), 34 %
photo-coverage
- ▶ Analysis ν energy: \sim MeV
- ▶ Energy resolution: 7.0 ± 0.1 %
- ▶ Vertex resolution: 13.8 ± 2.3 cm/ $\sqrt{E(\text{MeV})}$
- ▶ Directional sensitivity: NONE

KamLAND: features

- ▶ Commissioned: 2001
- ▶ Detector medium: liquid scintillator
- ▶ Size: 1 kt
- ▶ Photomultiplier tubes:
1325 17-inch, 779 20-inch (Hamamatsu), 34 %
photo-coverage
- ▶ Analysis ν energy: \sim MeV
- ▶ Energy resolution: 7.0 ± 0.1 %
- ▶ Vertex resolution: 13.8 ± 2.3 cm/ $\sqrt{E(\text{MeV})}$
- ▶ Directional sensitivity: NONE
- ▶ No analysis at higher energies

Directionality in water


Super-Kamiokande



- Cherenkov rings

Directionality in water


Super-Kamiokande



- ▶ Cherenkov rings
- ▶ Tell charged particle direction


Directionality in water

Super-Kamiokande




- ▶ Cherenkov rings
- ▶ Tell charged particle direction
- ▶ Can we do something similar in scintillator?


In scintillator...



In scintillator...




In scintillator...




- Cherenkov is emitted

In scintillator...



- ▶ Cherenkov is emitted
- ▶ Along with isotropic Scintillation


In scintillator...



- ▶ Cherenkov is emitted
- ▶ Along with isotropic Scintillation
- ▶ \implies Cannot simply use Cherenkov for directionality


Furthermore...

Inverse-beta decay



Furthermore...


Inverse-beta decay



- KamLAND is used to seeing simple kinematics at low energies (MeV)

Furthermore...


Inverse-beta decay




- ▶ KamLAND is used to seeing simple kinematics at low energies (MeV)
- ▶ single final-state lepton

But at higher energies, the kinematics is not so simple


Resonance production




Deep inelastic scattering




Many photons at high energy in scintillator




204 μ 's overlaid




Many photons at high energy in scintillator




204 μ 's overlaid



Many photons at high energy in scintillator




204 μ 's overlaid




- Prepulsing is few % effect.

Many photons at high energy in scintillator



204 μ 's overlaid



- ▶ Prepulsing is few % effect.
- ▶ fitters must be robust against these statistical outliers

Problems are...

- ▶ Light is produced isotropically

Problems are...

- ▶ Light is produced isotropically
- ▶ At high energies
 - ▶ complicated kinematics
 - ▶ multiple final-state particles

Problems are...

- ▶ Light is produced isotropically
- ▶ At high energies
 - ▶ complicated kinematics
 - ▶ multiple final-state particles
- ▶ Let's change perspective and think more simple


Problems are...

- ▶ Light is produced isotropically
- ▶ At high energies
 - ▶ complicated kinematics
 - ▶ multiple final-state particles
- ▶ Let's change perspective and think more simple
- ▶ There are two pieces of information arriving at PMTs

Problems are...


- ▶ Light is produced isotropically
- ▶ At high energies
 - ▶ complicated kinematics
 - ▶ multiple final-state particles
- ▶ Let's change perspective and think more simple
- ▶ There are two pieces of information arriving at PMTs
 - ▶ Time
 - ▶ Charge

Fit direction with charge and time




Fit direction with charge and time

- Use **center of charge** to fit middle of track




Fit direction with charge and time



- ▶ Use **center of charge** to fit middle of track
- ▶ Use **center of time** to fit near one end of track

Fit direction with charge and time



- ▶ Use **center of charge** to fit middle of track
- ▶ Use **center of time** to fit near one end of track
- ▶ And just connect dots to find direction!

Question:

Question:

- ▶ But, what do we use for the weights in the **weighted mean**:

$$\frac{\sum_i w_i x_i}{\sum_i w_i},$$

when calculating center of charge and time?

Question:


- ▶ But, what do we use for the weights in the **weighted mean**:

$$\frac{\sum_i w_i x_i}{\sum_i w_i},$$


when calculating center of charge and time?

- ▶ Let's review some basic physics

What weight is used for *center of gravity*?




What weight is used for *center of gravity*?



To find center of gravity:

$$\text{net torque} = -(m_1g)l_1 + (m_2g)l_2 = 0$$

What weight is used for *center of gravity*?




To find center of gravity:

$$\text{net torque} = -(m_1g)l_1 + (m_2g)l_2 = 0$$

$$\implies -m_1l_1 + m_2l_2 = 0$$

What weight is used for *center of gravity*?




To find center of gravity:

$$\text{net torque} = -(m_1g)l_1 + (m_2g)l_2 = 0$$


$$\implies -m_1l_1 + m_2l_2 = 0$$

\therefore weight is mass: $w_i = m_i$

What weight is used for *center of charge*?




What weight is used for *center of charge*?



$$q_1 \propto \frac{1}{l_1^2}, \quad q_2 \propto \frac{1}{l_2^2}$$


What weight is used for *center of charge*?



$$q_1 \propto \frac{1}{l_1^2}, \quad q_2 \propto \frac{1}{l_2^2}$$

$$\Rightarrow \sqrt{q_1} \propto \frac{1}{l_1}, \quad \sqrt{q_2} \propto \frac{1}{l_2}$$

What weight is used for *center of charge*?




$$q_1 \propto \frac{1}{l_1^2}, \quad q_2 \propto \frac{1}{l_2^2}$$

$$\Rightarrow \sqrt{q_1} \propto \frac{1}{l_1}, \quad \sqrt{q_2} \propto \frac{1}{l_2}$$

$$\Rightarrow -\sqrt{q_1}l_1 + \sqrt{q_2}l_2 = 0$$

What weight is used for *center of charge*?




$$q_1 \propto \frac{1}{l_1^2}, \quad q_2 \propto \frac{1}{l_2^2}$$

$$\Rightarrow \sqrt{q_1} \propto \frac{1}{l_1}, \quad \sqrt{q_2} \propto \frac{1}{l_2}$$


$$\Rightarrow -\sqrt{q_1}l_1 + \sqrt{q_2}l_2 = 0$$

$$\therefore \text{weight is square root of charge: } w_i = \sqrt{q_i}$$

What weight is used for *center of time*?




What weight is used for *center of time*?



Let $\Delta t_i \equiv t_i - t_0$

$$\implies \Delta t_1 = \frac{l_1}{c}, \quad \Delta t_2 = \frac{l_2}{c}$$

What weight is used for *center of time*?




Let $\Delta t_i \equiv t_i - t_0$

$$\implies \Delta t_1 = \frac{l_1}{c}, \quad \Delta t_2 = \frac{l_2}{c}$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)\frac{l_1}{c} + \left(\frac{1}{\Delta t_2}\right)\frac{l_2}{c} = 0$$

What weight is used for *center of time*?




Let $\Delta t_i \equiv t_i - t_0$

$$\implies \Delta t_1 = \frac{l_1}{c}, \quad \Delta t_2 = \frac{l_2}{c}$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)\frac{l_1}{c} + \left(\frac{1}{\Delta t_2}\right)\frac{l_2}{c} = 0$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)l_1 + \left(\frac{1}{\Delta t_2}\right)l_2 = 0$$

What weight is used for *center of time*?



Let $\Delta t_i \equiv t_i - t_0$

$$\implies \Delta t_1 = \frac{l_1}{c}, \quad \Delta t_2 = \frac{l_2}{c}$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)\frac{l_1}{c} + \left(\frac{1}{\Delta t_2}\right)\frac{l_2}{c} = 0$$

$$\implies -\left(\frac{1}{\Delta t_1}\right)l_1 + \left(\frac{1}{\Delta t_2}\right)l_2 = 0$$

\therefore weight is inverse of time: $w_i = \frac{1}{\Delta t_i}$

Conclusion

- ▶ Use **mass** as weight for *center of gravity*.


Conclusion

- ▶ Use **mass** as weight for *center of gravity*.
- ▶ Use $\sqrt{\text{charge}}$ as weight for *center of charge*.


Conclusion

- ▶ Use **mass** as weight for *center of gravity*.
- ▶ Use $\sqrt{\mathbf{charge}}$ as weight for *center of charge*.
- ▶ Use $\left(\frac{1}{\mathbf{time}}\right)$ as weight for *center of time*.



Test algorithm against μ (Data)



Deviation from
 μ -fitter




Test algorithm against ν (MC)





- ▶ Black line: 1σ of reconstructed angle from ν direction
- ▶ Red line: 1σ of lepton angle from ν direction

Test algorithm against T2K events (Data)



Test algorithm against T2K events (Data)



Track Reconstruction and Particle ID

Hellgartner's algorithm

$$h(\vec{x}, t) = \sum_{i=1}^{N_{\text{PMT}}} \Theta(q_i - q_{\text{threshold}}) \sum_{j=1}^{N_{\gamma}} f(t_{ij} - t_i^{\text{TOF}}, t)$$

where N_{PMT} : number of PMTs

N_{γ} : number of photon hits to count per PMT

q_i : charge on i -th PMT, $q_{\text{threshold}}$: minimum charge for analysis


t_{ij} : j -th hit time on i -th PMT

t_i^{TOF} : expected time-of-flight between i -th PMT and \vec{x}

$$f(\Delta t, t) \propto (t - \Delta t) \exp \left[-\frac{(\Delta t - t)^2}{2\sigma_{\text{tts}}} \right]$$


Figure of merit for each test point in space = $\int_{-\infty}^{\infty} |h(\vec{x}, t)|^2 dt$

Test Hellgartner on double 1 GeV muons (MC)






Dominikus Hellgartner

Test Hellgartner on 2 GeV ν_e (MC)



Test Hellgartner on T2K events (Data)



Lepton discrimination algorithm

Explanation is here.

Test lepton discrimination (MC)

Reconstructed Ellipticity

