

High Energy Analysis at KamLAND and Application to Dark Matter Search

Michinari Sakai

University of Hawaii, Manoa


michinar@hawaii.edu

July 18, 2015

Overview

Test Hellgartner on T2K events (Data)

$$E_{\text{reconstructed}} = 363 \text{ MeV}$$




Lepton discrimination algorithm

Explanation is here.

Test lepton discrimination (MC)


Reconstructed Ellipticity




Analysis on Dark Matter Search

Dark matter detection scheme


Signal: Dark matter (WIMP)
annihilation induced ν



Background: atmospheric ν



Earth Model (PREM)





Neutrino Oscillation Parameters

(normal hierarchy, PDG 2014)

- ▶ $\sin^2(2\theta_{12}) = 0.846 \pm 0.021$
 $\implies \theta_{12} = 33.45^\circ$
- ▶ $\sin^2(2\theta_{13}) = (9.3 \pm 0.8) \times 10^{-2}$
 $\implies \theta_{13} = 8.88^\circ$
- ▶ $\sin^2(2\theta_{23}) = 0.999^{+0.001}_{-0.018}$
 $\implies \theta_{23} = 44.09^\circ$
- ▶ $\Delta m_{21}^2 = 7.53 \pm 0.18 \times 10^{-5} \text{ eV}$
- ▶ $\Delta m_{31}^2 = 2.52 \pm 0.06 \times 10^{-3} \text{ eV}$


1 GeV ν_e oscillation probability

$P(\nu_e \rightarrow \nu_x)$ from inside Earth to KamLAND



Dark matter capture in Earth vs mass m_x

(Spin-independent cross-section $\sigma_{\text{SI}} = 1 \times 10^{-40} \text{ cm}^2$)



Data Selection

- ▶ Runs: 1330 to 12 475
- ▶ Total Live Time: 3671 days
- ▶ Selection Criteria:
 - ▶ Outer detector PMT hits < 5 (want fully contained events)
 - ▶ $E_{\text{reconstructed}} \geq 1 \text{ GeV}$