	I = -60 A	I = -100 A
G4Pions	$70.0 \pm 3.0 \%$	$81.9 \pm 2.1 \%$
G4Muons	$15.1 \pm 1.0 \%$	$13.4 \pm 0.6 \%$
G4Electrons	$14.9 \pm 1.0 \%$	$4.7 \pm 0.3 \%$

Table 1: Simulated beamline composition per magnet settings

$$E_{\text{Front Face}}^{\text{kin}} = \sqrt{p_{Beam}^2 + m_{Beam}^2 - m_{Beam}^2 - E_{Loss}}$$
 (1)

$$E_j^{\text{kin}} = E_{\text{Front Face}}^{\text{kin}} - \sum_{j < i} E_{\text{dep i}}$$
 (2)

$$E_i = \sqrt{p_{Beam}^2 + m_{Beam}^2 - m_{Beam}^2 - E_{Loss} - E_{dep FF-i}}$$
 (3)

$$\delta E_i = \sqrt{\delta p_{Beam}^2 + \delta E_{Loss}^2 + \delta E_{dep FF-i}^2}$$
 (4)

$$E_{\text{dep FF-i}} = \sum_{j < i} E_{\text{dep i}} \Rightarrow \delta E_{\text{dep FF-i}} = (i - 1)\delta E_{\text{dep i}}$$
 (5)

$$\sigma_{TOT}^{\pi^{-}}(E_i) = \frac{1}{n \ \delta X} \frac{\epsilon^{\text{Inc}}(E_i) \ C_{\text{Int}}^{\pi MC}(E_i) \ N_{\text{Int}}^{\text{TOT}}(E_i)}{\epsilon^{\text{Int}}(E_i) \ C_{\text{Inc}}^{\pi MC}(E_i) \ N_{\text{Inc}}^{\text{TOT}}(E_i)}.$$
 (6)

$$\sigma_{TOT}^{K^+}(E_i) = \frac{1}{n \ \delta X} \frac{\epsilon^{\text{Inc}}(E_i) \ C_{\text{Int}}^{KMC}(E_i) \ N_{\text{Int}}^{\text{TOT}}(E_i)}{\epsilon^{\text{Int}}(E_i) \ C_{\text{Inc}}^{KMC}(E_i) \ N_{\text{Inc}}^{\text{TOT}}(E_i)}.$$
 (7)

$$\mathcal{L}(\mu_0; \sigma_0^2; \Delta\theta_0, \Delta\theta_1) = \prod_{i=0}^1 f_X(\Delta\theta_i, \mu_0, \sigma_0^2) \Rightarrow \tag{8}$$

$$\log \mathcal{L} = -\frac{1}{2}\log(2\pi) - \log\sigma_0 - \frac{1}{2}\frac{(\Delta\theta_0 - \mu_0)^2}{\sigma_0^2} + \text{same for } \Delta\theta_1$$
 (9)

	Run-II Neg Pol
1. Events Reconstructed in Beamline	158396
2. Events with Plausible Trajectory	147468
3. Beamline $\pi^-/\mu^-/e^-$ Candidate	138481
4. Events Surviving Pile Up Filter	108929
5. Events with WC2TPC Match	41757
6. Events Surviving Shower Filter	40841
7. Available Events For Cross Section	40841

Table 2: Number of data events for Run-II Negative and Positive polarity