


Accepted Manuscript

Multinational retrospective case-control study of risk factors for the development of late invasive pulmonary aspergillosis following kidney transplantation

Francisco López-Medrano, Mario Fernández-Ruiz, José Tiago Silva, Peggy L. Carver, Christian van Delden, Esperanza Merino, María José Pérez-Saez, Milagros Montero, Julien Coussement, Milene de Abreu Mazzolin, Carlos Cervera, Lidia Santos, Nuria Sabé, Anne Scemla, Elisa Cordero, Leónidas Cruzado-Vega, Paloma Leticia Martín-Moreno, Óscar Len, Eddison Rudas, Alfredo Ponce de León, Mariano Arriola, Ricardo Lauzurica, Miruna D. David, Claudia González-Rico, Fernando Henríquez-Palop, Jesús Fortún, Marcio Nucci, Oriol Manuel, José Ramón Paño-Pardo, Miguel Montejo, Antonio Vena, Beatriz Sánchez-Sobrinho, Auxiliadora Mazuecos, Julio Pascual, Juan Pablo Horcajada, Thanh Lecompte, Asunción Moreno, Jordi Carratalà, Marino Blanes, Domingo Hernández, Erick Alejandro Hernández-Méndez, María Carmen Fariñas, Manuel Perelló-Carrascosa, Patricia Muñoz, Amado Andrés, José María Aguado



PII: S1198-743X(17)30335-X

DOI: [10.1016/j.cmi.2017.06.016](https://doi.org/10.1016/j.cmi.2017.06.016)

Reference: CMI 983

To appear in: *Clinical Microbiology and Infection*

Received Date: 31 March 2017

Revised Date: 14 June 2017

Accepted Date: 15 June 2017

Please cite this article as: López-Medrano F, Fernández-Ruiz M, Silva JT, Carver PL, van Delden C, Merino E, Pérez-Saez MJ, Montero M, Coussement J, de Abreu Mazzolin M, Cervera C, Santos L, Sabé N, Scemla A, Cordero E, Cruzado-Vega L, Martín-Moreno PL, Len Ó, Rudas E, Ponce de León A, Arriola M, Lauzurica R, David MD, González-Rico C, Henríquez-Palop F, Fortún J, Nucci M, Manuel O, Paño-Pardo JR, Montejo M, Vena A, Sánchez-Sobrinho B, Mazuecos A, Pascual J, Horcajada JP, Lecompte T, Moreno A, Carratalà J, Blanes M, Hernández D, Hernández-Méndez EA, Fariñas MC, Perelló-Carrascosa M, Muñoz P, Andrés A, Aguado JM, the Spanish Network for Research in Infectious Diseases (REIPI), the Group for the Study of Infection in Transplant Recipients (GESITRA) of the Spanish Society of Clinical Microbiology and Infectious Diseases (SEIMC), the Study Group for Infections in Compromised Hosts (ESGICH) of the European Society of Clinical Microbiology and

Infectious Diseases (ESCMID), the Swiss Transplant Cohort Study (STCS), Multinational retrospective case-control study of risk factors for the development of late invasive pulmonary aspergillosis following kidney transplantation, *Clinical Microbiology and Infection* (2017), doi: 10.1016/j.cmi.2017.06.016.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Revised paper (CLM-11767.R1) [marked version]**

2 **Original article**

3 **Title page**

4 **Complete title:** Multinational retrospective case-control study of risk factors for the
5 development of late invasive pulmonary aspergillosis following kidney transplantation.

6 **Running title:** Late pulmonary aspergillosis in kidney transplantation

7 **Authors' names and affiliations:**

8 Francisco López-Medrano (1)*; Mario Fernández-Ruiz (1)*; José Tiago Silva (1); Peggy L.
9 Carver (2); Christian van Delden (3); Esperanza Merino (4); María José Pérez-Saez (5)
10 Milagros Montero (6); Julien Coussement (7); Milene de Abreu Mazzolin (8), Carlos Cervera (9);
11 Lidia Santos (10); Nuria Sabé (11); Anne Scemla (12); Elisa Cordero (13); Leónidas Cruzado-
12 Vega (14); Paloma Leticia Martín-Moreno (15); Óscar Len (16); Eddison Rudas (17); Alfredo
13 Ponce de León (18); Mariano Arriola (19); Ricardo Lauzurica (20); Miruna D. David (21);
14 Claudia González-Rico (22); Fernando Henríquez-Palop (23); Jesús Fortún (24); Marcio Nucci
15 (25); Oriol Manuel (26); José Ramón Paño-Pardo (27); Miguel Montejo (28); Antonio Vena (29);
16 Beatriz Sánchez-Sobrinó (30); Auxiliadora Mazuecos (31); Julio Pascual (5); Juan Pablo
17 Horcajada (6); Thanh Lecompte (3); Asunción Moreno (9); Jordi Carratalà (11); Marino Blanes
18 (32); Domingo Hernández (17); Erick Alejandro Hernández-Méndez (18); María Carmen
19 Fariñas (22); Manuel Perelló-Carrascosa (33); Patricia Muñoz (29); Amado Andrés (34); José
20 María Aguado (1); and the Spanish Network for Research in Infectious Diseases (REIPI), the
21 Group for the Study of Infection in Transplant Recipients (GESITRA) of the Spanish Society of
22 Clinical Microbiology and Infectious Diseases (SEIMC), the Study Group for Infections in
23 Compromised Hosts (ESGICH) of the European Society of Clinical Microbiology and Infectious
24 Diseases (ESCMID), and the Swiss Transplant Cohort Study (STCS).

- 25 1. Unit of Infectious Diseases, Hospital Universitario "12 de Octubre", Instituto de
26 Investigación Hospital "12 de Octubre" (i+12), Department of Medicine, School of
27 Medicine, Universidad Complutense, Madrid, Spain
- 28 2. University of Michigan College of Pharmacy, Department of Clinical Pharmacy,
29 University of Michigan Health System, Ann Arbor, Michigan

- 30 3. Unit of Transplant Infectious Diseases, Department of Medical Specialities, University
31 Hospitals Geneva, Geneva, Switzerland
- 32 4. Unit of Infectious Diseases, Hospital Universitario General, Alicante, Spain
- 33 5. Department of Nephrology, Hospital del Mar, Hospital del Mar Medical Research
34 Institute (IMIM), Barcelona, Spain
- 35 6. Department of Infectious Diseases, Hospital del Mar, Hospital del Mar Medical
36 Research Institute (IMIM), Barcelona, Spain
- 37 7. Department of Nephrology, Dialysis and Kidney Transplantation, CUB-Hôpital Erasme,
38 Université Libre de Bruxelles, Brussels, Belgium
- 39 8. Division of Nephrology, Department of Medicine, Universidade Federal de São Paulo-
40 UNIFESP and Hospital do Rim e Hipertensão, Fundação Oswaldo Ramos, São Paulo,
41 Brazil
- 42 9. Department of Infectious Diseases, Hospital Clinic, Institut d'Investigacions
43 Biomèdiques August Pi i Sunyer (IDIBAPS), School of Medicine, University of
44 Barcelona, Barcelona, Spain
- 45 10. Unit of Renal Transplantation, Department of Urology and Kidney Transplantation,
46 Coimbra Hospital and University Centre, Coimbra, Portugal
- 47 11. Department of Infectious Diseases, Hospital Universitari de Bellvitge, Institut
48 d'Investigació Biomèdica de Bellvitge (IDIBELL), Barcelona, Spain
- 49 12. Service de Néphrologie et Transplantation Adulte, Hôpital Necker Enfants Malades,
50 Assistance Publique-Hôpitaux de Paris, Université Paris Descartes Sorbonne Paris
51 Cité, RTRS Centaure, Paris, France
- 52 13. Department of Infectious Diseases, Hospitales Universitarios "Vigen del Rocío",
53 Instituto de Biomedicina de Sevilla (IBIS), Seville, Spain
- 54 14. Department of Nephrology, Hospital Universitario "La Fe", Valencia, Spain
- 55 15. Department of Nephrology, Clínica Universitaria de Navarra, Pamplona, Spain
- 56 16. Department of Infectious Diseases, Hospital Universitari Vall d'Hebrón, Vall d'Hebron
57 Research Institute (VHIR), Barcelona, Spain
- 58 17. Department of Nephrology, Hospital Universitario "Carlos Haya", Málaga, Spain

- 59 18. Department of Transplantation, Instituto Nacional de Ciencias Médicas y Nutrición
60 "Salvador Zubirán", México DF, México
- 61 19. Clínica de Nefrología, Urología y Enfermedades Cardiovasculares, Santa Fe,
62 Argentina
- 63 20. Department of Nephrology, University Hospital Germans Trias i Pujol, Badalona,
64 Barcelona, Spain
- 65 21. Department of Microbiology, University Hospitals Birmingham NHS Foundation Trust,
66 Birmingham, UK
- 67 22. Department of Infectious Diseases, University Hospital "Marqués de Valdecilla",
68 Santander, Spain
- 69 23. Department of Nephrology, University Hospital "Doctor Negrín", Las Palmas de Gran
70 Canaria, Spain
- 71 24. Department of Infectious Diseases, University Hospital "Ramón y Cajal", Madrid,
72 Spain
- 73 25. Department of Internal Medicine, Hematology Service and Mycology Laboratory,
74 Hospital Universitário Clementino Fraga Filho, Universidade Federal do Rio de
75 Janeiro, Rio de Janeiro, Brasil
- 76 26. Department of Infectious Diseases and Transplantation Center, University Hospital
77 (CHUV) and University of Lausanne, Switzerland
- 78 27. Infectious Diseases and Clinical Microbiology Unit, Department of Internal Medicine,
79 Hospital Universitario "La Paz", School of Medicine, Universidad Autónoma de Madrid,
80 Madrid, Spain
- 81 28. Department of Infectious Diseases, Hospital Universitario Cruces, Barakaldo, Bilbao,
82 Spain
- 83 29. Department of Clinical Microbiology and Infectious Diseases, Hospital General
84 Universitario Gregorio Marañón; Instituto de Investigación Sanitaria Hospital Gregorio
85 Marañón, Madrid, Spain; CIBER Enfermedades Respiratorias - CIBERES
86 (CB06/06/0058), Madrid Spain; Medicine Department, School of Medicine,
87 Universidad Complutense de Madrid, Madrid, Spain

30. Department of Nephrology, Hospital Universitario Puerta de Hierro-Majadahonda,
School of Medicine, Universidad Autónoma de Madrid, Madrid, Spain
31. Department of Nephrology, Hospital Universitario "Puerta del Mar", Cádiz, Spain
32. Unit of Infectious Diseases, Hospital Universitario "La Fe", Valencia, Spain
33. Department of Nephrology, Hospital Universitari Vall d'Hebrón, Vall d'Hebron
Research Institute (VHIR), Barcelona, Spain
34. Department of Nephrology, Hospital Universitario "12 de Octubre", Instituto de
Investigación Hospital "12 de Octubre" (i+12), Department of Medicine, School of
Medicine, Universidad Complutense, Madrid, Spain

* Both authors contributed equally to this work

- **Manuscript word length** (excluding title page, abstract, references, tables and figure legends): 2,582
- **Number of tables:** 3
- **Number of figures:** 1
- **Number of references:** 28
- **Keywords:** kidney transplantation; late invasive pulmonary aspergillosis; risk factors; case-control study.
- **Corresponding author:** Francisco López-Medrano, MD, PhD, MSc. Unit of Infectious Diseases, Hospital Universitario "12 de Octubre", Centro de Actividades Ambulatorias, 2ª planta, bloque D. Avda. de Córdoba, s/n. Postal code 28041. Madrid, Spain. Phone: +34 913908000. Fax: +34 914695775. E-mail address: **flmedrano@yahoo.es**

Abstract (250 words)

Objectives: To assess the risk factors for the development of late-onset invasive pulmonary aspergillosis (IPA) after kidney transplantation (KT).

Methods: We performed a multinational case-control study that retrospectively recruited 112 KT recipients diagnosed with IPA between 2000 and 2013. Controls were matched (1:1 ratio) by center and date of transplantation. Immunosuppression-related events (IREs) included the occurrence of non-ventilator-associated pneumonia, tuberculosis, cytomegalovirus disease and/or *de novo* malignancy.

Results: We identified 61 cases of late (>180 days after transplantation) IPA from 24 participating centers (accounting for 54.5% [61/112] of all cases included in the overall study). Most diagnoses (54.1% [33/61]) were established within the first 36 post-transplant months, although 5 cases occurred more than 10 years after transplantation. Overall mortality among cases was 47.5% (29/61). Compared to controls, cases were significantly older (P -value = 0.010) and more likely to have pre-transplant chronic obstructive pulmonary disease (P -value = 0.001) and a diagnosis of bloodstream infection (P -value = 0.016) and IRE (P -value <0.001) within the 6 months prior to the onset of late IPA. After multivariate adjustment, previous occurrence of IRE (odds ratio: 19.26; 95% confidence interval: 2.07 - 179.46; P -value = 0.009) was identified as an independent risk factor for late IPA.

Conclusion: More than half of IPA cases after KT occur beyond the sixth month, with some of them presenting very late. Late IPA entails a poor prognosis. We identified some risk factors that could help the clinician to delimit the subgroup of KT recipients at the highest risk for late IPA.

Introduction

Invasive pulmonary aspergillosis (IPA) constitutes one of the most feared complications occurring in patients undergoing solid organ transplantation (SOT) in terms of both patient and graft survival [1-3]. Apart from local susceptibility associated with specific surgical procedures (e.g., ulcerative aspergillus tracheobronchitis at the bronchial anastomosis site after lung transplantation) [4], it is conventionally assumed that the lifelong use of immunosuppression to avoid graft rejection confers the most relevant risk for this event [5].

The intensity of the immunosuppressive therapy is usually higher during the first 6 months following SOT, and therefore this period has been traditionally considered as carrying the maximum risk for opportunistic infection including IPA [6]. Nevertheless, kidney transplant (KT) recipients require potent triple-drug regimens —often containing steroids, calcineurin inhibitors and antiproliferative agents— for indefinite time periods [7]. Although the relative risk of post-transplant IPA after KT is lower compared to other types of grafts [1,3,8], KT recipients suffer from the highest absolute disease burden due to the large number of procedures performed worldwide [9,10]. In addition, recent decades have witnessed a continuous improvement in long-term graft survival [11], thus increasing the population of aged KT recipients chronically exposed to a high degree of immunosuppression.

By using a multicenter case-control design, we have recently analyzed the risk factors for the occurrence of early IPA (i.e., diagnosed within the first 180 days) after KT [12]. Only one previous study has analyzed the predisposing conditions for the late forms of infection, although its results were limited by its single-center nature and by the inclusion of only 26 cases of late IPA [13].

Transplant physicians may benefit from identifying, among the increasing population of long-term KT recipients, that subgroup of patients at increased risk for late IPA in order to implement individualized follow-up and prevention strategies. Unfortunately, such an approach remains an unmet clinical need. To the best of our knowledge, this is the first study specifically aiming to ascertain the predisposing factors for the development of late IPA from a large representative population of KT recipients.

Materials and Methods

Study design

This is a sub-analysis of a multinational retrospective case-control study performed in 29 hospitals from 10 European (Spain, Switzerland [6 centers included in the Swiss Transplant Cohort Study [14]], Belgium, Portugal, France and United Kingdom) and American institutions (United States, Brazil, Mexico and Argentina). Participating centers included cases of IPA diagnosed in KT recipients between January 1, 2000 and December 31, 2013 [12,15]. In the present *a priori* designed sub-analysis we focused on late episodes of IPA, defined as those diagnosed beyond the first 180 days after transplantation ("IPA cases"). The "control group" was selected (in an 1:1 ratio) among those patients that underwent transplantation at the same center within a 3-month period before or after the calendar date of the corresponding case but without the diagnosis of IPA throughout the post-transplant period. In addition, controls must have survived at least until the time of diagnosis of IPA in the index case. To take into account the effect of post-transplant events on the occurrence of late IPA, controls were assigned a "pseudo-date of diagnosis" to match their cases with the aim of ensuring comparable risk exposure periods in both groups. The criteria used to establish the date of IPA diagnosis is available as **Supplementary Methods**. This research adhered to the STROBE guidelines for observational studies. The study protocol was approved by the local Ethics Committee of the coordinating center and of other participating sites as required.

Study definitions

IPA was defined according to the revised criteria proposed in 2008 by the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group (details provided as **Supplementary Methods**) [16]. It should be noted that we added a modified radiological criterion (beyond the classic dense, well-circumscribed lesions with or without halo sign or cavitation) based on the presence of certain lung patterns that have been specifically associated with post-transplant IPA (peribronchial consolidation or tree-in-bud pattern) [17]. Additional study definitions (including IPA-attributable mortality, cytomegalovirus [CMV] disease, tuberculosis, pneumonia, respiratory tract viral infection, bloodstream infection [BSI] or post-transplant lymphoproliferative disorder [PTLD]) are

available in **Supplementary Methods**.

To encompass the different post-transplant complications that may be attributable to over-immunosuppression we constructed a composite variable (termed “immunosuppression-related event” [IRE]) that included the occurrence of any of the following: non-ventilator-associated pneumonia, tuberculosis, CMV disease and/or post-transplant *de novo* malignancy (both PTLD and solid organ tumors). Community-acquired pneumonia has been previously recognized to be more frequent among SOT recipients due to immunosuppression [18] and therefore pneumococcal vaccination is strongly recommended for this population [19]. We did not consider within the definition of IRE certain post-transplant infections (such as BSI or ventilator-associated pneumonia) that may be arguably attributable to invasive procedures, instrumentation (i.e., indwelling catheters) or anatomical abnormalities rather than to the recipient’s immune status.

Statistical analysis

Continuous variables were summarized by the mean \pm standard deviation (SD) or the median with interquartile ranges (IQR), while categorical variables were summarized using absolute counts and percentages. Categorical variables were compared using the McNemar test, whereas the Student's t-test for repeated measures or the Wilcoxon signed-ranks test were applied for continuous variables. Conditional logistic regression was used to identify independent risk factors for the development of late IPA. Those variables found to be significant (P -value ≤ 0.1) at the univariate level were included into the multivariable models in a backward stepwise fashion. Collinearity among explanatory variables was assessed by means of the variance inflation factor (VIF), with VIF values over 3 suggesting significant collinearity. Results are given as odds ratios (ORs) with 95% confidence intervals (CIs). As a secondary outcome, we compared patient survival from the date (for cases) or the “pseudodate” (for controls) of IPA diagnosis. Survival curves were plotted by the Kaplan-Meier method and differences between groups were compared with the log-rank test. All the significance tests were two-tailed. Statistical analysis was performed with SPSS version 20.0 (IBM Corp., Armonk, NY) and graphics were generated with Prism v. 6.0 (GraphPad Software Inc., La Jolla, CA).

Results

We included 61 cases of late IPA (14/61 [23.0%] proven and 47/61 [77.0%] probable) and their corresponding controls from 24 out of 29 participating centers (i.e., 5 centers did not contribute to the present sub-analysis). This figure accounts for 54.5% (61/112) of all the cases enrolled in the overall study. Twenty-nine out of 61 cases (47.5%) were diagnosed between 2010 and 2013. The median time interval between transplantation and diagnosis was 34.4 months (IQR: 11.8 - 78.5). Most diagnoses (54.1% [33/61]) were established within the first 36 months, although this period spanned more than 27 years (with 5 very late-onset cases occurring after the tenth year) (**Figure 1**). The median follow-up from the date (for cases) or the “pseudo-date” of diagnosis (for controls) was 476 days (IQR: 70.0 - 1298.5). Overall and IPA-attributable mortality among IPA cases was 47.5% (29/61) and 21.3% (13/61) and occurred at a median of 53.5 days (IQR: 14.5 - 171.5) and 15 days (IQR: 7.3 - 33.3), respectively, from diagnosis. There were no significant differences in one-year survival rates between cases occurring in months 6 to 36 or >36 months after transplantation (55.0% versus 41.0%, respectively; log-rank test P -value = 0.619). Among survivors, 9.4% (3/32) patients experienced definitive graft failure requiring return to permanent dialysis. None of the patients in the control group died during the follow-up. One-year survival was significantly lower among cases than controls (49.0% versus 100.0%; log-rank test P -value = 0.021).

The demographics and pre-transplant factors of patients who developed late IPA and their controls are compared in **Table 1**. Cases were significantly older (54.6 ± 14.2 versus 48.6 ± 15.5 years; P -value = 0.010) and more likely to have pre-transplant chronic obstructive pulmonary disease (COPD) (18.0% [11/61] versus 0.0% [0/61]; P -value = 0.001) than control counterparts. The prevalence of underlying diabetic nephropathy as a reason for end-stage renal disease requiring transplantation was also higher among cases, although not achieving statistical significance (19.7% [12/61] versus 6.6% [4/61]; P -value = 0.077).

Donor- and transplant-related and post-transplant variables are compared in **Table 2**. Cases were more likely to have been diagnosed with an IRE during the 6 months prior to the onset of IPA (34.4% [21/61] versus 3.3% [2/61]; P -value <0.001), with significant (for non-ventilator-associated pneumonia and CMV disease) or near significant differences (for post-transplant *de novo* malignancy) observed for each of the different individual events included in this composite

variable. PTLD was the predominant type of malignancy diagnosed. A prior occurrence of BSI was also more common among cases than controls (11.5% [7/61] versus 0.0% [0/61]; P -value = 0.016). No significant differences were observed between the groups regarding the prior occurrence of acute graft rejection or the requirement of steroid boluses. None of these episodes were treated with lymphocyte-depleting agents as anti-rejection therapy, and only one of them (in the control group) received rituximab.

Finally, age at transplantation, pre-transplant COPD, underlying diabetic nephropathy, and the diagnosis of an IRE or BSI within the preceding 6 months were entered into the conditional logistic regression model (**Table 3**). Linear regression analysis showed no significant collinearity among these explanatory variables, with all VIF values <1.5 (data not shown). After multivariate adjustment, prior diagnosis of IRE (OR: 19.26; 95% CI: 2.07 - 179.46; P -value = 0.009) was identified as the only independent risk factor associated with late IPA.

Discussion

To our knowledge, our multinational retrospective case-control study represents the largest effort to date to explore the clinical outcome of and risk factors for IPA in the specific population of KT recipients. Our experience highlights the poor prognosis conferred by the late forms of this opportunistic infection, since more than half of the included patients had died at a median time of less than two months from diagnosis. In addition, IPA-attributable mortality was assumed in more than 20% of cases. Notwithstanding such an ominous picture, in our previous study we reported even worse figures for early IPA (first 180 days), with global and attributable mortality of 60.8% and 45.1%, respectively [12]. We hypothesize that this difference may be explained by the relatively more intensive immunosuppression among patients in their first post-transplant months [15].

Remarkably, although most of the episodes of late IPA occurred within the first three years, almost 10% of them were diagnosed across a large time period covering more than a decade after transplantation, including some very late onset episodes occurring more than ten years post-transplantation. In a previous series of IPA among KT recipients [13], 43% of the 41 cases were diagnosed beyond the sixth month, and 6 (14%) beyond the fifth year post-transplantation. These concordant results reinforce the previously stated concept [20] that the period at risk for severe opportunistic infection continues far beyond the classical time scheme proposed for SOT recipients.

Despite the wide range of time between KT and the onset of late IPA, we were still able to identify some factors associated to this event. Cases were more likely to have been diagnosed with COPD, although such association only showed borderline univariate significance. The presence of pre-transplant COPD may reflect underlying injury to the lung parenchyma [12,21] or act as a surrogate marker for prolonged corticosteroid exposure. BSI during the six preceding months was also more frequent among cases. Comparable associations have been previously reported for the overall SOT population [8] or, specifically, KT recipients [12]. The occurrence of BSI may identify patients commonly suffering from invasive procedures, impaired graft function and antibiotic therapy exposure, which overall reflect increased patient frailty.

Following the example of previous studies [22], we created a composite variable (IRE) that summarized post-transplant complications —such as severe non-device-associated infections,

CMV disease or *de novo* cancer— that are consistently assumed to indicate an excess of immunosuppression. In the regression model this condition displayed a significant association with the development of IPA during the following six months. Other authors have also reported the observation of episodes of pneumonia preceding the onset of IPA [23,24]. On the other hand, the deleterious impact exerted by CMV on the risk of IPA has been well established for the SOT recipient [8,25,26]. In accordance with this rationale, the incidence of CMV disease in our experience was ten times higher among cases than controls (16.4% versus 1.6%, respectively). In a similar way, a recent diagnosis of *de novo* cancer (either PTLN or solid organ tumor) had been made in almost one out of every ten cases as compared to none of the controls. In a French nationwide epidemiological study, both hematologic and solid organ malignancies have been described as an important risk factor for invasive aspergillosis [3]. In addition to the direct deleterious effect of the oncologic therapies (B-cell-depleting agents such as rituximab or cytotoxic chemotherapy) on the host's response and infection susceptibility, the function of natural killer cells (which significantly contribute to the protective immunity against fungi [27]) has been shown to be impaired in KT recipients with post-transplant cancer [28].

The design of our study (case-control study) prevents us from estimating the actual incidence of late IPA among KT recipients that develop an episode of IRE. Case-control studies can generate plausible associations rather than demonstrate direct causality. In our opinion, such a circumstance and the heterogeneous distribution of IPA cases over a very long post-transplant period would make it unreasonable to propose the use of antifungal prophylaxis for those recipients fulfilling the characterized risk factors. Nevertheless, our findings do support the recommendation of maintaining a low threshold for suspicion of post-transplant IPA in patients with compatible respiratory symptoms and underlying COPD or recently diagnosed with a serious infection, CMV disease or post-transplant cancer. In addition, this clinical awareness should be maintained even for very long-term KT recipients, as IPA may occur many years after transplantation. In this context, we have previously shown the protean clinical features of IPA among KT recipients and the correlation between the timely initiation of antifungal therapy and the outcome [15].

Strengths of the present collaborative effort include its multicenter nature, the use of uniform diagnostic criteria, and the standardized collection of a large number of variables. However,

some limitations must be acknowledged, such as its retrospective design and the relatively low sample size that may have limited statistical power. Therefore, confidence intervals for risk estimates were wide. Most IPA cases were categorized as “probable” rather than “proven” [16]. The protracted inclusion period imposes heterogeneity among participating centers in immunosuppression and standard of care. Nonetheless, the low incidence among KT recipients of late-onset IPA made this approach the only practical method to collect a meaningful number of cases. We lacked detailed data on certain relevant factors (such as the receipt of rituximab or cytotoxic chemotherapy among patients with PTLT). Finally, we were unable to estimate the incidence of late IPA due to the lack of denominator figures (i.e., number of transplant procedures performed at each center or number of at-risk recipients during the study period) since our research was conceived exclusively to ascertain the risk factors for developing such condition. Thus, we chose a case-control design instead than other approaches (i.e., nested case-control study within a multicenter cohort).

In conclusion, late IPA may develop among KT recipients even more than 10 years after transplantation and entails a very poor prognosis. The preceding diagnosis of post-transplant adverse events reflecting an excess of immunosuppression, such as serious or opportunistic infection or *de novo* malignancy, may be useful to identify those patients at the highest risk for this complication.

- Acknowledgements:** The authors would like to acknowledge Mariano Barroso for his kind assistance with the creation of the database. The present study was performed under the scientific auspice of the Group for the Study of Infection in Transplant Recipients (GESITRA) of the Spanish Society of Clinical Microbiology and Infectious Diseases (SEIMC) and the Spanish Network Research of Infectious Diseases (REIPI).
- Authors' contribution:** FLM designed the study; FLM and MFR analyzed data and wrote the manuscript; FLM, MFR, JTS and JMA coordinated the study. FLM, JTS, PLC, CvD, EM, MJPS, MM, JC, MAM, CC, LS, NS, AS, EC, LCV, PLMM, OL, ER, APL, MA, TL, MD, CGR, FHP, JF, MN, OM, JRPP, MM, AV, BSS, AM, JP, JPH, TL, AM, JC, MB, DH, EAHM, MCF and MPC collected data and critically reviewed the final version of the manuscript.
- Transparency declaration:** Francisco López-Medrano has been paid for talks on behalf of Pfizer, Gilead Sciences and Astellas Pharma. Mario Fernández-Ruiz has been paid for talks on behalf of Pfizer and Gilead Sciences. Christian van Delden has been consultant for Basilea, Debiopharm, Gilead Sciences, Merck Sharp and Dohme, Pfizer and Astellas Pharma. Óscar Len has been paid for talks on behalf of Astellas Pharma and Merck, Sharp and Dohme and has received grants from Merck Sharp and Dohme and Astellas. Oriol Manuel has received unrestricted grants for research from Roche and Lophius Bioscience. Mariano Arriola has been consultant for Novartis, Pfizer and Astellas Pharma. Miruna D. David has been a consultant for Merck Sharp and Dohme and has received travel grant to attend conferences from Astellas Pharma and Gilead. Jesús Fortún has received grant support from Astellas Pharma, Gilead Sciences, Merck Sharp and Dohme, Pfizer and Instituto de Salud Carlos III. Ricardo Lauzurica has been paid for talks on behalf of Novartis and Astellas Pharma. Marino Blanes has been paid for talks on behalf of Astellas, Pfizer, Gilead and Merck Sharp and Dohme. José María Aguado has been a consultant to and on the speakers' bureau for Astellas Pharma, Pfizer, Gilead, Merck Sharp and Dohme, and Roche. The remaining authors of this manuscript have no conflicts of interest to disclose.
- Funding sources:** This research was supported by Plan Nacional de I+D+i and Instituto de Salud Carlos III (Proyecto Integrado de Excelencia [PIE] 13/00045), Subdirección

General de Redes y Centros de Investigación Cooperativa, Spanish Ministry of Economy and Competitiveness, Spanish Network for Research in Infectious Diseases (REIPI RD12/0015) - co-financed by the European Development Regional Fund (EDRF) "*A way to achieve Europe*". This study was also co-funded by an unrestricted grant from Pfizer Pharmaceutical (REI-ANT-2013-01). Mario Fernández-Ruiz holds a clinical research contract "Juan Rodés" (JR14/00036) from the Instituto de Salud Carlos III, Spanish Ministry of Economy and Competitiveness. These funding institutions had no involvement in the study design; in the collection, analysis, and interpretation of data; in the writing of the paper; or in the decision to submit it for publication.

References

1. Cornet M, Fleury L, Maslo C, Bernard JF, Brucker G, Invasive Aspergillosis Surveillance Network of the Assistance Publique-Hopitaux de P. Epidemiology of invasive aspergillosis in France: a six-year multicentric survey in the Greater Paris area. *J Hosp Infect* 2002;51:288-96.
2. Morgan J, Wannemuehler KA, Marr KA, Hadley S, Kontoyiannis DP, Walsh TJ, et al. Incidence of invasive aspergillosis following hematopoietic stem cell and solid organ transplantation: interim results of a prospective multicenter surveillance program. *Med Mycol* 2005;43 Suppl 1:S49-58.
3. Lortholary O, Gangneux JP, Sitbon K, Lebeau B, de Monbrison F, Le Strat Y, et al. Epidemiological trends in invasive aspergillosis in France: the SAIF network (2005-2007). *Clin Microbiol Infect* 2011;17:1882-9.
4. Fernández-Ruiz M, Silva JT, San-Juan R, de Dios B, García-Luján R, López-Medrano F, et al. *Aspergillus* tracheobronchitis: report of 8 cases and review of the literature. *Medicine (Baltimore)* 2012;91:261-73.
5. Singh N, Husain S, AST Infectious Diseases Community of Practice. Aspergillosis in solid organ transplantation. *Am J Transplant* 2013;13 Suppl 4:228-41.
6. Fishman JA, Issa NC. Infection in organ transplantation: risk factors and evolving patterns of infection. *Infect Dis Clin North Am* 2010;24:273-83.
7. Kidney Disease: Improving Global Outcomes Transplant Work G. KDIGO clinical practice guideline for the care of kidney transplant recipients. *Am J Transplant* 2009;9 Suppl 3:S1-155.
8. Gavalda J, Len O, San Juan R, Aguado JM, Fortún J, Lumberras C, et al. Risk factors for invasive aspergillosis in solid-organ transplant recipients: a case-control study. *Clin Infect Dis* 2005;41:52-9.
9. Pappas PG, Alexander BD, Andes DR, Hadley S, Kauffman CA, Freifeld A, et al. Invasive fungal infections among organ transplant recipients: results of the Transplant-Associated Infection Surveillance Network (TRANSNET). *Clin Infect Dis* 2010;50:1101-11.
10. Neofytos D, Treadway S, Ostrander D, Alonso CD, Dierberg KL, Nussenblatt V, et al. Epidemiology, outcomes, and mortality predictors of invasive mold infections among

transplant recipients: a 10-year, single-center experience. *Transpl Infect Dis* 2013;15:233-42.

11. Ojo AO, Hanson JA, Wolfe RA, Leichtman AB, Agodoa LY, Port FK. Long-term survival in renal transplant recipients with graft function. *Kidney Int* 2000;57:307-13.

12. López-Medrano F, Silva JT, Fernández-Ruiz M, Carver PL, van Delden C, Merino E, et al. Risk factors associated with early invasive pulmonary aspergillosis in kidney transplant recipients: results from a multinational matched case-control study. *Am J Transplant* 2016;16:2148-57.

13. Heylen L, Maertens J, Naesens M, Van Wijngaerden E, Lagrou K, Bammens B, et al. Invasive aspergillosis after kidney transplant: case-control study. *Clin Infect Dis* 2015;60:1505-11.

14. Koller MT, van Delden C, Muller NJ, Baumann P, Lovis C, Marti HP, et al. Design and methodology of the Swiss Transplant Cohort Study (STCS): a comprehensive prospective nationwide long-term follow-up cohort. *Eur J Epidemiol* 2013;28:347-55.

15. López-Medrano F, Fernández-Ruiz M, Silva JT, Carver PL, van Delden C, Merino E, et al. Clinical presentation and determinants of mortality of invasive pulmonary aspergillosis in kidney transplant recipients: a multinational cohort study. *Am J Transplant* 2016;16:3220-34.

16. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE, Calandra T, et al. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. *Clin Infect Dis* 2008;46:1813-21.

17. Muñoz P, Vena A, Ceron I, Valerio M, Palomo J, Guinea J, et al. Invasive pulmonary aspergillosis in heart transplant recipients: two radiologic patterns with a different prognosis. *J Heart Lung Transplant* 2014;33:1034-40.

18. Giannella M, Muñoz P, Alarcon JM, Mularoni A, Grossi P, Bouza E, et al. Pneumonia in solid organ transplant recipients: a prospective multicenter study. *Transpl Infect Dis* 2014;16:232-41.

19. Blasi F, Akova M, Bonanni P, Dartois N, Sauty E, Webber C, et al. Community-acquired pneumonia in adults: Highlighting missed opportunities for vaccination. *Eur J Intern Med* 2017;37:13-8.
20. San Juan R, Aguado JM, Lumbreras C, Díaz-Pedroche C, López-Medrano F, Lizasoain M, et al. Incidence, clinical characteristics and risk factors of late infection in solid organ transplant recipients: data from the RESITRA study group. *Am J Transplant* 2007;7:964-71.
21. Guinea J, Torres-Narbona M, Gijon P, Muñoz P, Pozo F, Pelaez T, et al. Pulmonary aspergillosis in patients with chronic obstructive pulmonary disease: incidence, risk factors, and outcome. *Clin Microbiol Infect* 2010;16:870-7.
22. San-Juan R, De Dios B, Navarro D, Garcia-Reyne A, Lumbreras C, Bravo D, et al. Epstein-Barr virus DNAemia is an early surrogate marker of the net state of immunosuppression in solid organ transplant recipients. *Transplantation* 2013;95:688-93.
23. Perez-Saez MJ, Mir M, Montero MM, Crespo M, Montero N, Gomez J, et al. Invasive aspergillosis in kidney transplant recipients: a cohort study. *Exp Clin Transplant* 2014;12:101-5.
24. Hoyo I, Sanclemente G, de la Bellacasa JP, Cofan F, Ricart MJ, Cardona M, et al. Epidemiology, clinical characteristics, and outcome of invasive aspergillosis in renal transplant patients. *Transpl Infect Dis* 2014;16:951-7.
25. Sahin SZ, Akalin H, Ersoy A, Yildiz A, Ocakoglu G, Cetinoglu ED, et al. Invasive fungal infections in renal transplant recipients: epidemiology and risk factors. *Mycopathologia* 2015;180:43-50.
26. Muñoz P, Valerio M, Palomo J, Giannella M, Yanez JF, Desco M, et al. Targeted antifungal prophylaxis in heart transplant recipients. *Transplantation* 2013;96:664-9.
27. Fernández-Ruiz M, López-Medrano F, San Juan R, Allende LM, Paz-Artal E, Aguado JM. Low natural killer cell counts and onset of invasive fungal disease after solid organ transplantation. *J Infect Dis* 2016;213:873-4.
28. Hope CM, Troelnikov A, Hanf W, Jesudason S, Coates PT, Heeger PS, et al. Peripheral natural killer cell and allo-stimulated T-cell function in kidney transplant recipients

465 associate with cancer risk and immunosuppression-related complications. Kidney Int
466 2015;88:1374-82.

ACCEPTED MANUSCRIPT

Tables

Table 1. Comparison of demographics and pre-transplant variables between KT recipients with and without late IPA.

Variable	Late IPA group (n = 61)	Control group (n = 61)	P-value ^a
Age, years [mean \pm SD]	54.6 \pm 14.2	48.6 \pm 15.5	0.010
Gender (male) [n (%)]	33 (54.1)	38 (62.3)	0.458
Pre-transplant conditions [n (%)]			
Diabetes mellitus	18 (29.5)	9 (14.7)	0.093
Chronic obstructive pulmonary disease	11 (18.0)	0 (0.0)	0.001
Pre-transplant corticosteroid therapy [n (%)] ^b	6 (10.3)	5 (8.8)	0.754
BMI at transplantation, Kg/m ² [mean \pm SD] ^c	24.3 \pm 3.6	26.7 \pm 7.3	0.074
Previous kidney transplantation [n (%)]	7 (11.5)	8 (13.1)	1.000
Underlying end-stage renal disease [n (%)]			
Glomerulonephritis	14 (23.0)	14 (23.0)	1.000
Diabetic nephropathy	12 (19.7)	4 (6.6)	0.077
Nephroangiosclerosis	8 (13.1)	8 (13.1)	1.000
Polycystic kidney disease	8 (13.1)	11 (18.0)	0.824
Chronic interstitial nephropathy	3 (4.9)	3 (4.9)	1.000
Congenital nephropathy	2 (3.3)	3 (4.9)	1.000
Lupus nephropathy	1 (1.6)	1 (1.6)	1.000
Reflux nephropathy	0 (0.0)	1 (1.6)	1.000
Unknown	6 (9.8)	9 (14.8)	0.388
Other	7 (11.5)	7 (11.5)	0.549
Pre-transplant positive serostatus [n (%)]			
Hepatitis C virus	6 (9.8)	1 (1.6)	0.125
Hepatitis B virus (surface antigen)	2 (3.3)	4 (6.6)	0.625
Epstein-Barr virus (anti-EBNA) ^d	49 (87.5)	47 (83.9)	0.754
CMV ^e	45 (73.8)	45 (75.0)	1.000
Pre-transplant maintenance dialysis [n (%)]	55 (90.2)	54 (88.5)	1.000
Duration, months [median (IQR)]	23 (15 - 41)	19.5 (12 - 45.8)	1.000

CMV: cytomegalovirus; EBNA: Epstein-Barr virus nuclear antigen; HBc: hepatitis B core antigen; ICU: intensive care unit; IPA: invasive pulmonary aspergillosis; IQR: interquartile range; SD: standard deviation.

^a Significant *P*-values (<0.05) are expressed in bold.

^b Data available for 58 cases and 57 controls.

^c Data available for 43 cases and 43 controls.

^d Data available for 56 cases and 56 controls.

^e Data available for 61 cases and 60 controls.

Table 2. Comparison of donor- and transplant-related factors, post-transplant events and outcomes.

Variable	Late IPA group (n = 61)	Control group (n = 61)	P-value ^a
Age of donor, years [mean ± SD]	49.8 ± 16.3	46.8 ± 13.5	0.283
Living donor [n (%)]	12 (19.7)	12 (19.7)	1.000
Double kidney transplantation [n (%)]	3 (4.9)	0 (0.0)	0.250
Induction therapy [n (%)] ^b			
None	22 (36.7)	20 (33.9)	1.000
Anti-CD25 (basiliximab or daclizumab)	22 (36.7)	20 (33.9)	0.815
Anti-thymocyte globulin	16 (26.7)	19 (32.2)	0.648
Primary immunosuppression regimen including [n (%)] ^b			
Steroids	54 (88.5)	57 (93.4)	0.375
Tacrolimus	29 (48.3)	30 (50.8)	1.000
Cyclosporine	19 (31.7)	20 (33.9)	1.000
MMF / MPA	47 (78.3)	50 (84.7)	0.375
Azathioprine	5 (8.5)	7 (11.9)	0.375
mTOR inhibitor	6 (10.0)	2 (3.4)	0.219
Length of hospital admission for transplantation, days [median (IQR)]	12 (8 - 18.8)	11 (6.3 - 18.8)	0.314
Delayed graft function [n (%)]	13 (21.3)	8 (13.1)	0.388
Surgical reintervention [n (%)] ^c	6 (10.2)	2 (3.7)	0.687
eGFR at month 3 after transplantation, mL/min/1.72 m ² [mean ± SD] ^d	23.8 ± 3.2	25.6 ± 3.4	0.873
eGFR at month 6 after transplantation, mL/min/1.72 m ² [mean ± SD] ^e	22.9 ± 3.1	20.5 ± 2.8	0.159
Leukopenia (<3.0 × 10 ⁹ cells/L) [n (%)] ^{f,g}	10 (16.9)	6 (10.2)	0.388
Neutropenia (<1.5 × 10 ⁹ cells/L) [n (%)] ^{f,h}	6 (12.2)	3 (6.2)	0.687
Serum IgG levels, mg/dL [mean ± SD] ⁱ	879 ± 627	763 ± 571	0.750
Post-transplant events within the previous 6 months [n (%)] ^j			
IRE ^{k,l}	21 (34.4)	2 (3.3)	0.000
CMV disease	10 (16.4)	1 (1.6)	0.004
Non ventilator-associated pneumonia	9 (14.8)	1 (1.6)	0.021
De novo malignancy ^m	5 (8.2)	0 (0.0)	0.063
Laboratory-confirmed respiratory tract viral infection ⁿ	5 (8.2)	0 (0.0)	0.063
Bloodstream infection ^o	7 (11.5)	0 (0.0)	0.016
ICU admission for ≥72 hours	2 (3.3)	0 (0.0)	0.500
Acute graft rejection	4 (6.6)	5 (8.2)	1.000
Episode treated with steroid boluses	4 (4.9)	5 (8.2)	0.687
Overall mortality [n (%)]	29 (47.5)	0 (0.0)	0.001

IPA-attributable mortality [n (%)]

13 (21.3)

-

NA

CMV: cytomegalovirus; eGFR: estimated glomerular filtration rate; ICU: intensive care unit; **IgG**: immunoglobulin G; IPA: invasive pulmonary aspergillosis; IQR: interquartile range; IRE: immunosuppression-related event; MMF / MPA: mofetil mycophenolate / mycophenolate acid; mTOR: mammalian target of rapamycin; **NA**: not applicable; SD: standard deviation.

^a Significant *P*-values (<0.05) are expressed in bold.

^b Data available for 60 cases and 59 controls.

^c Data available for 59 cases and 54 controls.

^d Data available for 56 cases and 56 controls.

^e Data available for 54 cases and 54 controls.

^f At any point during the first 6 months after transplantation.

^g Data available for 59 cases and 59 controls.

^h Data available for 49 cases and 48 controls.

ⁱ Serum IgG levels measured within the 6-month period prior to or following the date of diagnosis of IPA (for cases) or the analogous "pseudo-date" of diagnosis (for controls). Data available for 10 cases and 4 controls.

^j Events occurring within the 6-month period prior to the date or the "pseudo-date" of diagnosis of IPA.

^k The total number of IREs may be less than the sum of each conditions since more than one event was consecutively present in some patients.

^l There were 3 cases of post-transplant tuberculosis, although none of them occurred within the 6-month period prior to the date or the "pseudo-date" of diagnosis of IPA.

^m Includes PTLN (3 cases), colorectal adenocarcinoma and metastatic adenocarcinoma of unknown primary origin (one case each).

ⁿ Includes influenza virus infection (4 cases).

^o Includes BSI due Enterobacteriaceae (3 cases), *Pseudomonas aeruginosa*, *Staphylococcus epidermidis*, *S. aureus* and *Candida albicans* (one case each).

Table 3. Uni- and multivariable analyses (conditional logistic regression) of risk factors predicting the occurrence of late IPA.

Variables	Univariate analysis			Multivariate analysis		
	OR	95% CI	P-value	OR	95% CI	P-value
Age at transplantation, years ^a	1.04	1.01 - 1.08	0.017	-	-	-
Diabetic nephropathy	3.00	0.97 - 9.30	0.057	-	-	-
Pre-transplant COPD	65.29	0.51 - 8324.28	0.091	-	-	-
Prior IRE ^{b,c}	20.00	2.68 - 149.02	0.003	19.26	2.07 - 179.46	0.009
Prior BSI ^b	7.00	0.86 - 56.89	0.069	-	-	-

BSI: bloodstream infection; CI: confidence interval; COPD: chronic obstructive pulmonary disease; IPA: invasive pulmonary aspergillosis; IRE: immunosuppression-related event; OR: odds ratio.

^a OR per unitary increment.

^b Events occurring within the 6 months previous to the date of diagnosis of IPA for cases or the analogous "pseudo-date of diagnosis" for corresponding controls.

^c Includes non-ventilator-associated pneumonia, CMV disease and post-transplant *de novo* malignancy.

474 **Figure legend**

- 475 • **Figure 1.** Temporal distribution of cases of late invasive pulmonary aspergillosis occurring
476 according to post-transplant month of diagnosis.

477 **Supporting Information**

478 Additional Supporting Information may be found in the online version of this article:

- 479 • **Supplementary Materials and Methods:** Definitions used for date of IPA diagnosis, IPA-
480 attributable diagnosis, CMV disease, tuberculosis, pneumonia, respiratory tract viral
481 infection, BSI, PTLD, delayed graft function, acute graft rejection and eGFR.

Figure 1.

