

# Quizzes as a measure of instructors' style of teaching

Elena Shevchenko

## Introduction

Quizzes is an essential part of students' academic assessment, which can be used to measure growth in knowledge and skills. However, education is interaction between students and instructor and, therefore, success or failure of a student is related to instructor's method of teaching. Some instructors choose conservative approaches while others prefer to use loyal methods of teaching. These method may impact students' ability to understand information and develop skills, where quizzes show quantitative assessment of these processes. Thus, quizzes also can be considered as a measure of instructors' style of teaching. In this paper instructors' style of teaching will be explored based on the results of quizzes.

## Data

The data includes 293 average quiz scores for 5 weekly quizzes in Statistics and Probability course and exam results for the same course, which was conducted after the last quiz. Each student was assigned to one instructor (A, B, C or D). Instructor A was assigned to 69 students, instructor B - 64, instructor C - 90 and instructor D - 70. Additionally, students were divided into groups (1 through 13). Most of the groups contain 23 students, while the others contain 18, 22 or 24 students. Data set also includes extra information that can be used in further analysis and will not be discussed in this paper. Quizzes were graded by each instructors, while exams were graded by other experts where exams were assigned randomly to different graders.

## EDA

Descriptive statistics for each variable is presented below. Quiz grades scale is 0-10 with almost equal mean and median.

	<b>group</b>	<b>quiz_average</b>	<b>Exam_1</b>	<b>Total</b>	<b>Grade</b>
count	281.000000	281.000000	281.000000	281.000000	281.000000
mean	7.209964	5.988968	74.227758	104.172598	71.892981
std	3.667722	2.359612	22.395585	30.204061	20.999297
min	1.000000	0.400000	10.000000	32.000000	17.495500
25%	4.000000	4.200000	59.000000	83.000000	57.173500
50%	7.000000	5.800000	78.000000	110.000000	76.341000
75%	10.000000	7.800000	93.000000	128.000000	89.332000
max	13.000000	10.000000	105.000000	155.000000	104.167500

Figure 1: Descriptive statistics

It can be seen from the table below that instructor A has the lowest quiz average , while instructor C has the highest quiz average. Immediate conclusion is that teaching methods of instructor A do not work on the students, while instructor C has more efficient approaches in teaching their students. However, there may be some other explanations of such results. One is that instructor C is more loyal in grading than instructor A. Another explanation is that it might happened that instructor A had students with lower level of mathematical skills who did not have experience in statistics before, while instructor C had more experienced students in this area. To control for difference between students' abilities to learn the material SAT scores can be considered. However, no extra data is available to instructors of this course. Moreover, the number of students assigned to each instructor and randomization of assignment allow to conclude that students should be approximately equally distributed between instructors according to their maths abilities.

<b>quiz_average</b>	
<b>instructor</b>	
A	3.993846
B	5.216949
C	8.086667
D	5.786567

Figure 2: Quiz average by instructor

Figure 3 shows that even though quizzes average was the lowest for instructor A, exams average is not the lowest among instructors. Additionally, instructor C has the highest exam average. The difference in exam grades for instructors A, C and D is negligible, while instructor B has significantly low exam average.

<b>Exam_1</b>	
<b>instructor</b>	
A	75.276923
B	65.966102
C	77.300000
D	76.358209

Figure 3: Exam average by instructor

Boxplots presented below show that exam grades for three instructors A, C and D look similar. Boxplot for instructor B is different from others and the box is located slightly lower than boxes for other instructors. Half of students for all three instructors received grades between 55 and 95, while for instructor B half of students received grades between 50 and 80. It is clear that the minimum exam grade is the highest for instructor A. The range for each instructors' grade is presented below two boxplots and shows that instructor A has the lowest range, while instructor C has the highest range. Standard deviation is also lowest for instructor A, while instructor D has the highest standard deviation.




Figure 4: Quiz averages, boxplot




Figure 5: Exam grades, boxplot

<b>Exam_1</b>	
<b>instructor</b>	
A	77
B	88
C	95
D	90

Figure 6: Range

		<b>instructor</b>
<b>Exam_1</b>	A	21.529419
	B	22.176382
	C	22.256144
	D	22.330643

Figure 7: Standard deviation

## Analysis

ANOVA is an appropriate analysis here since the interest is to test whether there is difference between quiz averages among instructors controlling for exams that students received. To use ANOVA it is necessary that the relationship between independent and dependent variables is linear. Both Figure 6 and correlation matrix show that there is linear positive relationship between two variables. Correlation matrix by instructor shows that the highest linear relationship between quiz grades and exam grades is for instructor A, while the lowest correlation coefficient is for instructor B.




Figure 8: Scatterplot

	quiz_average	Exam_1
quiz_average	1.000000	0.513822
Exam_1	0.513822	1.000000

Figure 9: Correlation matrix

		quiz_average	Exam_1
instructor			
A	quiz_average	1.000000	0.711916
	Exam_1	0.711916	1.000000
B	quiz_average	1.000000	0.568907
	Exam_1	0.568907	1.000000
C	quiz_average	1.000000	0.658733
	Exam_1	0.658733	1.000000
D	quiz_average	1.000000	0.609444
	Exam_1	0.609444	1.000000

Figure 10: Correlation matrix by instructor

Figure 11 presents the results of ANOVA. The conclusion of ANOVA is that means of all quiz averages are not all equal. Post hoc test TukeyHSD is conducted to determine pairs that are not equal. According to Figure 12, all pairs of means can be considered equal except pair for instructor B and D.

	sum_sq	df	F	PR(>F)
<b>instructor</b>	<b>692.673301</b>	<b>3.0</b>	<b>73.82737</b>	<b>4.065494e-35</b>
<b>Residual</b>	<b>866.302500</b>	<b>277.0</b>	<b>NaN</b>	<b>NaN</b>

Figure 11: ANOVA

```

Multiple Comparison of Means - Tukey HSD, FWER=0.05
=====
group1 group2 meandiff p-adj    lower    upper  reject
-----
      A      B   1.2231  0.001   0.4011   2.0451   True
      A      C   4.0928  0.001   3.3488   4.8369   True
      A      D   1.7927  0.001   0.9969   2.5885   True
      B      C   2.8697  0.001    2.104   3.6354   True
      B      D   0.5696  0.2737  -0.2465   1.3857  False
      C      D  -2.3001  0.001  -3.0377  -1.5625   True
-----

```

Figure 12: TukeyHSD post hoc test