

A weighted principal curve approach to inferring 3D chromatin architecture

Elena Tuzhilina


(joint work with Trevor Hastie and Mark Segal)

Stanford University, Department of Statistics

elenatuz@stanford.edu

November 21, 2019


Chromatin



Chromatin = DNA + nucleosomes

① DNA


Chromatin



Chromatin = DNA + nucleosomes

- 1 DNA
- 2 'Beads-on-a-string'


Chromatin



Chromatin = DNA + nucleosomes

- 1 DNA
- 2 'Beads-on-a-string'
- 3 Chromatin fiber


Chromatin



Chromatin = DNA + nucleosomes

- ① DNA
- ② 'Beads-on-a-string'
- ③ Chromatin fiber
- ④ Chromatin loop


Chromatin



Chromatin = DNA + nucleosomes

- ① DNA
- ② 'Beads-on-a-string'
- ③ Chromatin fiber
- ④ Chromatin loop
- ⑤ Chromosome


Contact matrix



Terminology

- genomic locus = 'piece'
- resolution = 'size of a piece'
- contact (formaldehyde + cross-linking + sequencing)

Contact matrix




Terminology

- genomic locus = 'piece'
- resolution = 'size of a piece'
- contact (formaldehyde + cross-linking + sequencing)

Notations

- $n = \#$ genomic loci
- $C_{ij} = \#$ contacts between loci i and j

Contact matrix



Terminology

- genomic locus = 'piece'
- resolution = 'size of a piece'
- contact (formaldehyde + cross-linking + sequencing)



Notations

- $n = \#$ genomic loci
- $C_{ij} = \#$ contacts between loci i and j

Contact matrix $C = [C_{ij}] \in \mathbb{Z}_+^{n \times n}$

Chromatin reconstruction problem

Goal: Use the information contained in C to reconstruct the locus spatial coordinates $x_1, \dots, x_n \in \mathbb{R}^3$.



Main ingredients

- loss function $\ell(x_1, \dots, x_n)$
- optimization problem minimizing/maximizing $\ell(x_1, \dots, x_n)$
w.r.t. $x_1, \dots, x_n \in \mathbb{R}^3$

Main ingredients

- loss function $\ell(x_1, \dots, x_n)$
- optimization problem minimizing/maximizing $\ell(x_1, \dots, x_n)$
w.r.t. $x_1, \dots, x_n \in \mathbb{R}^3$

Example (Deterministic model, Metric MDS)

- 1 Convert C to a distance matrix D , e.g. $D_{ij} = \begin{cases} (C_{ij})^{-\alpha} & \text{if } C_{ij} > 0 \\ \infty & \text{if } C_{ij} = 0 \end{cases}$
- 2 Minimize Stress objective

$$\ell(x_1, \dots, x_n) = \sum_{i,j=1}^n W_{ij} (D_{ij} - \|x_i - x_j\|)^2$$

w.r.t. $x_1, \dots, x_n \in \mathbb{R}^3$

Main ingredients

- loss function $\ell(x_1, \dots, x_n)$
- optimization problem minimizing/maximizing $\ell(x_1, \dots, x_n)$
w.r.t. $x_1, \dots, x_n \in \mathbb{R}^3$

Example (Probability model, Poisson)

- 1 $C_{ij} \sim \text{Pois}(\lambda_{ij})$, where $\lambda_{ij} = \lambda_{ij}(x_1, \dots, x_n) = \beta \|x_i - x_j\|^\alpha$
- 2 Minimize negative log-likelihood

$$\ell(x_1, \dots, x_n) = \sum_{1 \leq i, j \leq n} \beta \|x_i - x_j\|^\alpha - C_{ij} \log(\beta \|x_i - x_j\|^\alpha)$$

w.r.t. $x_1, \dots, x_n \in \mathbb{R}^3$

(Previous approaches) Add a smoothness penalty!

$$\begin{aligned} \text{minimize } & \ell(x_1, \dots, x_n) + \lambda h(x_1, \dots, x_n) \\ \text{w.r.t. } & x_1, \dots, x_n \in \mathbb{R}^3 \end{aligned}$$

(Previous approaches) Add a smoothness penalty!

$$\begin{aligned} &\text{minimize } \ell(x_1, \dots, x_n) + \lambda h(x_1, \dots, x_n) \\ &\text{w.r.t. } x_1, \dots, x_n \in \mathbb{R}^3 \end{aligned}$$

Problems

- non convexity
- complexity of the objective, gradient and hessian

Controlling reconstruction smoothness

(Previous approaches) Add a smoothness penalty!

$$\begin{aligned} &\text{minimize } \ell(x_1, \dots, x_n) + \lambda h(x_1, \dots, x_n) \\ &\text{w.r.t. } x_1, \dots, x_n \in \mathbb{R}^3 \end{aligned}$$

Problems

- non convexity
- complexity of the objective, gradient and hessian

(Our approach) Add a constraint!

$$\begin{aligned} &\text{minimize } \ell(x_1, \dots, x_n) \text{ w.r.t. } x_1, \dots, x_n \in \mathbb{R}^3 \\ &x_1, \dots, x_n \in \text{smooth one-dimensional curve} \end{aligned}$$

Smooth curve constraint

① $x_1, \dots, x_n \in \gamma(t)$, where $\gamma(t) = \begin{pmatrix} \gamma_1(t) \\ \gamma_2(t) \\ \gamma_3(t) \end{pmatrix}$

Smooth curve constraint

- ① $x_1, \dots, x_n \in \gamma(t)$, where $\gamma(t) = \begin{pmatrix} \gamma_1(t) \\ \gamma_2(t) \\ \gamma_3(t) \end{pmatrix}$
- ② $h_1(t), \dots, h_k(t)$ – cubic spline basis functions, $\gamma_j(t) = \sum_{\ell=1}^k \Theta_{\ell j} h_{\ell}(t)$


k = spline degrees of freedom (df), hyperparameter

Smooth curve constraint

- ① $x_1, \dots, x_n \in \gamma(t)$, where $\gamma(t) = \begin{pmatrix} \gamma_1(t) \\ \gamma_2(t) \\ \gamma_3(t) \end{pmatrix}$
- ② $h_1(t), \dots, h_k(t)$ – cubic spline basis functions, $\gamma_j(t) = \sum_{\ell=1}^k \Theta_{\ell j} h_{\ell}(t)$
- ③ $x_i = \gamma(i) = \begin{pmatrix} \sum_{\ell=1}^k \Theta_{\ell 1} h_{\ell}(i) \\ \sum_{\ell=1}^k \Theta_{\ell 2} h_{\ell}(i) \\ \sum_{\ell=1}^k \Theta_{\ell 3} h_{\ell}(i) \end{pmatrix}$

Smooth curve constraint

- 1 $x_1, \dots, x_n \in \gamma(t)$, where $\gamma(t) = \begin{pmatrix} \gamma_1(t) \\ \gamma_2(t) \\ \gamma_3(t) \end{pmatrix}$
- 2 $h_1(t), \dots, h_k(t)$ – cubic spline basis functions, $\gamma_j(t) = \sum_{\ell=1}^k \Theta_{\ell j} h_{\ell}(t)$
- 3 $x_i = \gamma(i) = \begin{pmatrix} \sum_{\ell=1}^k \Theta_{\ell 1} h_{\ell}(i) \\ \sum_{\ell=1}^k \Theta_{\ell 2} h_{\ell}(i) \\ \sum_{\ell=1}^k \Theta_{\ell 3} h_{\ell}(i) \end{pmatrix}$



$$\longrightarrow H = \begin{pmatrix} | & & | \\ h_1 & \dots & h_k \\ | & & | \end{pmatrix} \in \mathbb{R}^{n \times k}$$

Smooth curve constraint

- ① $x_1, \dots, x_n \in \gamma(t)$, where $\gamma(t) = \begin{pmatrix} \gamma_1(t) \\ \gamma_2(t) \\ \gamma_3(t) \end{pmatrix}$
- ② $h_1(t), \dots, h_k(t)$ – cubic spline basis functions, $\gamma_j(t) = \sum_{\ell=1}^k \Theta_{\ell j} h_{\ell}(t)$
- ③ $x_i = \gamma(i) = \begin{pmatrix} \sum_{\ell=1}^k \Theta_{\ell 1} h_{\ell}(i) \\ \sum_{\ell=1}^k \Theta_{\ell 2} h_{\ell}(i) \\ \sum_{\ell=1}^k \Theta_{\ell 3} h_{\ell}(i) \end{pmatrix}$

$$X = \begin{pmatrix} -x_1^T - \\ \dots \\ -x_n^T - \end{pmatrix} = \begin{pmatrix} | & | & | \\ \gamma_1 & \gamma_2 & \gamma_3 \\ | & | & | \end{pmatrix} \in \mathbb{R}^{n \times 3} \quad H = \begin{pmatrix} | & & | \\ h_1 & \dots & h_k \\ | & & | \end{pmatrix} \in \mathbb{R}^{n \times k}$$

Smooth curve constraint

- ① $x_1, \dots, x_n \in \gamma(t)$, where $\gamma(t) = \begin{pmatrix} \gamma_1(t) \\ \gamma_2(t) \\ \gamma_3(t) \end{pmatrix}$
- ② $h_1(t), \dots, h_k(t)$ – cubic spline basis functions, $\gamma_j(t) = \sum_{\ell=1}^k \Theta_{\ell j} h_{\ell}(t)$
- ③ $x_i = \gamma(i) = \begin{pmatrix} \sum_{\ell=1}^k \Theta_{\ell 1} h_{\ell}(i) \\ \sum_{\ell=1}^k \Theta_{\ell 2} h_{\ell}(i) \\ \sum_{\ell=1}^k \Theta_{\ell 3} h_{\ell}(i) \end{pmatrix}$

$$X = \begin{pmatrix} -x_1^T - \\ \dots \\ -x_n^T - \end{pmatrix} = \begin{pmatrix} | & | & | \\ \gamma_1 & \gamma_2 & \gamma_3 \\ | & | & | \end{pmatrix} \in \mathbb{R}^{n \times 3} \quad H = \begin{pmatrix} | & & | \\ h_1 & \dots & h_k \\ | & & | \end{pmatrix} \in \mathbb{R}^{n \times k}$$

$x_1, \dots, x_n \in \text{smooth one-dimensional curve} \iff$

$\exists \Theta \in \mathbb{R}^{k \times 3}$ such that $X = H\Theta$

PCMS = Classical MDS + Smooth curve constraint

PCMS = Classical MDS + Smooth curve constraint

Classical MDS

$$\ell(x_1, \dots, x_n) = \sum_{i=1}^n \sum_{j=1}^n (C_{ij} - \langle x_i, x_j \rangle)^2 \iff \ell(X) = \|C - XX^T\|_F^2$$

PCMS = Classical MDS + Smooth curve constraint

Classical MDS

$$\ell(x_1, \dots, x_n) = \sum_{i=1}^n \sum_{j=1}^n (C_{ij} - \langle x_i, x_j \rangle)^2 \iff \ell(X) = \|C - XX^T\|_F^2$$

Smooth curve constraint $X = H\Theta$

PCMS = Classical MDS + Smooth curve constraint

Classical MDS

$$\ell(x_1, \dots, x_n) = \sum_{i=1}^n \sum_{j=1}^n (C_{ij} - \langle x_i, x_j \rangle)^2 \iff \ell(X) = \|C - XX^T\|_F^2$$

Smooth curve constraint $X = H\Theta$

PCMS optimization problem

$$\text{minimize } \ell_{PCMS}(\Theta) = \|C - H\Theta\Theta^T H^T\|_F^2 \text{ w.r.t. } \Theta \in \mathbb{R}^{k \times 3}$$

PCMS = Classical MDS + Smooth curve constraint

Classical MDS

$$\ell(x_1, \dots, x_n) = \sum_{i=1}^n \sum_{j=1}^n (C_{ij} - \langle x_i, x_j \rangle)^2 \iff \ell(X) = \|C - XX^T\|_F^2$$

Smooth curve constraint $X = H\Theta$

PCMS optimization problem

$$\text{minimize } \ell_{PCMS}(\Theta) = \|C - H\Theta\Theta^T H^T\|_F^2 \text{ w.r.t. } \Theta \in \mathbb{R}^{k \times 3}$$

Solution via Eigen Decomposition of $H^T C H$

PCMS = Classical MDS + Smooth curve constraint

Classical MDS

$$\ell(x_1, \dots, x_n) = \sum_{i=1}^n \sum_{j=1}^n W_{ij} (C_{ij} - \langle x_i, x_j \rangle)^2 \iff \ell(X) = \|\sqrt{W} * (C - XX^T)\|_F^2$$

Smooth curve constraint $X = H\Theta$

PCMS optimization problem

$$\text{minimize } \ell_{PCMS}(\Theta) = \|\sqrt{W} * (C - H\Theta\Theta^T H^T)\|_F^2 \text{ w.r.t. } \Theta \in \mathbb{R}^{k \times 3}$$

Solution iterative algorithm using PCMS as a building block

Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

Transformation


$$C^{\log} = \frac{\log(C+\epsilon)-\beta}{\alpha} \text{ for } \epsilon = 0.001, \alpha = 1 \text{ and } \beta = \log\left(\frac{\sum_{i,j=1}^n C_{ij}}{n}\right)$$

PCMS examples ($df = 10$)


Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

Transformation


$$C^{\log} = \frac{\log(C+\epsilon)-\beta}{\alpha} \text{ for } \epsilon = 0.001, \alpha = 1 \text{ and } \beta = \log\left(\frac{\sum_{i,j=1}^n C_{ij}}{n}\right)$$



Original data



Reconstruction




3D conformation

PCMS examples ($df = 25$)


Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

Transformation


$$C^{\log} = \frac{\log(C+\epsilon)-\beta}{\alpha} \text{ for } \epsilon = 0.001, \alpha = 1 \text{ and } \beta = \log\left(\frac{\sum_{i,j=1}^n C_{ij}}{n}\right)$$



Original data



Reconstruction




3D conformation

PCMS examples ($df = 50$)


Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

Transformation


$$C^{\log} = \frac{\log(C+\epsilon)-\beta}{\alpha} \text{ for } \epsilon = 0.001, \alpha = 1 \text{ and } \beta = \log\left(\frac{\sum_{i,j=1}^n C_{ij}}{n}\right)$$



Original data



Reconstruction



3D conformation

PoisMS = Weighted PCMS + Poisson Model

PoisMS = Weighted PCMS + Poisson Model

Model $C_{ij} \sim \text{Pois}(\lambda_{ij}), \quad \log(\lambda_{ij}) = \alpha \langle x_i, x_j \rangle + \beta$

PoisMS = Weighted PCMS + Poisson Model

Model $C_{ij} \sim \text{Pois}(\lambda_{ij})$, $\log(\lambda_{ij}) = \alpha \langle x_i, x_j \rangle + \beta$

Negative log-likelihood

$$\ell_{\text{PoisMS}}(X) = \sum_{1 \leq i, j \leq n} \left[e^{\alpha \langle x_i, x_j \rangle + \beta} - C_{ij} (\alpha \langle x_i, x_j \rangle + \beta) \right]$$

PoisMS = Weighted PCMS + Poisson Model

Model $C_{ij} \sim \text{Pois}(\lambda_{ij})$, $\log(\lambda_{ij}) = \alpha \langle x_i, x_j \rangle + \beta$

Negative log-likelihood

$$\ell_{\text{PoisMS}}(X) = \sum_{1 \leq i, j \leq n} \left[e^{\alpha \langle x_i, x_j \rangle + \beta} - C_{ij} (\alpha \langle x_i, x_j \rangle + \beta) \right]$$

Smooth curve constraint $X = H\Theta$

PoisMS = Weighted PCMS + Poisson Model

Model $C_{ij} \sim \text{Pois}(\lambda_{ij})$, $\log(\lambda_{ij}) = \alpha \langle x_i, x_j \rangle + \beta$

Negative log-likelihood

$$\ell_{\text{PoisMS}}(X) = \sum_{1 \leq i, j \leq n} \left[e^{\alpha \langle x_i, x_j \rangle + \beta} - C_{ij} (\alpha \langle x_i, x_j \rangle + \beta) \right]$$

Smooth curve constraint $X = H\Theta$

PoisMS optimization problem

minimize $\ell_{\text{PoisMS}}(X)$ w.r.t. X subject to $X = H\Theta$

PoisMS = Weighted PCMS + Poisson Model

Model $C_{ij} \sim \text{Pois}(\lambda_{ij})$, $\log(\lambda_{ij}) = \alpha \langle x_i, x_j \rangle + \beta$

Negative log-likelihood

$$\ell_{\text{PoisMS}}(X) = \sum_{1 \leq i, j \leq n} \left[e^{\alpha \langle x_i, x_j \rangle + \beta} - C_{ij} (\alpha \langle x_i, x_j \rangle + \beta) \right]$$

Smooth curve constraint $X = H\Theta$

PoisMS optimization problem

minimize $\ell_{\text{PoisMS}}(X)$ w.r.t. X subject to $X = H\Theta$

Solution use PCMS as a building block of the iterative algorithm!

PoisMS iterative algorithm

$$\ell_{PoisMS}(X) \approx \ell_{SOA}(X) = \|\sqrt{W} * (Z - X^T X)\|_F^2$$

$$\text{where } W = e^{\alpha X_0 X_0^T + \beta} \quad \text{and} \quad Z = X_0 X_0^T + \frac{1}{\alpha} \frac{C - W}{W}$$

PoisMS iterative algorithm

$$\ell_{PoisMS}(X) \approx \ell_{SOA}(X) = \|\sqrt{W} * (Z - X^T X)\|_F^2$$

$$\text{where } W = e^{\alpha X_0 X_0^T + \beta} \quad \text{and} \quad Z = X_0 X_0^T + \frac{1}{\alpha} \frac{C - W}{W}$$

\Rightarrow **can use Weighted PCMS to optimize SOA loss**

PoisMS iterative algorithm

$$\ell_{\text{PoisMS}}(X) \approx \ell_{\text{SOA}}(X) = \|\sqrt{W} * (Z - X^T X)\|_F^2$$

$$\text{where } W = e^{\alpha X_0 X_0^T + \beta} \quad \text{and} \quad Z = X_0 X_0^T + \frac{1}{\alpha} \frac{C - W}{W}$$

\Rightarrow can use **Weighted PCMS** to optimize **SOA** loss

1 **[Initialize]** Generate random $\Theta \in \mathbb{R}^{k \times 3}$

PoisMS iterative algorithm

$$\ell_{\text{PoisMS}}(X) \approx \ell_{\text{SOA}}(X) = \|\sqrt{W} * (Z - X^T X)\|_F^2$$

$$\text{where } W = e^{\alpha X_0 X_0^T + \beta} \quad \text{and} \quad Z = X_0 X_0^T + \frac{1}{\alpha} \frac{C - W}{W}$$

\Rightarrow **can use Weighted PCMS to optimize SOA loss**

❶ **[Initialize]** Generate random $\Theta \in \mathbb{R}^{k \times 3}$

❷ *Repeat until convergence:*

PoisMS iterative algorithm

$$\ell_{\text{PoisMS}}(X) \approx \ell_{\text{SOA}}(X) = \|\sqrt{W} * (Z - X^T X)\|_F^2$$

$$\text{where } W = e^{\alpha X_0 X_0^T + \beta} \quad \text{and} \quad Z = X_0 X_0^T + \frac{1}{\alpha} \frac{C - W}{W}$$

\Rightarrow can use **Weighted PCMS** to optimize **SOA** loss

① **[Initialize]** Generate random $\Theta \in \mathbb{R}^{k \times 3}$

② *Repeat until convergence:*

- **[SOA]** $\hat{C} = H\Theta\Theta^T H^T, \quad W = e^{\alpha \cdot \hat{C} + \beta}, \quad Z = \hat{C} + \frac{1}{\alpha} \left(\frac{C - W}{W} \right)$

PoisMS iterative algorithm

$$\ell_{\text{PoisMS}}(X) \approx \ell_{\text{SOA}}(X) = \|\sqrt{W} * (Z - X^T X)\|_F^2$$

$$\text{where } W = e^{\alpha X_0 X_0^T + \beta} \quad \text{and} \quad Z = X_0 X_0^T + \frac{1}{\alpha} \frac{C - W}{W}$$

\Rightarrow can use **Weighted PCMS** to optimize **SOA** loss

① **[Initialize]** Generate random $\Theta \in \mathbb{R}^{k \times 3}$

② *Repeat until convergence:*

- **[SOA]** $\hat{C} = H\Theta\Theta^T H^T$, $W = e^{\alpha \cdot \hat{C} + \beta}$, $Z = \hat{C} + \frac{1}{\alpha} \left(\frac{C - W}{W} \right)$
- **[WPCMS]** $\Theta := \text{PCMS}_W(Z, H)$

PoisMS examples

Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

PoisMS examples


Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

PoisMS parameters: $\alpha = 1$ and $\beta = \log \left(\frac{\sum_{i,j=1}^n C_{ij}}{n} \right)$


PoisMS examples ($df = 10$)

Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.


PoisMS parameters: $\alpha = 1$ and $\beta = \log \left(\frac{\sum_{i,j=1}^n C_{ij}}{n} \right)$



Original data



Reconstruction




3D conformation


PoisMS examples ($df = 25$)

Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

PoisMS parameters: $\alpha = 1$ and $\beta = \log \left(\frac{\sum_{i,j=1}^n C_{ij}}{n} \right)$



Original data



Reconstruction




3D conformation


PoisMS examples ($df = 50$)

Data: Hi-C data for IMR90 cells from the Gene Expression Omnibus, chromosome 20, probe resolution 100kb, $n = 625$.

PoisMS parameters: $\alpha = 1$ and $\beta = \log \left(\frac{\sum_{i,j=1}^n C_{ij}}{n} \right)$



Original data



Reconstruction



3D conformation

References



T. Stevens et al. *3D structures of individual mammalian genomes studied by single-cell Hi-C*. Nature, 544:59–64, 2017.



N. Varoquaux et al. *A statistical approach for inferring the three-dimensional structure of the genome*. Bioinformatics, 30(12):26-33, 2014.



Z. Zhang et al. *3D Chromosome Modeling with Semi-Definite Programming and Hi-C Data*. Journal of computational biology, 20(11):831–846, 2013.



C. Zou et al. *HSA: integrating multi-track Hi-C data for genome-scale reconstruction of 3D chromatin structure*. Genome Biology, 17(40), 2016.



M. Rosenthal et al. *Bayesian Estimation of 3D Chromosomal Structure from Single Cell Hi-C Data*. Journal of Computational Biology, published online, 2019.



A. Buja et al. *Data Visualization With Multidimensional Scaling*. Journal of Computational and Graphical Statistics, 17(2):444-472, 2008.



R. Mazumder et al. *Spectral Regularization Algorithms for Learning Large Incomplete Matrices*. Journal of Machine Learning Research, 11: 2287-2322, 2010.



N. Srebro et al. *Weighted Low-Rank Approximations*. Proceedings of the Twentieth International Conference on Machine Learning (ICML-2003), Washington DC, 2003.

Thank you for your attention!