

# Canonical Correlation Analysis in high dimensions with structured regularization

Elena Tuzhilina

(joint work with Trevor Hastie and Leonardo Tozzi)



Stanford University, Department of Statistics

*elenatuz@stanford.edu*

August 12, 2021


# Data

- ① **brain activations:** magnetic resonance imaging obtained during a gambling task designed to probe the brain circuits underlying reward
- ② **behavioral performance measures:** self-reports assessing various aspects of reward-related behaviors, depression symptoms and positive as well as negative affective states




**Question:** is there any correlation between brain activity and behavioral measures of performance during the gambling tasks?

# Notations



$X \in \mathbb{R}^{n \times p}$  – brain activations



$Y \in \mathbb{R}^{n \times q}$  – behavior scores

## Dimensions:

- $n = 153$  participants
- $p = 90,368$  greyordinates
- $q = 9$  scores

# Canonical Correlation Analysis

**Goal:** find  $\alpha \in \mathbb{R}^p, \beta \in \mathbb{R}^q$  such that

$$\text{maximize } \rho(\alpha, \beta) = \text{cor}(X\alpha, Y\beta) = \frac{\alpha^T \Sigma_{XY} \beta}{\sqrt{\alpha^T \Sigma_{XX} \alpha} \sqrt{\beta^T \Sigma_{YY} \beta}}$$

- canonical coefficients  $\alpha$  and  $\beta$
- canonical correlation  $\rho(\alpha, \beta) = \text{cor}(u, v)$

Find maximum via Singular Value Decomposition of  $\Sigma_{XX}^{-\frac{1}{2}} \Sigma_{XY} \Sigma_{YY}^{-\frac{1}{2}}$

# Regularization

**Problem:**  $\Sigma_{XX}$  is singular when  $p > n$ , so CCA does not work.

**Solution:** adjust the diagonal of  $\Sigma_{XX}$ .

Modified correlation coefficient

$$\rho(\alpha, \beta) = \frac{\alpha^T \Sigma_{XY} \beta}{\sqrt{\alpha^T (\Sigma_{XX} + \lambda_1 I) \alpha} \sqrt{\beta^T \Sigma_{YY} \beta}}$$

Find maximum via Singular Value Decomposition of  $(\Sigma_{XX} + \lambda_1 I)^{-\frac{1}{2}} \Sigma_{XY} \Sigma_{YY}^{-\frac{1}{2}}$

**RCCA shrinkage property:**  $\lambda_1 \uparrow$  shrinks  $\alpha \rightarrow 0$ , similar to ridge regression.

# Kernel trick

```
library(CCA)
rcc(X = activation, Y = behavior, lambda1 = 10, lambda2 = 0)
```

Error: cannot allocate vector of size 62.1 Gb  
Traceback:

1. rcc(X = activation, Y = behavior, lambda1 = 10, lambda2 = 0)
2. var(X, na.rm = TRUE, use = "pairwise")


**Problem:**  $\Sigma_{XX}$  is  $p \times p$ , so cannot store for  $p \approx 90K$ .

**Solution:** find a linear transformation such that RCCA for  $(X, Y)$  is equivalent to RCCA for  $(R, Y)$  and

$$V = \begin{array}{|c|} \hline p \times n \\ \hline \end{array} \quad R = XV = \begin{array}{|c|} \hline n \times p \\ \hline \end{array} \begin{array}{|c|} \hline p \times n \\ \hline \end{array} = \begin{array}{|c|} \hline n \times n \\ \hline \end{array}$$


# RCCA preformance

RCCA  
max correlation = 0.11 for  $\lambda_1=0.001$




# RCCA interpretability

**Visualization:** plot canonical coefficients  $\alpha$  for the optimal RCCA model.



(a) Cortical coefficients.




(b) Subcortical coefficients.


**Problem:** There is a lot of variation, so this plot is hard to interpret.

# Group structure

**Motivation:** brain features come in groups (aka brain regions). How to take into account the group structure?



(a) Cortical parcellation (210 regions).



(b) Subcortical parcellation (19 regions).

# Grouped structure

## Block structure:

$$X = (\underbrace{X_1}_{region_1}, \dots, \underbrace{X_K}_{region_K}) \text{ and } \alpha = (\underbrace{\alpha_1}_{region_1}, \dots, \underbrace{\alpha_K}_{region_K})$$

### Assumptions:

- ① group homogeneity  $\alpha_k \approx \bar{\alpha}_k$
- ② sparsity on a group level  $\bar{\alpha}_k \approx 0$


$$C = \begin{bmatrix} \frac{11^T}{p_1} & 0 & \dots & 0 \\ 0 & \frac{11^T}{p_2} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \frac{11^T}{p_K} \end{bmatrix}$$

Modified correlation coefficient

$$\rho(\alpha, \beta) = \frac{\alpha^T \Sigma_{XY} \beta}{\sqrt{\alpha^T (\Sigma_{XX} + \lambda_1(I - C) + \mu_1 C) \alpha} \sqrt{\beta^T \Sigma_{YY} \beta}}$$


**GRCCA shrinkage property:**  $\lambda_1 \uparrow$  shrinks  $\alpha_k \rightarrow \bar{\alpha}_k$ ,  
 $\mu_1 \uparrow$  shrinks  $\bar{\alpha}_k \rightarrow 0$ .

# GRCCA performance




# GRCCA interpretability

**Visualization:** plot canonical coefficients  $\alpha$  for the optimal GRCCA model.



(a) Cortical coefficients.



(b) Subcortical coefficients.

# References

- D. M. BARCH, G. C. BURGESS, M. P. HARMS, S. E. PETERSEN, B. L. SCHLAGGAR, M. CORBETTA, M. F. GLASSER, S. CURTISS, S. DIXIT, C. FELDT, D. NOLAN, E. BRYANT, T. HARTLEY, O. FOOTER, J. M. BJORK, R. PODRACK, S. SMITH, H. JOHANSEN-BERG, A. Z. SNYDER, AND D. C. VAN ESSEN, *Function in the human connectome: Task-fMRI and individual differences in behavior*, *NeuroImage*, 80 (2013), pp. 169–189.
- D. C. V. ESSEN, S. M. SMITH, D. M. BARCH, T. E. BEHRENS, E. YACOUB, K. UGURBIL, AND W.-M. H. CONSORTIUM, *The WU-Minn Human Connectome Project: an overview*, *Neuroimage*, 80 (2013), pp. 62–79.
- M. F. GLASSER, T. S. COALSON, E. C. ROBINSON, C. D. HACKER, J. HARWELL, E. YACOUB, K. UGURBIL, J. ANDERSSON, C. F. BECKMANN, M. JENKINSON, S. M. SMITH, AND D. C. V. ESSEN, *A multi-modal parcellation of human cerebral cortex*, *Nature*, 536 (2016), pp. 171–178.
- I. GONZALEZ, D. S. P. MARTIN, AND A. BACCINI, *CCA: An R Package to Extend Canonical Correlation Analysis*, *Journal of Statistical Software*, 23(12) (2008), pp. 1–14.
- H. HOTELLING, *Relations between two sets of variables*, *Biometrika*, 28 (1936), pp. 321—377.
- W. HÄRDLE AND L. SIMAR, *Applied Multivariate Statistical Analysis: Second edition*, Springer, 2007.
- S. LEURGANS, R. MOYEEED, AND B. SILVERMAN, *Canonical Correlation Analysis when the Data are Curves*, *Journal of the Royal Statistical Society*, 55(3) (1993), pp. 725—740.
- K. MARDIA, J. T. KENT, AND J. M. BIBBY, *Multivariate Analysis*, Academic Press, New York, NY, 1979.
- H. VINOD, *Canonical Ridge and Econometrics of Joint Production*, *Journal of Econometrics*, 4(2) (1976), pp. 147—166.
- S. WEINTRAUB, S. S. DIKME, R. K. HEATON, D. S. TULSKY, P. D. ZELAZO, P. J. BAUER, N. E. CARLOZZI, J. SLOTKIN, D. BLITZ, K. WALLNER-ALLEN, N. A. FOX, J. L. BEAUMONT, D. MUNGAS, C. J. NOWINSKI, J. RICHLER, J. A. DEOCAMPO, J. E. ANDERSON, J. J. MANLY, B. BOROSH, R. HAVLIK, K. CONWAY, E. EDWARDS, L. FREUND, J. W. KING, C. MOY, E. WITT, AND R. C. GERSHON, *Cognition assessment using the NIH Toolbox*, *Neurology*, 80 (2013), pp. S54–64.

Thank you for your attention!