the major caspases responsible for PIAS1 cleavage are caspase-3, -6 and -8 (Figure 4B and 4C, lanes 3, 6 and 8). By examining the size of these cleaved fragments, we noticed that caspase-3 mediated cleavage generates one N-terminal (Figure 4B, lane 3, 50 kDa) and one C-terminal (Figure 4C, lane 3, 25 kDa) fragments, caspase-6 mediated cleavage generates one N-terminal (Figure 4B, lane 6, 15 kDa) and two C-terminal (Figure 4C, lane 6, 70 kDa and 25 kDa) fragments, and the cleavage by caspase-8 leads to two N-terminal (Figure 4B, lane 6; 15 kDa and 50 kDa) and one C-terminal (Figure 4C, lane 8, 70 kDa) fragments. The 25 kDa fragment generated by caspase-3 and -6 (Figure 4C, lanes 3 and 6) matched with that observed in EBV-replicating

utans from humans, or vice versa, or could spread amongst orangutans in captivity or in the wild (Table 1). It was necessary to answer OIE and IUCN problem description questions separately for each translocation type (reinforcement, reintroduction, and wild-to-wild translocation), as each translocation type has a distinct set of risks and uncertainties. Element 3 required consideration of the conservation, policy and cultural contexts underpinning orangutan translocations, using published literature and author's unpublished data. Elements 1–3 are summarized in Table 2, a matrix of answers to the OIE and IUCN problem description questions. To exemplify how the problem description could inform decision making and action (element