

Memories of migrations past: Sociality and cognition in dynamic, seasonal environments

Eliezer Gurarie^{1,2,*}, Sriya Potluri¹, G. Christopher Cosner³, R. Stephen Cantrell³, William F. Fagan¹

¹*Department of Biology, University of Maryland, College Park, MD 20742, USA.*

²*Department of Environmental and Forest Biology, Syracuse, NY*

³*Department of Mathematics, The University of Miami, Coral Gables, FL 33146, USA.*

Correspondence*:
Eliezer Gurarie
egurarie@umd.edu

2 ABSTRACT

3 For full guidelines regarding your manuscript please refer to [Author Guidelines](#).

4 As a primary goal, the abstract should render the general significance and conceptual advance of the
5 work clearly accessible to a broad readership. References should not be cited in the abstract. Leave the
6 Abstract empty if your article does not require one, please see [Summary Table](#) for details according to
7 article type.

8 **Keywords:** keyword, keyword, keyword, keyword, keyword, keyword, keyword, keyword

1 INTRODUCTION

9 Seasonal migrations are widespread among terrestrial, aquatic, avian and invertebrate species
10 ([Dingle, 2014](#)). For many species, migration is an extremely successful strategy, allowing a far greater
11 number of individuals to inhabit landscapes which might not otherwise be able to support large
12 numbers year round ([Fryxell et al., 1988](#)). Essentially, the evolutionary stability of a migratory
13 strategy relies on the fitness benefits of accessing suitable seasonal resources, whether those are
14 for energetic gain, predator avoidance, or a suitable physical, biotic or social environment for
15 reproduction outweighing the energetic and survival related costs of migration ([Avgar et al., 2014](#)).

16 Proximate causes, drivers and mechanism can vary considerably across and even within species
17 ([Berthold, 1999](#); [Shaw, 2016](#)). Some migrants follow a “green wave” of spring vegetation as it
18 flowers across altitudinal or latitudinal gradients ([Bischof et al., 2012](#); [Kölzsch et al., 2015](#)). These
19 migrations can be considered “tactical”, as they can occur - as an extreme simplification - purely as
20 response to local conditions. Other migrants perform long-distance migrations in anticipation that
21 critical resources will be available at the time of arrival at the end point of migration ([Abrahms
22 et al., 2019](#)). This second behavior involves the greatest trade-off between the costs of migration
23 against the benefits of accessing resources, whether those are food, suitable habitat for breeding,
24 or refuge from predators, that are highly seasonal and localized. This approach can be considered
25 “strategic” in the sense that it is driven not by immediate cues but by an anticipation based on prior
26 experience ([Bracis and Mueller, 2017](#)).

Migratory species are often considered to be more vulnerable to environmental change, as a disruption in either of the seasonal ranges or along a migratory corridor can have significant negative impacts (Wilcove and Wikelski, 2008; Kauffman et al., 2021). On the other hand, it has been argued that migratory species might be more resilient to disruptions due to their wide-ranging mobility (Robinson et al., 2009). Clearly, the ability to perform a migration without local cues is only possible if the behavior is hard-programmed or remembered. On the other hand, a strictly programmed behavior can be maladaptive if conditions change. Because the scenarios underlying migration are manifold and complex, mathematical modeling may provide some insights and help clarify where, when, and under what conditions we might migration behavior to emerge, to be adaptive, to be maladaptive, or to collapse.

Diffusion-advection models have a long pedigree in animal movement modeling (Skellam, 1951; Turchin, 1998; Okubo and Levin, 2001). These models are grounded in the general idea that animal movements, somewhat like movements of physical particles, combine random (diffusive) components with directed (advective) components. While direct relationships between diffusion models and movement data are somewhat tenuous (Gurarie and Ovaskainen, 2011; Potts and Schlägel, 2020), as a theoretical tool for exploring processes they are invaluable for their versatility and the relative ease of numeric computation of the partial differential equations (PDEs) that describe them mathematically. Much theoretical and some applied work has been done on refining the basic assumptions of diffusion models, e.g. by including heterogeneity in populations (Skalski and Gilliam, 2003; Gurarie et al., 2009), fat-tailed dispersal kernels (Kot et al., 1996), non-linear or otherwise complex responses to resources and conspecifics. But perhaps the greatest difference between animals and randomly moving passive particles described by diffusion models is cognition, including social behavior and memory. Refinements to diffusion-advection equations have revealed conditions under which non-local information gathering (Fagan et al., 2017) and behavioral switching may confer foraging advantages (Fagan et al., 2019), in particular when resources are dynamic and patchy. While diffusion models that incorporate social movements (swarming, schooling, herds) have been well-studied (Okubo, 1986; Grünbaum, 1994; Mogilner and Edelstein-Keshet, 1999), the interacting role of memory and sociality in a specifically migratory framework has been relatively unexplored.

Here, we develop a social diffusion-advection model with memory to explore the resilience of a migratory population under various dynamic, seasonal resource distributions. In formulating the model, our goal was to identify the minimum set of movement and memory parameters required to generate an adaptive, migratory behavior. This includes the ability to learn to migrate from non-migratory initial conditions, simulating the release of naive animals in a seasonal environment (Jesmer et al., 2018), to lose the propensity to migrate if the resource distribution does not require it, also a commonly observed phenomenon (Wilcove and Wikelski, 2008), and to assess the resilience or fragility of a migratory population against changing resource distribution dynamics, including both stochasticity and trends in spatial and temporal distributions, mirroring the effects of climate change.

We anticipated that under many conditions a blending of *tactical* (i.e. direct response to resource availability or perception) and *strategic* (i.e. memory-driven and forward-thinking) behavior will help foragers navigate dynamic, seasonal environments. Over-reliance on either strategy should be maladaptive. We further anticipate that a shorter-term memory updating is needed to navigate trends in resource spatial distribution and temporal distribution (phenology), but that a longer-term

reference memory is needed to navigate resource distributions that are stochastic (Lin et al., 2021). Similarly, we anticipated that a balance between very low sociality and extreme sociality would lead to the most resilient migratory process.

2 METHODS

2.1 Memory movement model

In designing our study, our goal was to develop a minimal heuristic in which the following processes were explicitly modeled: (1) Random or exploratory movement, (2) attraction to resources, (3) a long-term (or *reference*) memory of large-scale movement behavior, (4) a short-term (or *working*) memory that updates movement behavior based on recent experience, and (5) some social aspect to the learned behavior.

A diffusion-advection equation provided a computationally efficient and versatile framework for examining just such a system. We consider a population moving in one dimension in a constrained domain D and distributing itself according to the following equation:

$$\frac{\partial u}{\partial t} = -\varepsilon \frac{\partial^2 u}{\partial x^2} + \alpha \frac{\partial}{\partial x} \left(u \frac{\partial h}{\partial x} \right) + \beta \frac{\partial}{\partial x} (v_s(u)) + v_m(t) \quad (1)$$

where u represents the population distributed in time and space. The first term is the diffusion term, capturing the fast time-scale exploration and “random” movements of individuals, with ε is the diffusion rate.

The second term represents the attraction to a dynamic resource h , with the proportionality of the advection to the gradient of the resource given by the parameter α (note, the population and resource distributions are functions of both space and time $u(x, t)$ and $h(x, t)$ - we omit the dependent variables in the notation for brevity). This is the well-studied standard chemotactic resource-following behavior. We borrow the general notation from our earlier related work (Fagan et al., 2017, 2019).

The third term captures the collective or social advection term of the population via a non-local, density dependent function $v_s(u, x)$. If this function takes the form of a convolution around a non-local kernel k , i.e. $v_s(u) = k(x) * u(x)$, and if that kernel is odd, an attractive or “swarming” behavior can be generated (Mogilner and Edelstein-Keshet, 1999). We use the kernel analyzed by Mogilner and Edelstein-Keshet (1999): $k(x) = \frac{x}{2\lambda^2} \exp(-x^2/2\lambda^2)$. The convolution of u with this kernel has the property of pushing the population in a positive direction when $x < \hat{u}$, and in a negative direction when $x > \hat{u}$. The parameter λ is a length scale of sociality - roughly one-half the size of the “swarm”, where β is a parameter that quantifies the overall strength of sociality.

Finally, the last term captures the direct advection that emerges from a memory-driven migratory behavior. This term evolves with a set of parameters θ_y that slowly change each year $y \in \{0, 1, 2, \dots\}$, i.e. the count of periods τ : $y = \lfloor t/\tau \rfloor$.

The migration speed is specified by six parameters θ : the timing of the start and duration of two anticipated seasons (nominally, winter and summer) $t_1, \Delta t_1, t_2, \Delta t_2$, and the spatial coordinates of

the population centroid for each season x_1 and x_2 . Thus the remembered migratory speed term is a simple step function given by:

$$v_m(t, \theta_y) = \begin{cases} 0; & t > t_1 \text{ and } t \leq t_1 + \Delta t_1 \\ s_{12}; & t > t_1 + \Delta t_1 \text{ and } t \leq t_2 \\ 0; & t > t_2 \text{ and } t \leq t_2 + \Delta t_2 \\ s_{21}; & t > t_2 + \Delta t_2 \text{ or } t \leq t_1 \end{cases} \quad (2)$$

where the migration speeds s_{12} and s_{21} from the respective ranges are set such that they arrive at x_1 at t_1 , leave at $t = t_1 + \Delta t_1$, arrive at x_2 at $t = t_2$, i.e. $s_{12} = \frac{x_2 - x_1}{t_2 - (t_1 + \Delta t_1)}$ and $s_{21} = \frac{x_1 - x_2}{t_1 - (t_2 - \tau + \Delta t_2)}$. This step-like migration function is a one-dimensional version of the migration parameters estimated in empirical studies for individuals (Gurarie et al., 2017) and, more relevantly, for populations (Gurarie et al., 2019) in empirical studies.

We consider these six parameters to be the known or remembered determinants of the migratory behavior, with an initial set θ_0 determining the reference migration behavior. This reference migration is updated each year by the experience of the population. To perform this updating, we estimate a new set of parameters $\widehat{\theta}_y$ after each year, and combine these new parameters with the reference parameters according to the following weighted mean:

$$\theta_{y+1} = \kappa^y \theta_o + (1 - \kappa^y) \widehat{\theta}_y$$

where each of the six parameters is updated identically. The estimates $\widehat{\theta}_y$ are obtained via a least-squares minimization of the migration track ($m(t, \theta) = \int_0^t v_m(t', \theta_y) \Delta t'$) against the spatial mean of the population process in year y (i.e. $\widehat{u}(t) = \int_X u_y(t, x) dx$). The parameter $\kappa \in (0, 1)$ captures the reliance on that long-term memory. When $\kappa = 0$, all of the actionable memory is from the preceding year. When $\kappa = 1$, the actionable memory is entirely the reference memory.

The model is confined to a one-dimensional bounded domain $[-\chi, \chi]$, with no flux outside of the boundaries i.e. $\partial u(-\chi, t)/\partial t = \partial u(\chi, t)/\partial t = 0$. As there are no birth or death processes, the total population remains fixed and constant, for convenience integrating to 1. Furthermore, the parameters remain constant throughout time, with no adaptation or mutation-selection process. Our interest is entirely in the ability of a fixed set of memory and movement parameters to navigate an intra- and interannually dynamic, seasonal environment.

2.2 Seasonal resource

We solved the model numerically on a spatial domain $x \in [-100, 100]$, and a periodicity $\tau = 100$ (i.e. ~years of 100 days). We were interested in an approximately periodic resource dynamic, i.e. one in which $h(x, t \approx h(x, t - \tau))$. We generated two types of resource distributions. A “non-surfable” resource (*island resource*), and weakly surfable resource (*drifting resource*). Both are characterized by a peak in time and space centered at m_x at m_t , and $-m_x$ at $\tau - m_t$ (for example, locations 30 and -30 at times 25 and 75, respectively). These pulses have a shared time scale of duration s_t and a spatial scale of extent s_x , the standard deviation in the time and space dimension respectively. The island resource is simply two uncorrelated bivariate normal distributions

$$h(x, t) = K (\Phi(m_x, m_t, s_x, s_t) + \Phi(-m_x, \tau - m_t, s_x, s_t))$$




Figure 1. Examples of various seasonal resource distribution functions, contrasting short duration, but wide pulses ($\sigma_t = 3, \sigma_x = 12$; left panels), long duration but spatially concentrated pulses ($\sigma_t = 12, \sigma_x = 3$; right panels), and isolated resource pulses (upper panels) from the weakly drifting resource (lower panels). The total amount of resource is identical across all scenarios. In the weakly drifting resources, the total amount is constant at all times, and uniform in the middle of the phase (time = 0, 50, 100).

where Φ is the bivariate Gaussian distribution function, and the normalizing constant K is selected such that the average total amount of resource throughout the year is 1, i.e. $\frac{1}{T} \int_T \int_X h(x, t) dx dt = 1$.

The drifting resource differs from the island resource in that the total amount of resource at any given time point is 1 ($\int_X h(x, t) dx = 1$). This property is attained by distributing the resource as a re-scaled beta distribution, where the shape and scale parameters vary sinusoidally in such a way as to make the standard deviations and means match the desired values of m_x, m_t, s_x, s_t (see appendix for details). Both types of resources are illustrated in figure 1.

Within a given year, the resource is entirely symmetric: $h_y(x, t) = h_y(-x, T - t)$. However, in scenarios exploring climate change we allow the peaks to vary with drift and stochasticity according to: $m_x(y) \sim N(\mu_x + \gamma_x y, \sigma_x)$ and $m_t(y) \sim N(\mu_t + \gamma_t y, \sigma_t)$, where the μ , γ and σ terms are the mean, slope and variance, respectively, for the location and time duration of the pulse. Thus, if $\gamma = 0$ and $\sigma = 0$, the conditions are constant across years and if $\gamma_x > 0$ there is a drift of the resource towards extremes, if $\beta_t < 0$ there is a shift towards earlier resource pulses. These trends model the pole-ward shift of peak resources and the earlier spring phenology occurring with a warming global climate. The spatial scales (s_x and s_t) remain constant in all of our simulations.

2.3 Metrics

The main metrics we are interested in are *migration mismatch*, *foraging efficiency* and *adaptation to trends*.

Migration mismatch captures the similarity between the migration phenology and the resource phenology. To do this, we compute the total mismatch, i.e. is a sum of the difference between the migration coefficients and the resource peaks. Spatial mismatch MM_x is given by the absolute difference between the migration targets and the resource peaks: $MM_x = |x_1 - m_x| + |x_2 + m_x|$. Temporal mismatch is the difference between the arrival time and the peak of the resource if arrival is post-peak, the difference between the departure time and the peak of the resource if departure is pre-peak, and 0 if the seasonal duration spans the peak, i.e. $MM_t = \max\{t_1 - m_1, m_1 - (t_1 + \Delta t_1), 0\} + \max\{t_2 - m_2, m_2 - (t_2 + \Delta t_2), 0\}$. Finally, the total mismatch is the sum of these: $TM = MM_x + MM_t$. A mismatch of less than 1 is essentially perfect, we consider a mismatch of 1-5 to be “good”, and beyond 50 the system can be said to have failed to keep track of the resource dynamics.

To quantify the foraging efficiency, i.e. the organisms’ ability to track the distribution of the resources over space and time, we use a continuous form of the Bhattacharyya coefficient (Bhattacharyya, 1943) which quantifies the similarity between two distributions. We compute this coefficient at every time point in a given year, and take the mean across the equilibrium year to determine foraging efficiency (FE). Thus, the foraging efficiency index is:

$$FE = \frac{1}{\tau} \int_0^\tau \int_{-\chi}^\chi \sqrt{u(x,t) h(x,t)} dx dt$$

where the spatial integral is taken over the domain. This metric is constrained to be between 0 and 1.

For simulations with a constant resource, we ran the model until a quasi-equilibrium (stationary) state was achieved, i.e. where the Bhattacharyya index of the population distribution across subsequent years reached a value of 0.9999. Once stationarity was attained, we computed the migration mismatch and foraging efficiency metrics, as well as the number of years required to reach stationarity.

For numerical runs with climate change, we first run a simulation with a given parameter set until stationarity, as above, and then begin shifting the location of the resource poleward with a slow, moderate or rapid trend ($\gamma_x = 0.25, 0.5$ and 1 , respectively), and / or by adding stochasticity (spatial standard deviation 3, 6, 9 or 12). For stochasticity analyses, we compare foraging efficiency across a range of the reference memory parameter κ . For analyses that include trend (with or without stochasticity), we quantified the ability of the system to keep track of climate change with a *spatial adaptation* (SA) index. This index is the ratio of the slope of the memory-based migration location over time (i.e. the fitted slope coefficient of $m_{x,i} = \hat{\gamma}_x i + m_{x,0}$, where i is the year) against the resource drift parameter γ_x , i.e. $SA = \hat{\gamma}_x / \gamma_x$. An SA equal to 1 suggests that the process is keeping up with climate change, an SA of 0 indicates that the process is not responding at all to climate change. Values greater than 1 (super-adaptation) are possible, as are values less than 1, which correspond to a loss of migration behavior.

All movement model parameters, resource parameters, and metrics are summarized in table 1.

2.4 Simulation studies

We explored this model using numerical differencing of a system of ordinary differential equations (ODE's) approximating the PDE in equation 1 with the Runge-Kutte algorithm using the `deSolve` (Soetaert et al., 2010) and `ReacTran` (Soetaert and Meysman, 2012) packages in R. We additionally used the `nlsLM` function in package `minpack.LM` (Elzhov et al., 2016) for robust and fast annual estimation of the migration parameters. The complete code is available as an R package (`memorymigration`) available on GitHub at <https://github.com/EliGurarie/memorymigration> and as an interactive Shiny application at <https://spot3512.shinyapps.io/memorymigrationshinyapp/>.

We assessed a wide range of parameter values and resource geometries and dynamics with the goal of answering the four main questions: (1) Can this model adapt to a discrete shift in peak resource location and timing? What is the relative role of memory and sociality for adaptation? (2) Can this model acquire a migratory behavior from a non-migratory initial condition? (3) What is the role of a reference memory for dealing with stochastic resource dynamics? (4) Can this model adapt when the resource peaks shifts in space?

3 RESULTS

3.1 Adaptation to resource phenology

The ability of this system to attain a stable, migratory state that matches the dynamics of the resource is illustrated in figure 2. In the illustrated scenario, it takes nearly 40 years to attain an equilibrium, and the eventual steady state is one where the centroid of the migration lines up exactly with the centroid of the resource, and the arrival timing coincides with the *peak* of resource

Table 1. Table of parameters, variables and metrics.

Memory migration model	
ϵ	Diffusion
α	Forage following
β	Strength of sociality
λ	Spatial scale of sociality
κ	Proportion of reference versus working memory
x_1, x_2	location of population centroids in summer and winter
$t_1, \Delta t_1$	start and duration of summer season
$t_2, \Delta t_2$	start and duration of winter season
κ	proportion memory allocated to reference
	(long-term) memory vs. working (short-term) memory
Resource dynamics	
τ	duration of period (year)
$m_x, -m_x$	spatial coordinate of resource peak for summer and winter
$m_t, \tau - m_t$	timing of resource peak for the summer and winter
σ_x, σ_t	time duration and and spatial scale of resource pulse
γ_x, γ_t	rate of change of peak location and timing of resource
ψ_x, ψ_t	standard deviation of peak location and timing
Metrics	
MM_x	spatial migration mismatch
MM_t	temporal migration mismatch
TM	total mismatch
FE	foraging efficiency
SA	spatial adaptation index




Figure 2. Example of adaptation to a shift in resource peak. The initial (year 0) behavior migrates to locations 50 and -50 at days 15 and 60, whereas the resource peak is at 30 and -30, peaking at times 25 and 75. The panels show (a) the first 14 years of the simulation; (b) the centroid of the annual movement of the population is shown in panel b, with dark blue to red colors indicating year 0 to year 40; (c) annual foraging efficiency across years; (d) migration timing parameters for each year, with orange segments indicating arrival and departure from the summering (northern) grounds, and the blue segments indicating timing of arrival and departure at the wintering grounds; (e) migration arrival and departure location across years, with blue and orange indicating winter (southern) and summer (northern) locations.

197 availability. Notably, the path to this equilibrium is somewhat indirect, with the later winter range
 198 taking more time to stabilize than the earlier summer range. The eventual steady state is one where
 199 the foraging efficiency is relatively high.

200 We ran this process for 8100 parameter combinations (figure 3). The fundamental dynamic of
 201 attaining and maintaining a well-matched migration phenology, can occur under many combinations

of parameter values, but all parameters play a role. Among the more intuitive results are that greater values of α (resource following) lead to an improved ability to match the migration. Resource peaks with larger spatial extent (higher σ_x) are generally better for migration matching. A larger set of parameters matched migration with low diffusion than high diffusion.

Less intuitive was the high importance of the sociality parameters, in particular the spatial scale of the swarming. Higher levels of social attraction (β) led to improved migration matching except in those cases where the sociality scale λ was high. Thus, for example, at $\lambda = 20$, no simulations at $\beta \geq 200$ managed to acquire or maintain a matched migration. However, at $\lambda = 50$ or 100 , the migration was slightly better matched at high values of β (figure 3). The spatial extent of the swarm was a remarkably significant variable. Smaller swarms were able to match migration only at low values of social attraction ($\beta = 200$), and relatively high values of resource attraction ($\alpha \geq 600$).

Random forest analyses, whether on the log of total mismatch or on the classification of a perfect match, uniformly show that the most important variables (Breiman, 2001) were α and λ (4.14 and 4.02 proportional increase in MSE), and the least important was σ_t , with a 0.5 proportional increase (table X).

Overall, foraging efficiency was strongly correlated with migration matching, as expected. At high mismatch (> 50), foraging efficiency was low (mean 0.29, s.d. 0.16) compared to the near-perfect matching migrations (mean 0.58, s.d. 0.14). However, somewhat higher mismatch (1 to 5) showed an even higher overall foraging efficiency (mean 0.62, s.d. 0.18 - see also figure 4).

3.2 Learning to migrate

Figure 5 illustrates the ability of the model animals to learn to migrate in a weakly drifting resource environment with a narrow pulse of resource peaking at 30 and -30 (at days 25 and 75), but a uniform distribution of resource at times 0 and 50. In order to learn to migrate, the system needed to have a higher exploratory impulse (higher diffusion constant ϵ), a stronger resource advection (higher α) and somewhat weaker sociality (lower β). The qualitative behavior of this process was to start drifting towards the summer resource, while slowly developing a weak pulse towards the winter resource as well. After first locking in on the summer resource, the winter migration, driven both by high diffusion and high resource following, slowly extended itself until both narrow peaks of resource can be consistently reached.

The model had, in general, a difficult time learning migration from a non-migratory initial condition. Thus, out of 4047 successful runs, only 4 attained mismatch below 1, and 130 below 5. Conditions that were more conducive to learning migration were pulses of *longer* duration (high σ_t), but *smaller* in scope (low σ_x), suggesting that the feedback that encourages migration needs to be compact in space but long enough in duration to lock in to the memory.

3.3 Adapting to climate change

To assess the ability of the system to adapt to a trending climate, we generated scenarios with slow (0.25 units / year), medium (0.5 units / year), and fast (1 unit / year) drift outward of the two resource pulses. We then assessed 40 parameter combinations for each of those scenarios, a high and low value of resource following ($\alpha = 400$ and 100) crossed with a high and low value of sociality ($\beta = 400$ and 100) crossed with 10 values of the spatial scale of sociality ($\lambda = 20$ to 200). The spatial and temporal scale of the resource pulses were fixed to a relatively “easy” to adapt to $\sigma_x = 12$ and $\sigma_t = 6$. We computed the adaptation index and foraging efficiency for each of the 120




Figure 3. Migration phenology matching across six model parameters. Low and high diffusion ($\epsilon = 1$ and 8 in upper and lower panel blocks), tight, medium and loose swarms ($\lambda = 20, 50, 100$) left to right panels. Within each of these blocks, high values of the resource following parameters α from 0 to 1000 are left to right, and higher values of the sociality parameter β are bottom to top. Within each of the combinations of $\epsilon, \lambda, \alpha, \beta$, we show results ranging across 6 values of resource duration (σ_t , 3 to 15 left to right), and 6 values of resource extent (σ_x , 3 to 15 bottom to top), as in the zoomed-in panel at (bottom right). The color scheme reflects the total mismatch, i.e. the sum of the absolute differences between the migration timing and locations from the resource peak. White squares represent runs that numerically failed to estimate migration parameters.

runs (figure 6). We were interested in the dynamics against λ due to the uniformly high importance of this parameter for determining the success of this process to match migration in steady states.

As figure 6 shows, higher values of resource following ($\alpha = 400$; orange circles) are nearly universally better for keeping up with climate change (SA values near 1). Furthermore, when combined with high sociality ($\beta = 400$; right panels), nearly all parameter combinations do a good job keeping up with climate change (SA values ranging between 0.53 and 0.85 for a swarm size greater than 50). However, that maximum value is still less than 1 , suggesting that truly matching a steadily drifting trend is very difficult. Small swarms ($\lambda < 50$) have a very hard time adapting when the social attraction is high, but do fairly well when social attraction is lower. But larger sized swarms do progressively worse across more parameterizations, e.g. in the most rapid climate change scenario, the SA drops from 0.83 to -0.13 as the swarm increases in size from 40 to 200 (essentially, the entire spatial domain).

A rather more dramatic pattern is visible for the lower foraging attraction scenario ($\alpha = 100$; blue circles). Notably, no parameter combination at this value comes close to keeping up with the rapid climate change (SA range -0.64 to 0.13). For slower climate change, however, there is a window of values for the swarm size between 40 and 80 , where the SA exceeds 1 , and then crashing quite rapidly to negative values of SA as that swarm size increases. These “super-adaptive” processes




Figure 4. Box-plots of foraging efficiency against mismatch across several values of foraging patch duration.

indicate a unique sweet spot where a swarm is large enough to capture and adapt to the drifting resource, but not so large that the information gathered in a given year is too weak to adjust the migratory behavior in a following year.

As anticipated, better adaptation to the drifting resource correlated strongly with higher foraging efficiency (inset boxplots).

3.4 Reference memory and stochasticity

While recent memory can be helpful for adapting to a single novelty or a smoothly changing conditions, we hypothesized that more conservative approach, relying on a “reference memory” may be more beneficial when conditions change stochastically. We tested this hypothesis by solving a set of models across a range of κ values from 0 (all recent memory) to 1 (all reference memory). In these scenarios, we ran the system for as many years as needed with no stochasticity to acquire a stationary state (i.e. similarity index greater than $1-1e-6$). We then used the stationary state as the reference memory, and then ran the process for an additional 50 years with a stochasticity (i.e. standard deviation in peak location of the resource) ranging from 0 to 12.

Figure 7 illustrates that, as predicted, the highest level of κ can significantly help foraging efficiency, with some variation across the spatial scale of sociality. When that scale of sociality is high enough (e.g. $\lambda = 120$, blue colors) there is greater probability of overlap with a stochastic resource, and a conservative, stable migratory regime is much more beneficial in the long run. Overall, as anticipated, the greater the stochasticity, the lower the foraging efficiency; and the greater the stochasticity, the greater the benefit of a conservative reference memory driven strategy.

3.5 Stochasticity and trends

We added trends to the stochastic process described above, adding 30 years of spatial outward trend in the resource. Our goal was to assess the ability of the process to keep up with climate change as a function of the reference memory parameter (figure 8). Over-reliance on reference memory ($\kappa = 1$) by definition does not allow the system to keep up with climate change, leading to an adaptation index of 0. However, in many cases a balancing of recent and reference memory (κ value between 0.6 and 0.8) in many cases was slightly but significantly better than relying entirely on recent memory. The smaller spatial scale (in the selected parameter space) does a generally better




Figure 5. Example of model learning to migrate. The resource is a “weakly drifting” resource and the initial (year 0) condition is non-migratory. The simulation was run for 100 years, and a sampling of those years (labeled) are presented in panel a: all years from 0 to 10, followed by 20, 40 and 100. Otherwise, panels are as in figure 2. Additional parameter values were $\epsilon = 5$, $\alpha = 500$, $\beta = 50$ and $\lambda = 40$.

289 job than the larger spatial scale at lower stochasticity. At higher level of stochasticity, however, the
 290 larger spatial scale outperforms the smaller spatial scale, which completely loses track of the climate
 291 change.

4 DISCUSSION

292 Animals navigate complex, dynamic, patchy environments. When there is a strongly localized
 293 and seasonal component to the resource dynamics, movement strategies limited to straightforward
 294 resource-following taxis necessarily fail to maximally exploit the dynamics. It is in these cases - quite
 295 common in the natural world – that seasonal migration becomes a viable, even necessary, strategy.




Figure 6. Adaptation to a steadily drifting resource. In three scenarios, the spatial coordinates of the resource drift by 0.25, 0.5, and 1 unit per year (top to bottom, respectively). The y-axis is the spatial adaptation index (SA), i.e. the trend of the memory-driven migration divided by the resource drift trend. Values near 1 indicate a behavior that keeps up with climate change, values near 0 indicate no change in migration behavior, and negative values indicate a trend that is opposite to the climate trend. We compare across spatial scales of sociality (λ - x-axis), for low and high values resource following ($\alpha = 400$ and 100 - orange and blue dots) and low and high values of sociality ($\beta = 100$ and 400 , left and right panels). The size of the circles is proportional to the foraging efficiency of the resulting parameter combinations. The bottom-right boxplots indicate the final year foraging efficiency against SA; purple and blue boxes indicate the highest values, orange and gray lower values.

296 However, when resources start shifting in space and time – as is occurring at an accelerated pace
 297 with recent global climate change – the migration phenology itself must exhibit some plasticity. It is
 298 our conjecture that this plasticity is greatly facilitated by a memory-driven process, in which recent
 299 experiences inform strategic behaviors in subsequent years. Here, we proposed a relatively simple,
 300 memory-driven mechanism by which a population can respond to long-term changes in resource
 301 dynamics, and a framework within which interacting memory, movement, social and resource
 302 dynamics can be explored.




Figure 7. Foraging efficiency (FE) across various values of reference memory κ (x-axis) for increasing amounts of interannual stochasticity (ψ_x , panels left to right), and two values of sociality spatial scale $\lambda = 80$ and 120. For the processes with non-zero stochasticity ($\psi_x > 0$), the process was run 90 times for values of κ . Points represent the average of the FE's across all 50 years and 90 replicates, and shaded areas represent a local regression (loess) smooth of FE against κ . In these scenarios, the resource following parameter $\alpha = 100$, the social attraction $\beta = 400$ and diffusion $\epsilon = 4$.

We found that the straightforward inclusion of a migration-specific memory in a social diffusion-advection framework was sufficient to capture many fundamental phenomena related to migration. In essence, our model allowed a population with a hybrid migratory and resource-following behavior to adjust the migratory behavior based on recent experiences with the resource location. The model, while simple, was able to emulate the successful navigation of an environment with temporally and spatially isolated seasonal resource patches, the spontaneous emergence of a migratory behavior, and intrinsic robustness to changes in those environmental resources, whether those changes were steadily shifting trends or inter-annual stochasticity. We postulated a process defined by relatively few parameters, and explored the ability of the process to navigate a fixed set of environments. Our goal was to understand the combinations of factors that lead to a resilient migration behavior. The foraging efficiency was a convenient metric of the utility of the migratory behavior, but was not a measure explicitly maximized by the model. Other metrics, such as foraging efficiency in a given season, may respond differently across model parameters and could be useful in understanding the relative success of alternative migratory strategies in different contexts (e.g., scenarios where there are differing constraints between 'low' and 'high' seasons or where good resource matching is particularly vital at certain stages in an animal's life history like gestation). However, we opted for an overall annually averaged metric in these initial explorations because that provides the broadest linkage between resource dynamics and the animals' locations and was consistent with the smooth and continuous nature of the resource distribution.

Our analyses afford several rather general conclusions. One is that migration in general can be acquired, as has been demonstrated for translocated ungulates (Jesmer et al., 2018), even in environments where resources are difficult to "surf" (figure 5). However, this requires a very strong attraction to resources, somewhat higher levels of exploratory behavior, and many years, findings that echo empirical observations (Jesmer et al., 2018). On the other hand, migrations can be




Figure 8. Role of reference memory in adapting to climate change for increasingly stochastic resource dynamics. We ran the model with a moderate rate of climate change (mean shift: 0.5/year) at five increasing levels of stochasticity (inter-annual standard deviation of resource peak 0, 3, 6 and 12, left to right panels). For non-zero stochasticity, we ran the process 30 times and present the mean and standard error of the spatial adaptation index across various values of the reference memory parameter κ : where $\kappa = 0$, the system modifies its migration based entirely on recent experiences; at $\kappa = 1$, the memory never changes from the reference memory. Other parameter values are resource following $\alpha = 100$, social attraction $\beta = 400$, and diffusion $\epsilon = 4$.

327 considered somewhat fragile, in the sense that under certain conditions, e.g. increasing stochasticity,
 328 rapidly shifting resources, or a shift in some of the system parameters, migration can collapse
 329 down into a localized non-migratory, residential behavior, at times with only a marginal cost to the
 330 foraging efficiency index. This sensitivity may explain why partially migratory populations composed
 331 of a mixture of resident and migratory individuals are so common and, apparently, evolutionary
 332 stable (Berthold, 1999; Chapman et al., 2011).

333 We also found that the ability to maintain, learn and adapt a migration depends strongly on
 334 properties of the resource dynamics. In particular, the reinforcement of memory and foraging is
 335 strongest when patches are concentrated in time, but relatively large in space. Interestingly, in
 336 most stable patterns, the eventual targeted migration arrival time coincided with the *peak*, rather
 337 than the beginning, of the resource dynamic. This indicates that the long-distance social migration
 338 behavior may be particularly reinforced when the targeted resource is very sudden (explosive). This
 339 is the case for the rapid green-up that occurs in high latitudes as snow recedes in tandem with
 340 extended day lengths leading to a rapid and intense local green-up, or, in a more extreme case, when
 341 resources are linked to the short-duration early blooming phenology of very particular plants (refs).

342 The sociality parameters – in particular, the spatial scale λ – were, somewhat unexpectedly, the
 343 most important for determining the resiliency of the process for stabilizing a migratory behavior. In
 344 particular, a population very small spatial scale tended to have a much more difficult time locking in
 345 to an adaptive migratory pattern, and only in conditions where the social attraction was relatively
 346 weak. In order to adapt to any shift in resources, *some* individuals must experience that resource,
 347 and the two sociality-related mechanisms to have that scope are either a relatively large spatial scale
 348 or a relatively weak social cohesion. On the other hand, overly large spatial scales compromised

the ability of the process to track climate change, due to a dilution of the population's ability to concentrate over available resource patches. *Multiple avenues by which 'sociality' (defined as swarm x attraction interplay) can structure migration depending on the relative balance of diffusion and resource following (figure ??). For example this could tie into the relative balance between individuals exploring the landscape to acquire new information about resources and their opportunities to interact with other individuals and communicate (or attract).*

All of the parameters in our model have well-defined biological interpretations. The diffusion (ϵ) captures short time-scaled randomness of movement, capturing the exploratory and short-term dispersive behavior. The foraging advection strength (α) captures the attraction of the population to better quality resources at a relatively large scale. These two parameters, the basic ingredients in diffusion-advection models of animal movement, have direct parallels to empirically estimated properties of animal behavior: diffusion is closely related to families of random walk models (Gurarie and Ovaskainen, 2011) while the advective taxis is related to the step- and resource selection functions that are routinely estimated from movement data (Potts and Schlägel, 2020). The spatial scale of the social group (λ) captures the spatial extent of the population, i.e. a population-level range. The sociality parameter (β) quantifies the strength an individual's desire to approach the center of the social group. This is not a parameter that is typically measured, though it would - in principle - be possible to estimate in a manner analogous to step-selection function using only other conspecifics as opposed to environmental variables as a covariate. Because both resources and populations were normalized to 1, the ratio between α and β can be interpreted as the relative importance of foraging to social cohesion. It does appear a certain balance between these two parameters is needed to successfully maintain, acquire, or adapt a migratory phenology.

Finally, the migration timing and location parameter can be similarly straightforwardly estimated from movement data (Cagnacci et al., 2015; Gurarie et al., 2019). Thus, for example, Gurarie et al. (2019) explicitly estimated the ranging area, timing, and seasonal range locations for migratory caribou (*Rangifer tarandus*), identifying the kind of inter-annual variation that is reflected in the stochastic scenarios explored here as well as some trends in the timing of migration. Finally, we note that our underlying assumption of the migration "urge" is consistent with the strong endogenous programs to migrate that are well-documented. Birds in particular exhibit a seasonal restlessness known as *Zugunruhe* (Berthold, 1999; Helm, 2006).

Individual based models have shown that collective knowledge is important, if not essential, to the evolution and process of migration (Shaw and Couzin, 2013; Guttal and Couzin, 2010; Berdahl et al., 2018). In our model, we assume that the migration process itself is driven by a collective trigger for the timing and locations of seasonal ranges and migration behavior. Synchrony of migration timing and high site fidelity are well documented for migratory species (Gurarie et al., 2019; Joly et al., 2021). We note, however, that diffusion-advection models can also be interpreted as a probabilistic description of a single individual's movement. In this case, λ would correspond to an individual home-range and β would be an individual's tendency to be drawn to the center of that home range, akin to an individual migratory Ornstein-Uhlenbeck process (Gurarie et al., 2017).

The reference memory parameter κ is, of course, impossible to observe directly. Our model does, however, allow us to explore in an heuristic way the conditions under which a strong cultural tendency to migrate with certain fixed patterns can help a population hedge against stochasticity (Abrahms et al., 2019; Fagan, 2019).

392 Importantly, our model did not include any selection, inheritance or birth or death processes,
393 i.e. there was no evolutionary maximization of a fitness, as in many models that explicitly explore the
394 evolution of migration (e.g. [Guttal and Couzin, 2010](#); [Shaw and Couzin, 2013](#); [Anderson et al., 2013](#)).
395 Nonetheless, it is instructive to compare the fundamental assumptions behind such evolutionary
396 individual based models and a social memory based model like ours. For example, [Anderson et al.](#)
397 (2013) explored the resilience of a population under selective pressure under persistent trends and
398 increased stochasticity of a drifting optimal resource window, showing that a certain amount of
399 heritable phenotypic plasticity is necessary to adapt successfully to climate change even at the
400 cost of efficiency. Our model underscores the fact that some level of resilience and adaptability
401 can be attained with a purely cognitive process that balances sociality with long and short term
402 collective memory. Importantly, this knowledge must be transmitted through social and cultural,
403 rather than genetic, pathways. The high level of sociality among migratory animals, as well as
404 multi-annual parent offspring bonds, are an evident pathway for that kind of transmission. As with
405 those evolutionary models, however, it is clear that when changes are too rapid, no amount of
406 cognition can help entirely mitigate against adverse outcomes. Furthermore, if behaviors are not
407 sufficiently plastic (i.e., if κ is too close to 1), then adaptation is that much more difficult.

408 Rapid environmental change, both global warming and increased anthropogenic development, is
409 causing severe and dramatic impacts to the widespread and generally successful strategy of seasonal
410 migration for many taxa, and the fate of many animal migrations is a topic of increasing concern
411 ([Kauffman et al., 2021](#); [Wilcove and Wikelski, 2008](#)). The ability of animals to respond to these
412 changes depends deeply on their behavioral plasticity and cognitive abilities. The importance of those
413 abilities is in direct proportion to the difficulty in studying them directly. By quantitatively exploring
414 the properties of a heuristic model that distill many of the main properties of wild populations in
415 dynamic and seasonal environments, we hope to have identified some broad patterns that might
416 guide further empirical exploration of the cognitive underpinnings of adaptability and resilience.

5 NOMENCLATURE

CONFLICT OF INTEREST STATEMENT

417 The authors declare that the research was conducted in the absence of any commercial or financial
418 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

419 EG and WFF provided the original idea. EG and SP developed and ran models and analysis. All
420 authors contributed to the writing.

FUNDING

421 EG, WFF, GCC and RSC, were supported in part by NSF award DMS 1853478. EG and WFF
422 were further partially supported by NSF grant IIBR 1915347.

ACKNOWLEDGMENTS

423 The authors are grateful to Quentin Read at SESYNC for computational support and advice for
424 multi-node multi-core model runs.

DATA AVAILABILITY STATEMENT

The only data analyzed in this manuscript was generated by simulation scripts made available in the GitHub repository at <https://github.com/EliGurarie/memorymigration>.

REFERENCES

- Abrahms, B., Hazen, E. L., Aikens, E. O., Savoca, M. S., Goldbogen, J. A., Bograd, S. J., et al. (2019). Memory and resource tracking drive blue whale migrations. *Proceedings of the National Academy of Sciences* 116, 5582–5587.
- Anderson, J. J., Gurarie, E., Bracis, C., Burke, B. J., and Laidre, K. L. (2013). Modeling climate change impacts on phenology and population dynamics of migratory marine species. *Ecological Modelling* 264, 83–97.
- Avgar, T., Street, G., and Fryxell, J. M. (2014). On the adaptive benefits of mammal migration. *Canadian Journal of Zoology* 92, 481–490. doi:10.1139/cjz-2013-0076
- Berdahl, A. M., Kao, A. B., Flack, A., Westley, P. A., Codling, E. A., Couzin, I. D., et al. (2018). Collective animal navigation and migratory culture: from theoretical models to empirical evidence. *Philosophical Transactions of the Royal Society B: Biological Sciences* 373, 20170009. doi:10.1098/rstb.2017.0009
- Berthold, P. (1999). A comprehensive theory for the evolution, control and adaptability of avian migration. *Ostrich* 70, 1–11. doi:10.1080/00306525.1999.9639744
- Bhattacharyya, A. (1943). On a measure of divergence between two statistical populations defined by their probability distributions. *Bull. Calcutta Math. Soc.* 35, 99–109
- Bischof, R., Loe, L. E., Meisingset, E. L., Zimmermann, B., Van Moorter, B., and Mysterud, A. (2012). A migratory northern ungulate in the pursuit of spring: jumping or surfing the green wave? *The American Naturalist* 180, 407–424
- Bracis, C. and Mueller, T. (2017). Memory, not just perception, plays an important role in terrestrial mammalian migration. *Proceedings of the Royal Society B: Biological Sciences* 284, 20170449
- Breiman, L. (2001). Random forests. *Machine learning* 45, 5–32
- Cagnacci, F., Focardi, S., Ghisla, A., van Moorter, B., Merrill, E. H., Gurarie, E., et al. (2015). How many routes lead to migration? comparison of methods to assess and characterize migratory movements. *Journal of Animal Ecology* 85, 54–68. doi:10.1111/1365-2656.12449
- Chapman, B. B., Brönmark, C., Nilsson, J.-Å., and Hansson, L.-A. (2011). The ecology and evolution of partial migration. *Oikos* 120, 1764–1775. doi:10.1111/j.1600-0706.2011.20131.x
- Dingle, H. (2014). *Migration: the biology of life on the move* (Oxford University Press, USA)
- Elzhov, T. V., Mullen, K. M., Spiess, A.-N., and Bolker, B. (2016). *minpack.lm: R Interface to the Levenberg-Marquardt Nonlinear Least-Squares Algorithm Found in MINPACK, Plus Support for Bounds*. R package version 1.2-1
- Fagan, W., Gurarie, E., Bewick, S., Howard, A., Cantrell, R., and Cosner, C. (2017). Perceptual ranges, information gathering, and foraging success in dynamic landscapes. *The American Naturalist* 189, 474–489. doi:10.1086/691099
- Fagan, W., Hoffman, T., Dahiya, D., Gurarie, E., Cantrell, R., and Cosner, C. (2019). Improved foraging by switching between diffusion and advection: benefits from movement that depends on spatial context. *Theoretical Ecology* 13, 127–136. doi:10.1007/s12080-019-00434-w
- Fagan, W. F. (2019). Migrating whales depend on memory to exploit reliable resources. *Proceedings of the National Academy of Sciences* 116, 5217–5219. doi:10.1073/pnas.1901803116

- Fryxell, J. M., Greever, J., and Sinclair, A. (1988). Why are migratory ungulates so abundant? *The American Naturalist* 131, 781–798
- Grünbaum, D. (1994). Translating stochastic density-dependent individual behavior with sensory constraints to an eulerian model of animal swarming. *Journal of Mathematical Biology* 33, 139–161. doi:10.1007/BF00160177
- Gurarie, E., Anderson, J. J., and Zabel, R. W. (2009). Continuous models of population-level heterogeneity inform analysis of animal dispersal and migration. *Ecology* 90, 2233–2242
- Gurarie, E., Cagnacci, F., Peters, W., Fleming, C. H., Calabrese, J. M., Mueller, T., et al. (2017). A framework for modelling range shifts and migrations: asking when, whither, whether and will it return. *Journal of Animal Ecology* 86, 943–959
- Gurarie, E., Hebblewhite, M., Joly, K., Kelly, A. P., Adamczewski, J., Davidson, S. C., et al. (2019). Tactical departures and strategic arrivals: Divergent effects of climate and weather on caribou spring migrations. *Ecosphere* 10, e02971
- Gurarie, E. and Ovaskainen, O. (2011). Characteristic spatial and temporal scales unify models of animal movement. *The American Naturalist* 178, 113–123
- Guttal, V. and Couzin, I. D. (2010). Social interactions, information use, and the evolution of collective migration. *Proceedings of the National Academy of Sciences* 107, 16172–16177. doi:10.1073/pnas.1006874107
- Helm, B. (2006). Zugunruhe of migratory and non-migratory birds in a circannual context. *Journal of Avian Biology* 37, 533–540. doi:10.1111/j.2006.0908-8857.03947.x
- Jesmer, B. R., Merkle, J. A., Goheen, J. R., Aikens, E. O., Beck, J. L., Courtemanch, A. B., et al. (2018). Is ungulate migration culturally transmitted? Evidence of social learning from translocated animals. *Science* 361, 1023–1025
- Joly, K., Gurarie, E., Hansen, D. A., and Cameron, M. D. (2021). Seasonal patterns of spatial fidelity and temporal consistency in the distribution and movements of a migratory ungulate. *Ecology and Evolution* 11, 8183–8200. doi:10.1002/ece3.7650
- Kauffman, M. J., Cagnacci, F., Chamaillé-Jammes, S., Hebblewhite, M., Hopcraft, J. G. C., Merkle, J. A., et al. (2021). Mapping out a future for ungulate migrations. *Science* 372, 566–569. doi:10.1126/science.abf0998
- Kölzsch, A., Bauer, S., De Boer, R., Griffin, L., Cabot, D., Exo, K.-M., et al. (2015). Forecasting spring from afar? Timing of migration and predictability of phenology along different migration routes of an avian herbivore. *Journal of Animal Ecology* 84, 272–283
- Kot, M., Lewis, M. A., and Driessche, P. v. d. (1996). Dispersal data and the spread of invading organisms. *Ecology* 77, 2027–2042
- Lin, H.-Y., Fagan, W. F., and Jabin, P.-E. (2021). Memory-driven movement model for periodic migrations. *Journal of Theoretical Biology* 508, 110486. doi:10.1016/j.jtbi.2020.110486
- Mogilner, A. and Edelstein-Keshet, L. (1999). A non-local model for a swarm. *Journal of Mathematical Biology* 38, 534–570. doi:10.1007/s002850050158
- Okubo, A. (1986). Dynamical aspects of animal grouping: Swarms, schools, flocks, and herds. *Advances in Biophysics* 22, 1–94. doi:10.1016/0065-227X(86)90003-1
- Okubo, A. and Levin, S. (2001). *Diffusion and Ecological Problems: Modern Perspectives* (New York: Springer Verlag)
- Potts, J. R. and Schlägel, U. E. (2020). Parametrizing diffusion-taxis equations from animal movement trajectories using step selection analysis. *Methods in Ecology and Evolution* 11, 1092–1105. doi:10.1111/2041-210X.13406

- 511 Robinson, R., Crick, H., Learmonth, J., Maclean, I., Thomas, C., Bairlein, F., et al. (2009).
512 Travelling through a warming world: climate change and migratory species. *Endangered Species*
513 *Research* 7, 87–99. doi:10.3354/esr00095
- 514 Shaw, A. K. (2016). Drivers of animal migration and implications in changing environments.
515 *Evolutionary Ecology* 30, 991–1007. doi:10.1007/s10682-016-9860-5
- 516 Shaw, A. K. and Couzin, I. D. (2013). Migration or residency? the evolution of movement behavior
517 and information usage in seasonal environments. *The American Naturalist* 181, 114–124
- 518 Skalski, G. T. and Gilliam, J. F. (2003). A diffusion-based theory of organism dispersal in
519 heterogeneous populations. *The American Naturalist* 161, 441–458
- 520 Skellam, J. G. (1951). Random dispersal in theoretical populations. *Biometrika* 38, 196–218
- 521 Soetaert, K. and Meysman, F. (2012). Reactive transport in aquatic ecosystems: Rapid model
522 prototyping in the open source software r. *Environmental Modelling and Software* 32, 49–60
- 523 Soetaert, K., Petzoldt, T., and Setzer, R. W. (2010). Solving differential equations in R: Package
524 deSolve. *Journal of Statistical Software* 33, 1–25. doi:10.18637/jss.v033.i09
- 525 Turchin, P. (1998). *Quantitative analysis of movement: measuring and modeling population*
526 *redistribution in animals and plants* (Sinauer Associates)
- 527 Wilcove, D. S. and Wikelski, M. (2008). Going, going, gone: Is animal migration disappearing.
528 *PLoS Biology* 6, e188. doi:10.1371/journal.pbio.0060188