Contents

A Introduction **1**

B Minkowski space and space-like surfaces **1**

Space-like immersions. 2
Entire surfaces. 2
The Gauss map in $\mathbb{M}^{2,1}$ 4
$\mathbb{H}^2$ is a model for the Hyperbolic plane. 5
Curvatures of the surface S 5

A Introduction

Goal of the work, maybe a simple example, some mention to preliminary information and a soft mention to what's the problem and the philosophy.

B Minkowski space and space-like surfaces

The environment we will work within is the Minkowki 3-space, defined as $\mathbb{R}^3$ endowed with the pseudometric

$$ds^2 = (dx^1)^2 + (dx^2)^2 - (dx^3)^2.$$

In order to distinguish the use of this metric from the standard euclidean metric of $\mathbb{R}^3$ we will use the notation $\mathbb{M}^{2,1}$. We will also denote the coordinates either as (x^1, x^2, x^3) or (x, y, t), choosing the notation according to convenience. We will also identify the slice $\{(x, y, 0)\}$ of $\mathbb{M}^{2,1}$ with $\mathbb{R}^2$ and with $\mathbb{C}$.

The space $\mathbb{M}^{2,1}$ is an example of a more general object called a pseudo-Euclidean vector space, a term that refers to a finite-dimensional vector space V endowed with a nondegenerate (possibly not positive-definite) symmetric bilinear form g. This kind of space serves as the "infinitesimal model" for pseudo-Riemannian manifolds, smooth manifolds endowed with a smooth nondegenerate $(0, 2)$-tensor. If the signature of this tensor is exactly $(n-1, 1)$, the manifold is called a Lorentzian manifold.
The following definition, borrowed from physics, will be useful later.

Definition. Given a Lorentzian manifold (M, g), we say that a vector $v \in T_pM$ is

- space-like if $\langle v, v\rangle > 0$
- light-like if $\langle v, v\rangle = 0$
- time-like if $\langle v, v\rangle < 0$

We will always denote with S a differentiable manifold of dimension 2 (a surface) immersed in $\mathbb{M}^{2,1}$, $\iota : S \hookrightarrow \mathbb{M}^{2,1}$. As usual, the immersion induces a scalar product on S, the first fundamental form or pull-back metric, which we indicate as $\mathbf{I} = \iota^*(g_{\mathbb{M}^{2,1}})$ and write in coordinates as $\mathbf{I} = g_{ij}dx^i dx^j$ (for simplicity we always forget the immersion ι).

Space-like immersions.
The object we will focus on are not arbitrary immersed surfaces but the so-called space-like surfaces of $\mathbb{M}^{2,1}$. This term refers to the immersed surfaces in $\mathbb{M}^{2,1}$ whose induced metric is positive-definite hence Riemannian, or in other terms, they're surfaces S whose tangent bundle $\{(p, v)\}$ is made only of space-like vectors.

A nice property of this kind of surface is that they are locally the graph of a function of the form $x^3 = f(x^1, x^2)$.
We can show this with a simple argument:
First, we assumed that the pseudometric restricted to S,$\langle \cdot, \cdot \rangle |_S \geq 0$, is definite positive, which means that the tangent vectors of S at any point p belong to the set of space-like vectors:

$$T_p \subset \mathscr{C} := \left\{ v \in \mathbb{M}^{2,1} : \langle v, v \rangle \geq 0 \right\} = \{t^2 \leq x^2 + y^2\}.$$

This is just the outside-region of a double cone in $\mathbb{M}^{2,1}$, whose symmetry axis is $\left\{x^1 = 0, x^2 = 0\right\}$. A plane is entirely made of space-like vectors iff it's contained between the two branches of the cone.
Then, having assumed that S is an immersed surface, hence an immersed submanifold, it is locally the solution of an equation of the form $F(x, y, z) = 0$. It is well known that from this local expression we can express the normal vector to the surface at a point p: $N_p = \vec{\nabla} F(p)$.
Furthermore, we observe that N_p is orthogonal to the space-like plane T_pS and so it cannot be horizontal. The last two statements taken together give us $\partial_z F(p) = (N_p)^3 \neq 0$, and so we can apply the implicit function theorem and express the surface as the graph of $z = f(x, y)$ near the point p.
Where the local representation is valid, we can parametrize the surface as $(x, y, f(x, y))$, and derive the local frame $V_x = (1, 0, \partial_x f(x, y))$, $V_y = (1, 0, \partial_y f(x, y))$, for which the first fundamental form is represented by the matrix

$$g = \begin{pmatrix} \langle V_x, V_x \rangle & \langle V_y, V_x \rangle \\ \langle V_x, V_y \rangle & \langle V_y, V_y \rangle \end{pmatrix} = \begin{pmatrix} 1 - f_x^2 & -f_x f_y \\ -f_x f_y & 1 - f_y^2 \end{pmatrix}.$$

Since the parametrization is obviously regular, the condition on g being a metric translates in having a strictly positive determinant,

$$0 < \det g = 1 - (f_x^2 + f_y^2) = 1 - |\vec{\nabla} f|^2,$$

which means $|\vec{\nabla} f|^2 < 1$.

Entire surfaces.
Given a surface S immersed in $\mathbb{M}^{2,1}$, we can define a projection $\Pi : S \to \mathbb{R}^2$ by forgetting the time coordinate, $\Pi(x^1, x^2, x^3) := (x^1, x^2)$ for any $(x^1, x^2, x^3) \in S$. We call the surface S is entire when the map Π is surjective. If the surface is globally a graph of a function φ, this condition is exactly stating that the function φ is an entire map from $\mathbb{R}^2$ to $\mathbb{R}$.

The situation for space-like surfaces is actually a lot nicer than what we presented up to now, as the representation as a graph is not only local but actually global as we prove in the proposition below.

Proposition 1. *Let Ω be a domain in $\mathbb{C}$ (that is, an open connected set) and let $S : \Omega \to \mathbb{M}^{2,1}$ be a space-like immersion. Suppose that S is complete with respect to the induced Riemannian metric. Then S is an entire graph and Ω is simply connected.*

Before laying out the proof let's recall a couple of results from the theory of coverings.

Proposition 2. *Let M and N be two differentiable manifolds and $f : M \to N$ a smooth map*

- *If M is simply connected, N is connected and f is a covering map, then it is a homeomorphism.*
- *We say that the map f satisfies the path lifting property if for every $p \in N$ and for every path $\alpha : I \to M$ such that $\alpha(0) = f(p)$ there exists a path $\tilde{\alpha} : I \to N$ such that $\tilde{\alpha}(0) = p$ and $f \circ \tilde{\alpha} = \alpha$. We can assume without loss of generality that the paths involved are differentiable.*
- *If f is a local homeomorphism and satisfies the path lifting property, then it's a covering map.*

Proof. The idea of the proof is to show that $\Pi \circ S$ is a global diffeomrphism. Clearly this implies that Π restricted to S is a diffeomeorphism on $\mathbb{R}^2$ and so that S is an entire graph. From that we obtain also that Ω is simply connected since it's diffeomorphic to $\mathbb{R}^2$.

Let's observe that the target of $\Pi \circ S$ is just $\mathbb{R}^2$, which is simply connected, and the domain is Ω, which is connected since it's a domain. We already proved, by showing that S is locally a graph, that the map $\Pi \circ S$ is a local homeomorphism.
Having met these conditions, if we show that $\Pi \circ S$ satisfies the path lifting property, we conclude that it is a diffeomorphism.

Let's fix a point $z \in \Omega$ and a path $\gamma : [0,1] \to \mathbb{R}^2$ of bounded length, $\ell(\gamma) < \infty$, such that $\gamma(0) = (\Pi \circ S)(z)$. Our goal is to show that γ admits a lift to a path γ^* on Ω starting at z.
To this end, let's consider the set of times up to which such a lift exists,

$$\mathcal{J} := \{\tau \in [0,1] \mid \exists \gamma^* : [0,t] \to \Omega \text{ such that } (\Pi \circ S) \circ \gamma^* = \gamma \text{ and } \gamma^*(0) = z\},$$

and let's define $\bar{\tau} := \sup \mathcal{J}$.

If we prove that $\bar{\tau} = 1$, then γ admits a lift on the whole interval $[0,1]$ and we have finished. Then let's suppose by contradiction that $\bar{\tau} < 1$ and define $\gamma^* : [0, \bar{\tau}) \to \Omega$ the lift of γ up to $\bar{\tau}$. This curve exists by definition of $\bar{\tau}$.

Let's observe that $0 \in \mathcal{J}$ and that $\mathcal{J}$ is open. This is true because, being $\Pi \circ S$ a local homeomorphism, if a certain τ is in $\mathcal{J} \cap (0,1)$ then the whole interval $(\tau - \varepsilon, \tau + \varepsilon)$ for $\varepsilon > 0$ sufficiently small, belongs to it.

Finally, we need to prove that the limit $\lim_{\tau \to \bar{\tau}} \gamma^*(t)$ exists. If this is true we can then extend γ^* up to $\bar{\tau}$, which means that $\bar{\tau} \in \mathcal{J}$. Since $\mathcal{J}$ is open, there exists an element $\bar{\tau} + \varepsilon \in \mathcal{J}$, $\varepsilon > 0$. This contradicts the definition of $\bar{\tau}$ and

proves $\bar{\tau} = 1$ since the alternative is absurd.

To prove that $\lim_{\tau \to \bar{\tau}} \gamma^*(\tau)$ exists we just need a couple of observations: First we note that the map Π dilates distances, as

$$\begin{aligned} \langle v, v \rangle_{\mathbb{M}^{2,1}} &= (v^1)^2 + (v^2)^2 - (v^3)^2 \\ &\leq (v^1)^2 + (v^2)^2 \\ &= \langle \Pi v, \Pi v \rangle_{\mathbb{R}^2} . \end{aligned}$$

From this it's clear that $\ell(\gamma^*) \leq \ell(\gamma)$.
Now, if we fix any sequence $\tau_k \to \bar{\tau}$ and observe that as γ is differentiable, it's derivative is bounded, so $\ell(\gamma|_{[\tau_k, \tau_m]}) \leq C|\tau_k - \tau_m|$. By observing that

$$\begin{aligned} d(\gamma^*(\tau_k), \gamma^*(\tau_m)) &\leq \ell(\gamma^*|_{[\tau_k, \tau_m]}) \\ &\leq \ell(\gamma|_{[\tau_k, \tau_m]}) \\ &\leq C|\tau_k - \tau_m| \end{aligned}$$

we deduce that also $(\gamma^*(\tau_k))_k$ is a Cauchy sequence, hence converging since S is complete and Ω is to be intended with the induced metric.
By the arbitrariness of the sequence $(\tau_k)_k$ we deduce the limit $\lim_{\tau \to \bar{\tau}} \gamma^*(\tau)$ exists. $\square$

In this section we have essentially proved that a complete space-like surface, immersed in $\mathbb{M}^{2,1}$ is actually the graph of a map with $|\vec{\nabla} f|^2 < 1$. It's easy to convince ourselves that the inverse result also holds and so that this is actually a characterization.

The Gauss map in $\mathbb{M}^{2,1}$
The surface S is an immersed submanifold of codimension 1. As we have a nondegenerate form on the ambient space, g, once fixed a $p \in S$, we can write the following decomposition, which varies smoothly as we change the point:

$$\mathbb{M}^{2,1} = T_p\mathbb{M}^{2,1} = (T_pS)^\perp \oplus T_pS$$

The codimension of the submanifold M is equal to the dimension of $(T_pS)^\perp$, that is, is exaclty one. In fact this space is generated only by the normal vector to S, N_p.
We can easily show that N must be a time-like vector, due to the fact that T_pS is made of space-like vectors.

Let's fix $e_1, e_2 \in T_pM$ two independent vectors. Since N belongs to $(T_pM)^\perp$ we have $\langle e_1, N \rangle = \langle e_2, N \rangle = 0$. This implies that when we look at the representative matrix of g_p written in the basis (N, e_1, e_2), we have

$$[g_p] = \begin{pmatrix} \langle n, n \rangle & 0 & 0 \\ 0 & \langle e_1, e_1 \rangle & \langle e_1, e_2 \rangle \\ 0 & \langle e_2, e_1 \rangle & \langle e_2, e_2 \rangle . \end{pmatrix} =: \begin{pmatrix} \langle n, n \rangle & 0 \\ 0 & [(g|_S)_p] \end{pmatrix} .$$

Where the block $[(g|_S)_p]$ represents the form g restricted to the subspace T_pM. We conclude that $\langle n, n \rangle$ is negative just by observing that $\det[g_p] < 0$ as g is a

pseudometric, while $\det[(g|_S)_p] > 0$ since g becomes a metric when restricted to S.

We can now repeat the classical construction of the Gauss map in this pseusoriemannian context.
The idea is just to take a unit vector in $(T_pM)^\perp$. On this space the norm of the vectors is negative so we will need to normalize them to -1 instead of 1.
So let's fix $w \in (T_pM)^\perp$ and scale it by a factor k such that $\langle kw, kw\rangle = -1$. If we solve this equation we obtain two distinct solutions, that send w to one of the two branches of $\hat{\mathbb{H}}^2 = \{v \in \mathbb{M}^{2,1} | \langle v, v\rangle = -1\}$, which means that a choice is needed. Here we consistently choose the normalized vector to to be in the space

$$\mathbb{H}^2 = \{v \in \mathbb{M}^{2,1} | \langle v, v\rangle = -1, v^3 > 0\}.$$

This way we have constructed a map

$$u : S \to \mathbb{H}^2,$$

the Gauss map of the surface S. Of course the only difference from the classical theory of surfaces is the codomain of the map u, where it is the sphere $\mathbb{S}^2$. Like in the classical case, the choice of the normal vector specifies an orientation for the surface, which implies that the Gauss map is gloabally defined only when the surface is orientable. In our case, this condition is trivially met since we're dealing with a graph surface.

$\mathbb{H}^2$ is a model for the Hyperbolic plane.

Curvatures of the surface S