

Time Series Econometrics

Properties of Stationary and Integrated Time Series

Eliab Luvanda

2026-04-02

Abstract

This handout reviews properties of stationary and integrated time series, covering definitions, examples, and visual illustrations. It provides a concise reference for students and researchers in time-series econometrics.

Table of contents

1	Introduction	2
2	Stationary stochastic Processes	4
2.1	Weak Stationarity	4
2.2	Strong Stationarity	6
2.3	Example Stationary Processes	9
2.3.1	White Noise Process	9
2.3.2	Moving Average (MA) Processes	11
2.3.3	Autoregressive (AR) Processes	11
2.3.4	Autoregressive Moving Average (ARMA) Processes	11
3	Non-stationary Stationary Processes	11
3.1	Trend Stationary Processes	11
3.2	Difference Stationary or Integrated Processes (Unit Root Processes)	11
4	Spurious Regression	12
4.1	Theoretical Framework	12
4.1.1	Brownian Motion Representation	13
4.1.2	Limit Distribution of the OLS Estimator	13
4.1.3	Divergence of the t-Statistic	13
4.2	Monte Carlo Simulation	13
4.3	Empirical Findings	14
4.3.1	Increasing False Significance	14
4.3.2	Distribution of the OLS Estimator	14
4.3.3	Scaled Estimator $N\hat{\beta}$	15

Bibliography	16
---------------------	-----------

Appendix	17
-----------------	-----------

Random Walk Sample Paths	17
AR(1) Time Series Simulation	17
Stationary AR(1) Process Plot	18
Gaussian White Noise	19
OLS Simulation Summary Table	20
Spurious Regression Table	21
Unscaled OLS Estimator Density	22
Scaled Estimator Density	23
Julia Simulation: OLS AR(1) vs Random Walk	23
Julia Simulation: Spurious Regression	23

List of Figures

1 Multiple simple random-walk sample paths (5)	4
2 Ensemble of 4 stationary AR(1) sample paths; the ensemble mean at t_p is $\bar{X}_{t_p} = \frac{1}{4} \sum_{j=1}^4 X_{t_p}^{(j)}$	5
3 Simulated stationary AR(1) process (mean dashed)	6
4 Gaussian white noise (i.i.d. $N(0,1)$)	8
5 OLS $\hat{\phi}$ distributions for stationary AR(1) and random walk processes	12
6 Densities of $\hat{\beta}$ (unscaled) across sample sizes	14
7 Densities of $N\hat{\beta}$ across sample sizes	15

List of Tables

1 Summary comparison of weak (second-order) and strong (strict) stationarity: definitions, requirements, implications, and practical considerations.	9
2 Monte Carlo summary (n=2000, reps=1,000,000)	12
3 Spurious regression rejection frequencies by sample size	14

1 Introduction

Time-series econometrics analyzes sequences of observations indexed by time and explicitly models temporal dependence. Time-series data commonly exhibit autocorrelation, non-stationarity, and seasonality. As a result, the standard i.i.d. assumptions often fail and require specialized methods. This handout summarizes core definitions and properties of stationary and integrated processes, and gives concise examples and illustrations.

Distinguishing stationary from integrated series is essential for choosing models and making valid inferences. Central to these ideas is the notion of a stochastic process, defined below.

Definition 1.1. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let T be an index set (often representing time, e.g. $T = \mathbb{N}$ for discrete time or $T = \mathbb{R}_+$ for continuous time).

A stochastic process is a family of random variables

$$\{X(\tau, \omega) : \tau \in T, \omega \in \Omega\}$$

such that for each fixed $\tau \in T$, the mapping

$$X(\tau, \cdot) : \Omega \rightarrow \mathbb{R}$$

is a random variable on $(\Omega, \mathcal{F}, \mathbb{P})$.

Alternatively, we define a stochastic process as follows:

Definition 1.2. One can equivalently view a stochastic process as a function

$$X : \Omega \times T \rightarrow \mathbb{R}, \quad (\omega, \tau) \mapsto X(\omega, \tau),$$

where:

- For each $\tau \in T$, the mapping $X(\cdot, \tau) : \Omega \rightarrow \mathbb{R}$ is measurable with respect to $\mathcal{F}$.
- For each $\omega \in \Omega$, the mapping $X(\omega, \cdot) : T \rightarrow \mathbb{R}$ is a trajectory (sample path) of the process.

Key Perspectives

- Random variable view: For each $\tau \in T$, $X(\tau)$ is a random variable.
- Sample path view: For each $\omega \in \Omega$, $X(\omega, \cdot)$ is a deterministic function of time.
- Distributional view: The finite-dimensional distributions $X(\tau_1), \dots, X(\tau_n)$ for $\tau_1, \dots, \tau_n \in T$ characterize the process.

Figure 1 illustrates the concept of a stochastic process through the simulation of multiple sample paths of a simple random walk, a canonical example within stochastic process theory. Each trajectory corresponds to a distinct realization of the process, associated with a particular outcome $\omega \in \Omega$. The figure depicts the temporal evolution of the process values across these realizations, thereby exemplifying the intrinsic randomness and variability that are fundamental characteristics of stochastic processes.




Figure 1: Multiple simple random-walk sample paths (5)

A stochastic process formalizes “random evolution over time”: given a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, the process assigns a random variable to each time point and, for a fixed outcome ω , a deterministic sample path. Figure 1 illustrates multiple sample paths (realizations) of a simple random walk. Each trajectory corresponds to a distinct realization. The figure depicts the temporal evolution of the process values across these realizations. It exemplifies the intrinsic randomness and variability that are fundamental characteristics of stochastic processes.

2 Stationary stochastic Processes

2.1 Weak Stationarity

Definition 2.1 (Weak (second-order) stationarity). Let $\{X_t : t \in T\}$ be a stochastic process with finite second moments. The process is said to be weakly stationary (or second-order stationary) if:

1. $\mathbb{E}[X_t] = \mu$ for all $t \in T$ (the mean is constant), and
2. $\text{Cov}(X_{t+h}, X_t) = \gamma(h)$ depends only on the lag h (the autocovariance is a function of lag only), for all t, h for which the moments exist.

Equivalently, weak stationarity implies that $\text{Var}(X_t) = \gamma(0)$ is constant over time. Many linear time-series models with finite variance (for example, ARMA models with appropriate coefficients) are weakly stationary. Weak stationarity is the common assumption used in spectral analysis, autocorrelation estimation, and linear forecasting methods.




Figure 2: Ensemble of 4 stationary AR(1) sample paths; the ensemble mean at t_p is $\bar{X}_{t_p} = \frac{1}{4} \sum_{j=1}^4 X_{t_p}^{(j)}$

Figure 2 plots an ensemble of four independent realizations of a stationary AR(1) process to illustrate how individual sample paths fluctuate around a constant equilibrium while their average tracks the model's theoretical mean: the red horizontal line marks the unconditional mean $\mu = \alpha/(1 - \phi)$, the vertical blue line marks the inspection time τ_p (here t_p), and the blue point gives the ensemble mean at that time $\bar{X}_{t_p} = \frac{1}{4} \sum_{j=1}^4 X_{t_p}^{(j)}$. The intuition is that, although each path exhibits idiosyncratic noise and short-run deviations, stationarity implies their first and second moments do not drift over time, so averaging across independent realizations reduces the noise and yields an estimate centered on μ (with sampling error that decreases as the ensemble size grows, by the law of large numbers).

Weak (or covariance) stationarity refers to a property of a time series whereby, notwithstanding its apparent randomness, its fundamental second-order characteristics remain invariant over time. Specifically, the mean of the series is constant, and the covariance between any two observations depends solely on the temporal distance (lag) separating them rather than on their specific positions in time. Intuitively, if one were to apply a fixed-length moving window across the series and compute summary statistics such as the mean, variance, and autocorrelations at various lags (e.g., 1, 2, or 3), these measures would remain stable and exhibit no systematic drift as the window progresses along the time axis. The series does not exhibit systematic increases in level, heightened volatility, or alterations in its correlation structure over calendar time; rather, it persistently replicates an identical variance–covariance configuration. This condition constitutes a less stringent requirement than full (strong) stationarity, as it disregards the complete distributional form and instead concentrates exclusively on the first and second moments. Such a focus is frequently sufficient for linear modeling frameworks and for a wide range of practical time-series methodologies.

The figure below illustrates a simulated weakly stationary AR(1) process, which is a common

example of a stationary time series. The plot demonstrates the constancy of the mean (indicated by the red dashed line) and the stable variance over time, visually confirming the properties of weak stationarity.

- Constancy of mean
- Stable variance




Figure 3: Simulated stationary AR(1) process (mean dashed)

The figure shows a simulated AR(1) series with coefficient $\phi = 0.8$ and unconditional mean approximately $\mu = 5$. The dashed red horizontal line marks the sample mean, and the dotted green lines mark the bands at approximately ± 2 sample standard deviations. Deviations from the mean are persistent but mean-reverting (due to $\phi < 1$), and the overall variance remains roughly constant over time — together these features visually illustrate weak (second-order) stationarity: a constant mean and a covariance structure that does not drift over time.

Remark 2.2. Weak stationarity requires only constant mean and autocovariance, whereas strong stationarity requires full distributional invariance under time shifts. Although strict stationarity is a stronger condition, weak stationarity is often sufficient in practice, especially for linear models and analyses focused on second-order properties.

2.2 Strong Stationarity

A strongly (strictly) stationary process has the same finite-dimensional distributions under any time shift. This is a stronger requirement than weak stationarity, which only constrains the first two moments; strict stationarity demands full distributional invariance.

Definition 2.3 (strong stationarity). Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let $\{X_t : t \in T\}$ be a stochastic process indexed by $T \subseteq \mathbb{R}$. The process $\{X_t\}$ is said to be strongly stationary (or strictly stationary) if, for every integer $n \geq 1$, for every choice of time

points $t_1, t_2, \dots, t_n \in T$, and for every shift $h \in \mathbb{R}$ such that $t_i + h \in T$, we have $(X_{t_1}, X_{t_2}, \dots, X_{t_n}) \stackrel{d}{=} (X_{t_1+h}, X_{t_2+h}, \dots, X_{t_n+h})$, where $\stackrel{d}{=}$ denotes equality in distribution.

Strong (or strict) stationarity denotes the property whereby the complete probabilistic structure of a stochastic process remains invariant under temporal translation. That is, not only the mean or variance, but the entire joint distribution of any finite collection of time-indexed random variables remains the same under time shifts. Intuitively, if one were to extract numerous short segments of the series—such as triplets or longer contiguous subsequences—and randomly permute their temporal order, it would be impossible to determine whether a given segment originated earlier or later in time. This is because all configurations of values, along with their associated probability distributions, remain invariant under temporal translation. This constitutes a highly stringent requirement: it precludes not only the presence of trends and temporal variation in variability, but also any modification in the distributional form or dependence structure of the data. Consequently, the data-generating process is, in a fundamental sense, temporally invariant.

Key Points

- Entire distribution invariance: Not just the mean, variance, or covariance, but the full joint distribution of any finite collection of random variables remains the same under time shifts.
- Stronger than weak stationarity: Weak stationarity only requires invariance of first and second moments, while strong stationarity requires invariance of all finite-dimensional distributions.
- Implication: Every strong stationary process is weakly stationary (if moments exist), but not every weakly stationary process is strongly stationary.

Intuitively, strict stationarity means that time shifts do not change the joint distribution of observations.

Example 2.4. An everyday example is i.i.d. white noise: independence plus identical marginals imply that every finite-dimensional joint distribution is invariant under time shifts, so the process is strictly stationary.




Figure 4: Gaussian white noise (i.i.d. $\mathcal{N}(0,1)$)

The figure shows a realization of Gaussian white noise: i.i.d. draws from $\mathcal{N}(0, 1)$. The series has (approximately) zero mean, constant variance, and no serial correlation at nonzero lags. Because the observations are independent with identical marginals, this process is a canonical example of a strictly (strongly) stationary process: every finite-dimensional distribution is invariant under time shifts. It is therefore also weakly stationary (second-order) when second moments exist. This strongly stationary example contrasts with the AR(1) above, where temporal dependence yields persistence while still satisfying weak stationarity whenever $|\phi| < 1$.

Example 2.5. The Bernoulli process (i.i.d. Bernoulli(p)) is likewise strictly stationary.

These examples illustrate that identical marginal distributions and independence across time imply strict stationarity.

A classic example of a process that is weakly stationary but not strongly stationary is a non-Gaussian process with time-invariant mean and covariance (often sharing the same covariance structure as a Gaussian stationary process), since in the non-Gaussian case weak stationarity need not imply strong stationarity. To make it concrete, consider the following:

Example 2.6. Define $X_t = Z_t Y$ with Z_t i.i.d. mean-zero noise and $Y \in \{\pm 1\}$ independent of the Z_t . The process has constant mean and lag-dependent covariance (weak stationarity), and finite-dimensional distributions are invariant under time shifts, so it is strictly stationary, as well.

In short: weak stationarity only cares about mean, variance, and covariance stability, while strong stationarity requires full distributional invariance. This example illustrates how a process can meet the weaker conditions without satisfying the stronger ones.

Table 1 presents a compact summary of the series and models analyzed: each row corresponds to a data series or simulated DGP and columns report the sample size, the sample

mean and variance, the estimated AR(1) coefficient (with its standard error), the first few autocovariances/autocorrelations, and the key test statistics with p-values used to assess weak stationarity and unit roots. Together these entries show the degree of persistence and variability for each series, allow quick comparison of estimated dynamics across models (e.g., stationary vs. near-unit-root behavior), and indicate which series reject or fail to reject stationarity at conventional significance levels.

% Replaced pipe-style markdown table with a LaTeX longtable to avoid % layout distortion in PDF output.

Table 1: Summary comparison of weak (second-order) and strong (strict) stationarity: definitions, requirements, implications, and practical considerations.

oprule Aspect	Weak (second-order) stationarity	Strong (strict) stationarity
Definition	Constant mean and covariance depend only on lag: $\mathbb{E}[X_t] = \mu$ and $\text{Cov}(X_{t+h}, X_t) = \gamma(h)$ for all t, h .	All finite-dimensional distributions are invariant under time shifts: for every $k, t_1, \dots, t_k$ and shift h , $(X_{t_1+h}, \dots, X_{t_k+h}) \stackrel{d}{=} (X_{t_1}, \dots, X_{t_k})$.
Required moments	Requires existence of first and second moments.	No moment requirement in the definition.
Condition on distributions	Only first two moments must be time-invariant.	Entire joint distributions must be time-invariant.
Implication relation	Does not imply strong stationarity in general.	Implies weak stationarity if first and second moments exist.
Gaussian case	For Gaussian processes, weak $\Leftrightarrow$ strong (mean covariance determine law).	Same as left column under Gaussianity.
Typical examples	ARMA processes with finite variance.	IID processes; time-homogeneous Markov processes.
Detectability / tests	Tests focus on mean/covariance (ADF, KPSS, sample ACF).	Hard to test directly; compares finite-dimensional distributions.
Practical use	Sufficient for spectral analysis and linear forecasting.	Stronger theoretical guarantees; rarely required in practice.
Vulnerabilities	Can hide nonstationary higher moments or non-Gaussian features.	More robust but harder to verify.

2.3 Example Stationary Processes

2.3.1 White Noise Process

Definition (White noise). Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let $\{\varepsilon_t\}_{t \in \mathbb{Z}}$ be a real-valued stochastic process with finite second moments. The process is a white noise process with variance σ^2 if, for every $t, s \in \mathbb{Z}$,

$$\mathbb{E}[\varepsilon_t] = 0, \quad \mathbb{E}[\varepsilon_t^2] = \sigma^2 < \infty, \quad \text{Cov}(\varepsilon_t, \varepsilon_s) = 0 \text{ for } t \neq s.$$

Equivalently, its autocovariance function is

$$\gamma(h) = \text{Cov}(\varepsilon_t, \varepsilon_{t-h}) = \begin{cases} \sigma^2, & h = 0, \\ 0, & h \neq 0. \end{cases}$$

This is the weak, wide-sense, or second-order notion of white noise: the process is centered, has constant finite variance, and is uncorrelated across distinct times. If the ε_t are also independent and identically distributed (i.i.d.) with mean zero and variance σ^2 , then the process is a strong (strictly) stationary white noise process, because all finite-dimensional distributions are invariant under time shifts. The Gaussian white noise process is a special case where $\varepsilon_t \sim \mathcal{N}(0, \sigma^2)$ i.i.d., which is also both weakly and strongly stationary.

Key Points

Two stronger notions are often separated from that basic definition:

- i.i.d. white noise / strong white noise: ε_t are independent and identically distributed with mean zero and variance σ^2 .
- Gaussian white noise: in addition, the process is jointly Gaussian; in that case, zero covariance already implies independence.

So, when one says:

$$\varepsilon_t \sim \text{i.i.d. } \mathcal{N}(0, \sigma^2),$$

It means that the process is a Gaussian white noise process, which is both weakly and strongly stationary. The i.i.d. assumption ensures strong stationarity, while the Gaussianity implies that the process is fully characterized by its mean and covariance structure.

And, when one says:

$$\varepsilon_t \sim \mathcal{WN}(0, \sigma^2),$$

In time-series analysis, the most cautious assumption is typically the second-order definition mentioned earlier, unless the author clearly specifies that the process is i.i.d. or Gaussian. This difference is important because being uncorrelated is a less restrictive condition than being independent. For instance, a process may display zero autocovariances at every nonzero lag—thereby meeting the criteria for weak white noise—yet still show higher-order dependencies that contradict the i.i.d. assumption. Consequently, in time-series analysis, it is essential to specify which definition of white noise is being used, since this choice affects the behavior of estimators and the reliability of statistical inference.

2.3.2 Moving Average (MA) Processes

Definition (MA(q) process). Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let $\{\varepsilon_t\}_{t \in \mathbb{Z}}$ be a white noise process with variance σ^2 . A stochastic process $\{X_t\}_{t \in \mathbb{Z}}$ is a moving average process of order q (MA(q)) if it can be represented as

$$X_t = \mu + \sum_{j=0}^q \theta_j \varepsilon_{t-j}.$$

Equivalently, in lag-operator notation,

$$X_t = \mu + \theta(B)\varepsilon_t, \quad \theta(B) = \theta_0 + \theta_1 B + \dots + \theta_q B^q.$$

This is the conventional formal definition employed in time-series analysis.

The term “moving average” indicates that the variable X_t is constructed as a finite linear combination of the present and previous shocks $\varepsilon_t, \varepsilon_{t-1}, \dots, \varepsilon_{t-q}$. The source of randomness lies in the white-noise innovations, while the parameters θ_j control the extent to which each shock influences the process over time.

According to this definition, an MA(q) process is inherently weakly stationary, and its autocovariance function satisfies

$$\gamma(h) = 0 \quad \text{for } |h| > q,$$

Thus, correlations disappear for lags greater than q . More precisely,

$$\gamma(h) = \sigma^2 \sum_{j=0}^{q-|h|} \theta_j \theta_{j+|h|} \quad \text{for } |h| \leq q,$$

adopting the convention that $\theta_0 = 1$.

One frequently encountered special instance is the MA(1) process:

$$X_t = \mu + \varepsilon_t + \theta_1 \varepsilon_{t-1}.$$

2.3.3 Autoregressive (AR) Processes

2.3.4 Autoregressive Moving Average (ARMA) Processes

3 Non-stationary Stationary Processes

3.1 Trend Stationary Processes

3.2 Difference Stationary or Integrated Processes (Unit Root Processes)


Figure 5: OLS $\hat{\phi}$ distributions for stationary AR(1) and random walk processes

Table 2: Monte Carlo summary (n=2000, reps=1,000,000)

Process	ϕ	Mean of $\hat{\phi}$	Std of $\hat{\phi}$	Bias of $\hat{\phi}$
Stationary AR(1)	0.7	0.6985	0.0160	-0.0015
Random Walk	1.0	0.9973	0.0023	-0.0027

4 Spurious Regression

A fundamental problem in time-series econometrics arises when nonstationary series are regressed on one another. Even when the underlying stochastic processes are independent, ordinary least squares (OLS) estimation may produce statistically significant relationships. This phenomenon is known as **spurious regression**, first documented by Granger and Newbold (1974) and later formalized by Phillips (1986).

In this section, we combine theory and Monte Carlo simulation to illustrate both the mechanism and the consequences of spurious regression.

4.1 Theoretical Framework

$$X_t = X_{t-1} + u_t, \quad Y_t = Y_{t-1} + v_t,$$

where u_t and v_t are independent white-noise processes with zero mean and finite variance.

We estimate the regression:

$$Y_t = \alpha + \beta X_t + e_t.$$

Since X_t and Y_t are independent, the true relationship is null. However, OLS estimation produces nonstandard behavior.

4.1.1 Brownian Motion Representation

Define the partial-sum processes:

$$\frac{1}{\sqrt{T}}X_{[Tr]} \Rightarrow W_1(r), \quad \frac{1}{\sqrt{T}}Y_{[Tr]} \Rightarrow W_2(r),$$

where $(W_1(r))$ and $(W_2(r))$ are independent Brownian motions on $([0,1])$.

4.1.2 Limit Distribution of the OLS Estimator

Theorem 4.1 (Phillips (1986)). *In the regression of one independent random walk on another, the OLS estimator satisfies*

$$\hat{\beta} \Rightarrow \frac{\int_0^1 W_1(r)W_2(r) dr}{\int_0^1 W_1(r)^2 dr}.$$

Remark 4.2. This limiting distribution is non-normal and depends on functionals of Brownian motion. Importantly, $\hat{\beta}$ itself converges to a nondegenerate distribution—no additional scaling is required.

4.1.3 Divergence of the t-Statistic

Proposition 4.3. *The conventional t-statistic for testing $(H_0 : \beta = 0)$ diverges as $T \rightarrow \infty$, leading to spurious statistical significance.*

This explains why standard inference fails in regressions involving nonstationary variables.

4.2 Monte Carlo Simulation

We simulate two independent random walks and estimate the regression:

$$Y_t = \alpha + \beta X_t + e_t.$$

The design uses:

- Sample sizes: $N = 50, 100, 500, 1000, 10,000, 100,000, 500,000$
- Replications: $R = 20,000$

The Julia code above generates the simulation results, which are then analyzed in R.

4.3 Empirical Findings

4.3.1 Increasing False Significance

The table below shows the percentage of simulations where the null hypothesis of no relationship is rejected at the 5% significance level. As sample size increases, the percentage of significant results approaches 100%, illustrating the spurious regression phenomenon.

Table 3: Spurious regression rejection frequencies by sample size

N	Percent Significant
50	67.9
100	74.5
500	89.5
1,000	91.9
10,000	97.2
100,000	98.8
500,000	99.9

4.3.2 Distribution of the OLS Estimator

After examining the rejection frequencies, we can also look at the distribution of the OLS estimator $\hat{\beta}$ across simulations. The figure below shows the density of $\hat{\beta}$ for different sample sizes. The distribution is for the unscaled OLS estimator $\hat{\beta}$.




Figure 6: Densities of $\hat{\beta}$ (unscaled) across sample sizes

The simulation shows that the distribution of $\hat{\beta}$ appears approximately normal. This is a **finite-sample illusion**. The true asymptotic distribution is non-normal, but finite samples can produce visually symmetric, bell-shaped histograms. The illusion of normality can be misleading, especially when standard inference is applied without recognizing the non-standard asymptotic behavior.




Figure 7: Densities of $N\hat{\beta}$ across sample sizes

4.3.3 Scaled Estimator $N\hat{\beta}$

The density plots above show the distribution of $N\hat{\beta}$. As sample size increases, the distribution becomes more dispersed and non-normal, consistent with the theoretical limit involving Brownian functionals. This illustrates the non-standard behavior of OLS estimators in spurious regressions.

Bibliography

- Brockwell, Peter J., and Richard A. Davis. 1991. *Time Series: Theory and Methods*. 2nd ed. Springer.
- Davidson, James. 1994. *Stochastic Limit Theory*. Oxford University Press.
- Davidson, James. 2000. *Econometric Theory*. Blackwell Publishing.
- Engle, R. F., and C. W. J. Granger, eds. 1991. *Long Run Economic Relationships: Readings in Cointegration*. Oxford University Press.
- Granger, C. W. J., and Paul Newbold. 1974. "Spurious Regressions in Econometrics." *Journal of Econometrics* 2 (2): 111–20.
- Hamilton, James D. 1994. *Time Series Analysis*. Princeton University Press.
- Hansen, Bruce E. 2022. *Econometrics*. Princeton University Press.
- Hayashi, Fumio. 2000. *Econometrics*. Princeton University Press.
- Lütkepohl, Helmut. 2005. *New Introduction to Multiple Time Series Analysis*. Springer.
- Madala, G. S. M., and In-Moo Kim. 1998. *Unit Roots, Cointegration and Structural Change*. Cambridge University Press.
- Neusser, Klaus. 2017. *Time Series Econometrics*. Springer.
- Pesaran, M. Hashem. 2015. *Time Series and Panel Data Econometrics*. Oxford University Press.
- Phillips, Peter C. B. 1986. "Understanding Spurious Regressions in Econometrics." *Journal of Econometrics* 33 (3): 311–40.
- Tsay, Ruey S. 2014. *Multivariate Time Series Analysis: With r and Financial Applications*. Wiley.
- Vance, Martin, Stan Hurn, and David Harris. 2013. *Econometric Modelling with Time Series: Specification, Estimation and Testing*. Cambridge University Press.
- White, Halbert. 2000. *Asymptotic Theory for Econometricians*.

Appendix

Random Walk Sample Paths

AR(1) Time Series Simulation

```
library(ggplot2)
set.seed(123)
n <- 100
library(ggpubr) # Load ggpubr for theme_classic2

# Function to generate a stationary time series
#  $y_t = c + \phi * y_{t-1} + \epsilon_t$ 
ar1_time_series <- function(n, c, phi) {
  # Generate white noise
  epsilon <- rnorm(n, mean = 0, sd = 1)

  # Initialize the time series
  y <- numeric(n)
  y[1] <- c + epsilon[1] # Initial value

  # Generate the time series
  for (t in 2:n) {
    y[t] <- c + phi * y[t - 1] + epsilon[t]
  }

  return(y)
}

# Define n and parameters
n <- 200
c <- 0.2
phi <- 0.6

# Generate an AR(1) time series and plot
data <- data.frame(time = 1:n, time_series = ar1_time_series(n, c, phi))

ggplot(data, aes(x = time, y = time_series)) +
  geom_line(color = "navy") +
  labs(x = "Time", y = "Value") +
  theme_classic2() +
  theme(text = element_text(family = "serif"), plot.title =
    ↪ element_text(hjust = 0.5))
```

Stationary AR(1) Process Plot

```
# Simulate a stationary AR(1) process
set.seed(123)
n <- 500
phi <- 0.8
mu <- 5
sigma <- 1
x <- numeric(n)
x[1] <- rnorm(1, mean = mu, sd = sigma / sqrt(1 - phi^2))
for (t in 2:n) {
  x[t] <- mu + phi * (x[t-1] - mu) + rnorm(1, 0, sigma)
}
library(ggplot2)

mean_x <- mean(x)
sd_x <- sd(x)
upper_band <- mean_x + 2 * sd_x
lower_band <- mean_x - 2 * sd_x

# position for mean label (place near top-right)
y_range <- range(x)
label_y <- y_range[2] - 0.06 * diff(y_range)
label_x <- n - 80
arrow_xend <- n - 60

ggplot(data.frame(Time = 1:n, Value = x), aes(Time, Value)) +
  geom_line(color = 'black', linewidth = 0.6) +
  geom_hline(yintercept = mean_x, linetype = 'dashed', color = 'red',
    ↪ linewidth = 1) +
  geom_hline(yintercept = upper_band, linetype = 'dotted', color =
    ↪ 'darkgreen', linewidth = 0.8) +
  geom_hline(yintercept = lower_band, linetype = 'dotted', color =
    ↪ 'darkgreen', linewidth = 0.8) +
  annotate("curve", x = label_x, xend = arrow_xend, y = label_y, yend =
    ↪ mean_x, curvature = 0.2, colour = 'red', arrow = grid::arrow(length
    ↪ = grid::unit(0.18, 'cm')), size = 0.6) +
  annotate("label", x = label_x, y = label_y, label = expression(E(X[t])
    ↪ == mu), color = 'red', fill = 'white', hjust = 0, vjust = 0.5, size =
    ↪ 3, label.r = grid::unit(0.15, 'lines')) +
```

```

labs(y = expression(X(t)), x = 'Time') +
theme_classic(base_size = 14) +
theme(
  axis.title.x = element_text(size = 10),
  axis.title.y = element_text(size = 10),
  axis.text.x = element_text(size = 8),
  axis.text.y = element_text(size = 8)
) +
scale_x_continuous(breaks = seq(0, n, by = 50)) +
scale_y_continuous(breaks = scales::pretty_breaks(n = 8))

```

Gaussian White Noise

```

set.seed(42)
n <- 500
wn <- rnorm(n, mean = 0, sd = 1)
library(ggplot2)

# compute summary lines
mean_wn <- mean(wn)
sd_wn <- sd(wn)
upper <- mean_wn + 2 * sd_wn
lower <- mean_wn - 2 * sd_wn

ggplot(data.frame(Time = 1:n, Value = wn), aes(x = Time, y = Value)) +
  geom_line(color = 'black') +
  geom_hline(yintercept = mean_wn, linetype = 'dashed', color = 'red',
    ↪ linewidth = 0.6) +
  geom_hline(yintercept = upper, linetype = 'dotted', color = 'darkgreen',
    ↪ linewidth = 0.5) +
  geom_hline(yintercept = lower, linetype = 'dotted', color = 'darkgreen',
    ↪ linewidth = 0.5) +
  labs(x = 'Time', y = expression(epsilon[t])) +
  scale_x_continuous(breaks = seq(0, n, by = 50)) +
  scale_y_continuous(breaks = seq(floor(min(wn)), ceiling(max(wn))), by =
    ↪ 1)) +
  theme_bw() +
  theme(plot.title = element_text(hjust = 0.5))

```

OLS Simulation Summary Table

```

if (file.exists("sim_run_1e6.log")) {
  lines <- readLines("sim_run_1e6.log")

  get_first <- function(pattern) {
    l <- grep(pattern, lines, value = TRUE)
    if (length(l) == 0) return(NA)
    sub(".*:\\s*", "", l[1])
  }

  sample_n <- as.integer(get_first("Sample size"))
  reps <- as.integer(get_first("Replications"))
  burnin <- as.integer(get_first("Burn-in"))

  get_nth_num <- function(pattern, n) {
    idx <- grep(pattern, lines)
    if (length(idx) < n) return(NA)
    as.numeric(sub(".*:\\s*", "", lines[idx[n]]))
  }

  mean_phi_ar1 <- get_nth_num("Mean\\(phi_hat\\)", 1)
  std_phi_ar1 <- get_nth_num("Std\\(phi_hat\\)", 1)
  bias_phi_ar1 <- get_nth_num("Bias\\(phi_hat\\)", 1)

  mean_phi_rw <- get_nth_num("Mean\\(phi_hat\\)", 2)
  std_phi_rw <- get_nth_num("Std\\(phi_hat\\)", 2)
  bias_phi_rw <- get_nth_num("Bias\\(phi_hat\\)", 2)

  true_phi_vals <- as.numeric(sub(".*:\\s*", "", grep("True phi", lines,
↪ value = TRUE)))

  df <- data.frame(
    Process = c("Stationary AR(1)", "Random Walk"),
    True_phi = true_phi_vals,
    Mean_phi_hat = c(mean_phi_ar1, mean_phi_rw),
    Std_phi_hat = c(std_phi_ar1, std_phi_rw),
    Bias_phi_hat = c(bias_phi_ar1, bias_phi_rw)
  )

  library(knitr)
  # Use kableExtra for LaTeX/booktabs styling and scale table to fit page

```

```

if (!requireNamespace("kableExtra", quietly = TRUE)) {
  # kableExtra not available: fall back to pandoc kable silently
  names(df) <- c("Process", "True  $\phi$ ", "Mean of  $\hat{\phi}$ ", "Std
    ↪ of  $\hat{\phi}$ ", "Bias of  $\hat{\phi}$ ")
  # Use a compact pandoc table as a fallback (omit burn-in from caption)
  print(knitr::kable(df, caption = paste0("Monte Carlo summary (n=",
    ↪ sample_n, ", reps=", format(reps, big.mark=','), ")"), digits = 4,
    ↪ format = "pandoc", escape = FALSE))
} else {
  col_names <- c("Process", " $\phi$ ", "Mean of  $\hat{\phi}$ ", "Std of
    ↪  $\hat{\phi}$ ", "Bias of  $\hat{\phi}$ ")
  # Use a plain LaTeX caption (avoid Quarto-style '#'-label which breaks
    ↪ LaTeX when
  # emitting raw LaTeX). If you need a Quarto crossref, switch to a
    ↪ pandoc table.
  # Omit burn-in from LaTeX caption to keep caption concise
  caption_text <- paste0("Monte Carlo summary (n=", sample_n, ", reps=",
    ↪ format(reps, big.mark=','), ")")
  # Produce a LaTeX longtable and scale down to fit the page to avoid
    ↪ distortion
  tbl <- knitr::kable(df, format = "latex", booktabs = TRUE, longtable =
    ↪ TRUE, digits = 4,
    caption = caption_text,
    align = c("l", "r", "r", "r", "r"), col.names = col_names, escape =
    ↪ FALSE)
  tbl <- kableExtra::kable_styling(tbl, latex_options =
    ↪ c("hold_position", "scale_down"), font_size = 8)
  cat(tbl)
}
} else {
  # no simulation log found; remain silent during rendering
}

```

Spurious Regression Table

```

library(readr)
library(dplyr)
library(knitr)

sim <- read_csv("spurious_rw_sim_results.csv", show_col_types = FALSE)

```

```
tbl <- sim |>
  mutate(significant = if_else(!is.na(pval) & pval < 0.05, 1, 0)) |>
  summarize(PercentSignificant = 100 * mean(significant), .by = N) |>
  arrange(N) |>
  mutate(
    N = format(N, big.mark = ",", scientific = FALSE, trim = TRUE),
    PercentSignificant = sprintf("%.1f", PercentSignificant)
  )

kable(
  tbl,
  caption = "Spurious regression rejection frequencies by sample size",
  col.names = c("N", "Percent Significant"),
  align = c("l", "r")
)
```

Unscaled OLS Estimator Density

```
library(readr)
library(dplyr)
library(ggplot2)

sim <- read_csv("spurious_rw_unscaled_sim.csv", show_col_types = FALSE)

plot_df <- sim |>
  filter(N %in% c(50, 500, 10000, 500000)) |>
  mutate(
    N_f = factor(N,
      levels = c(50, 500, 10000, 500000),
      labels = format(c(50, 500, 10000, 500000), big.mark = ",",
        scientific = FALSE, trim = TRUE)
    )
  )

ggplot(plot_df, aes(x = beta)) +
  geom_density(fill = "steelblue", alpha = 0.35, color = "steelblue4",
    linewidth = 0.4) +
  facet_wrap(~N_f, nrow = 2, ncol = 2, scales = "free") +
  labs(x = expression(hat(beta)), y = "Density") +
  theme_classic()
```

Scaled Estimator Density

```

library(readr)
library(dplyr)
library(ggplot2)

sim <- read_csv("spurious_rw_sim_results.csv", show_col_types = FALSE)

plot_df <- sim |>
  filter(N %in% c(50, 500, 10000, 500000)) |>
  mutate(
    N_beta_hat = N * beta,
    N_f = factor(
      N,
      levels = c(50, 500, 10000, 500000),
      labels = format(c(50, 500, 10000, 500000), big.mark = ",", scientific
        ↪ = FALSE, trim = TRUE)
    )
  )

ggplot(plot_df, aes(x = N_beta_hat)) +
  geom_density(fill = "steelblue", alpha = 0.35, color = "steelblue4",
    ↪ linewidth = 0.4) +
  facet_wrap(~N_f, nrow = 2, ncol = 2, scales = "free") +
  labs(x = expression(N * hat(beta)), y = "Density") +
  theme_classic()

```

Julia Simulation: OLS AR(1) vs Random Walk

```
include("simulate_ols_ar1_random_walk.jl")
```

Julia Simulation: Spurious Regression

```

using Random, Distributions, CSV, DataFrames, Statistics

outfile = "spurious_rw_sim_results.csv"
run_sim = get(ENV, "RUN_SPURIOUS_SIM", "0") == "1"

if !run_sim && isfile(outfile)
    @info "Skipping simulation. Using existing file: $outfile"
else

```

```
Random.seed!(2026)
Ns = [50, 100, 500, 1000, 10000, 100000, 500000]
R = 20000
# ... full simulation code as above ...
end
```