

Potentials

Elias Xu

March 8, 2025

Contents

1	Electrical Potentials	1
1.1	Formulas and Dimensional Analysis	2
1.2	Equipotentials	2
1.3	Calculating Electrical Potential from Electric Field	3
2	The Electric Potential of a Point Charge	3
2.1	Calculating the Point Charge's Potential Through Integration	4
3	Electric Flux	4
3.1	Finding formula for Electric Flux	4
3.2	Closed Areas vs. Open Areas	5
4	Gauss's Law	5
4.1	Gauss' Theorem	5
4.2	Nice Gaussian Surfaces	5
4.3	Electrical Fields vs Flux	5
5	Charges on a Conductor	5
5.1	Proof by Gauss' law	6
5.2	Charge inside a cavity	6
6	Electric Fields Infinitely Long Charged Rod	6
7	Electric Fields inside a NONCONDUCTING cylinder	7
7.1	Find E ($r \leq R$)	8
7.2	Find E ($r > R$)	8
8	Solving for stuff with symmetry	8
9	Potentials of a (Uniformly Charged, Infinite) Plane	8
10	Uniformly charged Infinitely Large Slabs with a width	9

1 Electrical Potentials

Electric potentials are measures of the potential energy created by an electric field. You can think of them as such:

Potential: Potential Energy :: Field: Force Energy

For the potential energy of point charges, there is a very big positional element here. As long as you have two charged particles, as long as they want to move somewhere, there will always be potential energy in that system. Usually, systems want to reduce that energy to zero.

1.1 Formulas and Dimensional Analysis

Let's look at the formulas for volts and field, and see what we can make for them. Let's have the formula for volts, and put of the formula for electric field for comparison.

$$V = \frac{U}{q}$$

$$\vec{E} = \frac{\vec{F}}{q}$$

Let's do a quick dimensional analysis too for these *volts*:

$$[V] = \frac{[J]}{[C]}$$

Differences between field and potential:

- Field is a vector quantity, voltage is a scalar quantity
- Field lines are always perpendicular across vector lines

Let's have a visualization changes of potential over time:




Figure 1: The change of charge across a number line

Notes:

- Voltage decreases as one goes through a field lines

1.2 Equipotentials




Figure 2: Example of electric fields vs electric potentials

- Lines across these show the areas of equidistant potentials
- Circular around point charge
- *Perpendicular to field lines*, (e.g. Electric field is the negative gradient of potential), or $E = -\Delta V$

1.3 Calculating Electrical Potential from Electric Field




Figure 4-3 A test charge q_0 moves from point i to point f along the path shown in a nonuniform electric field. During a displacement $d\vec{s}$, an electric force $q_0\vec{E}$ acts on the test charge. This force points in the direction of the field line at the location of the test charge.

Figure 3: Visualization of the different differentials here

To try to find a formula for this, starting from the givens and working up.

$$\vec{F} = q_0\vec{E}$$

$$W = \int \delta W$$

$$\begin{aligned} W &= \int \delta w = \int_i^f \vec{F} \cdot d\vec{s} \\ &= -\Delta U = -q_0\Delta V \end{aligned}$$

$$\Delta v = \boxed{-\int_i^f \vec{E} \cdot d\vec{s}}$$

Given that electric field is a vector, and potentials are scalars, one might have to treat these as partial derivatives of x and y . So we have the original equation:

$$\vec{E} = \frac{Q}{4\pi\epsilon R^2} = -\frac{\delta V}{\delta s}$$

But we have to split it into components:

$$E_x = \frac{\delta V}{\delta x}$$

$$E_y = \frac{\delta V}{\delta y}$$

$$\vec{E} = -\frac{\Delta V}{\Delta S} = \frac{\delta V}{\delta x} + \frac{\delta V}{\delta y}$$

2 The Electric Potential of a Point Charge

Given a ring of radius R , which has a uniformly distributed charge Q , we can infer the following:

- The net electric field in the center is zero (common sense)
- The net electric field at every point is also zero (similar to gravitational field)
- What is the potential inside the ring, given $\Delta V = \int \vec{E} \cdot d\vec{s} = \int 0 \cdot d\vec{s} = \boxed{C}$.

2.1 Calculating the Point Charge's Potential Through Integration

Things to note:

- Potential is a function of charge and radius, or the distance from a point
- Charge creates a scalar potential field
- $u(q_1, q_0) = q_0 V(q_1)$, the relationship between charge and potential energy

To get definite zero, we can set the potential at $r \rightarrow \infty$ to be zero, or $\lim_{r \rightarrow \infty} V(q) = 0$ to be zero.

$$\Delta V = V_f - V_i = 0 - V_i = \int_r^\infty \vec{E} \cdot \delta \vec{s}$$

$$-V_r = - \int_r^\infty \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} [\hat{r} \cdot \delta \vec{s}] = - \int_r^\infty \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \delta r$$

$$V_r = \frac{Q}{4\pi\epsilon r}$$

Note that the property is *scalar*. This makes stuff like addition and subtraction better!

3 Electric Flux

A measure of how many field lines pierces a certain surface / goes through a surface

3.1 Finding formula for Electric Flux

Flux is proportional to

- Cross sectional area of the surface [A]
- Strength of the wires [E]
- orientation of the wires [θ]

Max flux occurs when $\theta = 0^\circ$ (the angle between the normal of the surface and the field lines), and min flux occurs when $\theta = 90^\circ$ (e.g. when normal of the surface and the field lines are perpendicular).




Figure 4: Visualization of max and min flux

Thus, you can compute the flux here:

$$\Phi = \vec{E} \cdot \vec{A} = |\vec{E}| |\vec{A}| \cos \theta$$

This is with the assumption that that

- Plane is regular
- Field is constant

For an irregular plane / field, you can use this formula

$$\Phi = \int \vec{E} \cdot \delta \vec{A}$$

3.2 Closed Areas vs. Open Areas

Gaussian Surfaces are closed figures, and that allows you to find a direction for the area vector, which is always outwards. Some additional points about close areas and flux:

- The flux for an object that is hit by continious equipotentials and doesn't contain any of their own is zero
- For example, if you have flux that goes through a box, the flux is zero




Figure 5: Object with zero flux

4 Gauss' Law

Electric flux gives us **Gauss' Law**, which tells us that the electric flux through a closed surface is proportional to the enclosed charge.

4.1 Gauss' Theorem

Gauss' theorem relates flux to charge, field, and area:

$$\Phi = \frac{q_{\text{enc}}}{\epsilon_0} = \int \vec{E} \cdot \delta \vec{A}$$

This can also be related to potentials, through the calc idea of divergence (iirc)

$$\Delta \cdot E = \frac{\rho}{\epsilon_0}$$

These equations, though, are only good/useful if you have a good gaussian surface

4.2 Nice Gaussian Surfaces

An example is if it is symmetrical to a charged object. An example is a circular surface with a point charge in the center.

$$\begin{aligned}\Phi &= \frac{q_{\text{enc}}}{\epsilon_0} = \int \vec{E} \cdot \delta \vec{A} \\ &= E \int \delta A = \frac{Q}{\epsilon_0} \\ E(4\pi r^2) &= \frac{Q}{\epsilon_0} \\ E &= \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}\end{aligned}$$

(Interesting because this merges both empirical physics with pure mathematics)

4.3 Electrical Fields vs Flux

Electric fields and flux can vary, for but flux you can calculate a total since it's a scalar value

5 Charges on a Conductor

Any charges placed in an insulated conductor would go to the edges of the conductor, none will remain.

5.1 Proof by Gauss' law

Let's solve with a copper (pretty good conductor) and inside it there's a gaussian surface.




Figure 6: Gaussian setup

We have to prove that the field everywhere except the surface is zero, which means that $E_{\text{Gaussian surface}} = 0$, because then, through Gauss' theorem, the charge would then be zero. Let's prove that through negation:

$$E \neq 0 \rightarrow \exists E \rightarrow \exists F_{\text{on any } q} \rightarrow \exists a_{\text{on any } q}$$

An acceleration means that there would be not at equilibrium, which is definitionally false.

5.2 Charge inside a cavity




Figure 7: Gaussian setup

The implications of this is that if there was a charge inside a cavity, it would be balanced out by a charge along the wall of the conductor to make sure that the charge is zero. If the charge inside the surface was q , then the charge on the edge would be $-q$.

6 Electric Fields Infinitely Long Charged Rod

It's really hard the original way, but we can use Gauss's theorem to find the charge. Let's start with a infinitely long charged rod with density λ :





Figure 8: Infinitely long charged rod with charged density λ , z perpendicular distance to point p

Let's construct a Gaussian cylinder (to mimic symmetry of the charged rod) with length L and radius z .



$$\Phi = \Phi_1 + \Phi_2 + \Phi_3 = \Phi_2 \text{ [surface 1 and 3 perpendicular to field]} = \frac{q_{\text{enc}}}{\epsilon_0} = \frac{\lambda L}{\epsilon_0}$$

And for z , the flux is uniform, because of the perpendicularity.

$$\Phi = \int \vec{E} \cdot \delta \vec{A} = EA = 2\pi 2LE = \frac{\lambda L}{\epsilon_0}$$

$$2\pi 2E = \frac{\lambda}{\epsilon_0}$$

$$E = \frac{\lambda}{2\pi\epsilon_0 z}$$

7 Electric Fields inside a NONCONDUCTING cylinder

This is different, because the charge can be anywhere (with a charged conductor, the charge was on the edges by the theorem). Before we solve, let's assume radial symmetry so that bad stuff doesn't happen. Let's first create it and then add a gaussian surface surrounding the point coaxially.




Figure 9: Nonconducting surface with a gaussian surface

The goal is to find E at point P .

$$\Phi_{\text{GS}} = \frac{q_{\text{enc}}}{\epsilon_0}$$

$$a = 2\pi rLE$$

Since the charge can be anywhere, you have to find q_{enc} by integrating.

$$EA = 2\pi rE = \frac{1}{\epsilon_0} \int \rho \delta V$$

There's two options different cases here, one where $r \leq R$, and one where $r > R$.

7.1 Find E ($r \leq R$)

$$2\pi r L E = \frac{1}{\epsilon_0} \int \rho \delta V = \pi r^2 L \frac{\rho}{\epsilon_0}$$


$$E = \frac{\rho r}{2\epsilon_0}$$

7.2 Find E ($r > R$)

$$2\pi r L E = \frac{1}{\epsilon_0} \int \rho \delta V = \frac{\rho}{\epsilon_0} \pi R^2 L$$

$$E = \frac{R^2 \rho}{\epsilon_0 2r}$$

In sum, we get this graph relating E with r :




8 Solving for stuff with symmetry

- No overthinking
- $A_{\text{sphere}} = 4\pi r^2$
- $\delta V_{\text{sphere}} = 4\pi r^2 \delta r$
- $\frac{q_{\text{enc}}}{\epsilon_0} = \int \vec{E} \cdot \delta \vec{A}$

9 Potentials of a (Uniformly Charged, Infinite) Plane

This is definitely impossible, but it describes the behavior that a plane has on an object that is very close to it relative its size.




To find an object with spherical symmetry, you just have to find an object whose ends are parallel to the plane and perpendicular to its normal. To calculate flux:

$$\Phi = \frac{q_{\text{enc}}}{\epsilon_0} = \frac{A\sigma}{\epsilon_0} = 2EA$$

$$E = \boxed{\frac{\sigma}{2\epsilon_0}}$$

10 Uniformly charged Infinitely Large Slabs with a width

Let's illustrate this problem through example. There is an infinitely charged slab with the length of 9.4 mm, with a uniform density of $\rho = 5.80 \frac{\text{fC}}{\text{m}^2}$. What is the field experienced at the distances a) $x = 0$ mm, b) $x = 2$ mm, c) $x = 4.7$ mm, d) $x = 26$ mm.



To solve this problem, you have think of special guassian pillbox with width equal to $2x$ centered at the middle of the space, in order to calculate the amount of charge enclosed.

$$\Phi = \frac{q_{\text{enc}}}{\epsilon_0} = 2EA = \frac{1}{\epsilon_0} \rho V = \frac{2Ax\rho}{\epsilon_0}$$

$$E = \frac{2Ax\rho}{A\epsilon_0 2A} = \frac{x\rho}{\epsilon_0}$$

This then gives you the formula to plug in:

- a) 0
- b) $1.3 \times 10^{-6} \frac{\text{N}}{\text{C}}$
- c) $3.08 \times 10^{-6} \frac{\text{N}}{\text{C}}$
- d) $3.08 \times 10^{-6} \frac{\text{N}}{\text{C}}$ (You don't gather any additional charge from the stuff outside the slab so the field doesn't change.)