Mon do we obtain the valuer

of
$$B = [\beta_0, \beta_1 - - \beta_{p-1}]^T$$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+0} & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+0} & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+0} & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+0} & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+0} & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 & - x_{m-1p-1} \end{bmatrix}$
 $X = \begin{bmatrix} x_{00} \times 0 & - x_{0p-1} \\ \vdots \\ x_{m+1} \times 0 &$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{\tilde{g}})^{T} (\mathbf{g} - \mathbf{\tilde{g}}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{\tilde{g}})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{xp}) \right]$$

$$= \frac{1}{m} \left[(\mathbf{g} - \mathbf{xp})^{T} (\mathbf{g} - \mathbf{x$$

- BPI Ya'p-1) in Matrix vector $\frac{\partial C}{\partial \beta} = 0 = \chi^{T} (g - \chi \beta)$ = x x p = x y $x \in R^{m \times p} x^{T} \in R$ $y \in R^{m}$ if we can invert $\Rightarrow B = (XX)^{-1}XY$ xu x (x2 x3 - - - xm-(