

Final

Fall 2020

Name: Eli Talcott Session (circle one): Morning Evening / DRC

Please copy and write the following statement:

I certify that I have neither given nor received unauthorized aid on this exam.

I certify that I have neither given nor received unauthorized aid on this exam.
(Please copy and write the above statement.)

Eli Talcott

(Signature)

Please state clearly each step you take to reach the final answer. A correct final answer without proper explanations will not receive the full credit.

Problem	Score
Q1	
Q2	
Q3	
Total	

Exercise 1. (30 POINTS)

(a) (6 points)

$$P_X(0) = \frac{1}{2}, P_X(1) = \frac{1}{2}, P_Y(0) = \frac{1}{2}, P_Y(1) = \frac{1}{2}, Z(t) = X_U(t - Y)$$

i) $X = 0, Y = 0$

ii) $X = 0, Y = 1$

iii) $X = 1, Y = 0$

iv) $X = 1, Y = 1$

(b) (8 points)

Find PMF of $Z(\frac{1}{2})$

For case	i) $Z(\frac{1}{2}) = 0$	$P[\text{case}] = \frac{1}{2}(\frac{1}{2}) = \frac{1}{4}$
	ii) $Z(\frac{1}{2}) = 0$	$\frac{1}{4}$
	iii) $Z(\frac{1}{2}) = 1$	$\frac{1}{4}$
	iv) $Z(\frac{1}{2}) = 0$	$\frac{1}{4}$

$$P[Z(\frac{1}{2}) = 1] = \frac{1}{4} \quad P[Z(\frac{1}{2}) = 0] = \frac{3}{4}$$

(c) (8 points)

Find $E[z(t)]$ ($\mu_z(t)$) $X = \text{Bernoulli w/ } p=0.5$
 $Y = \text{Bernoulli w/ } p=0.5$

$$E[z(t)] = \sum_{\text{cases}} X u(t-Y) P[\text{case}]$$

$$= 0\left(\frac{1}{4}\right) + 0\left(\frac{1}{4}\right) + u(t)\left(\frac{1}{4}\right) + u(t-1)\left(\frac{1}{4}\right)$$

$$E[z(t)] = \frac{1}{4}u(t) + \frac{1}{4}u(t-1)$$

(d) (8 points)

Find joint PMF of $z(\frac{1}{2})$ and $z(\frac{3}{2})$

For case	i) $z(\frac{1}{2}) = 0$	$z(\frac{3}{2}) = 0$	$P[\text{case}] = \frac{1}{4}$
	ii) $z(\frac{1}{2}) = 0$	$= 1$	$= \frac{1}{4}$
	iii) $z(\frac{1}{2}) = 1$	$= 0$	$= \frac{1}{4}$
	iv) $z(\frac{1}{2}) = 1$	$= 1$	$= \frac{1}{4}$

$$P[z(\frac{1}{2})=0, z(\frac{3}{2})=0] = \frac{1}{2}, P[z(\frac{1}{2})=0, z(\frac{3}{2})=1] = \frac{1}{4}$$

$$P[z(\frac{1}{2})=1, z(\frac{3}{2})=0] = 0, P[z(\frac{1}{2})=1, z(\frac{3}{2})=1] = \frac{1}{4}$$

Exercise 2. (30 POINTS)

(a) (10 points)

$$f_T(t) = \begin{cases} \lambda e^{-\lambda t}, & t \geq 0 \\ 0, & t < 0 \end{cases} \quad F_T(t) = \begin{cases} 1 - e^{-\lambda t}, & t \geq 0 \\ 0, & t < 0 \end{cases}$$

$$U = \max\{T_1, T_2\}, \quad V = \min\{T_1, T_2\}, \quad \text{Find } f_v(t)$$

$$F_v(t) = (1 - e^{-\lambda T_1})(v(T_1 - T_2)) + (1 - e^{-\lambda T_2})(1 - v(T_1 - T_2)) \quad \begin{matrix} \text{if } T_1 \geq T_2: v(t_1 - t_2) = 1 \\ T_1 < T_2: v(t_1 - t_2) = 0 \end{matrix}$$

~~$$F_v(t) = v(T_1 - T_2) - e^{-\lambda T_1} v(T_1 - T_2) + 1 - v(T_1 - T_2) - e^{-\lambda T_2}$$~~

$$F_v(t) = \max\{(1 - e^{-\lambda T_1}), (1 - e^{-\lambda T_2})\}(t)$$

(b) (12 points)

$$V = \min\{T_1, T_2\}$$

$$F_V(t) = \min\{(1 - e^{-\lambda T_1}), (1 - e^{-\lambda T_2})\}(t)$$

(c) (4 points)

$$p_1 = P[T_1 \geq T_2], p_2 = P[T_1 < T_2]$$

$$E[V] = p_1 \left(\frac{1}{\lambda} \right) \dots$$

(d) (4 points)

Exercise 3. (40 POINTS)

(a) (4 points)

$$X(t, \epsilon) = A(\epsilon)t + B(\epsilon), \quad -\infty < t < \infty$$

$$\mu_X = E[X(t)] = E[A(\epsilon)t + B(\epsilon)] = tE[A(\epsilon)] + E[B(\epsilon)]$$

$$\mu_X(t) = \mu_A t + \mu_B$$

(b) (12 points)

Find $R_X(t_1, t_2)$, Is it WSS?

$$R_X(t_1, t_2) = E[X(t_1)X(t_2)] = E[(A(\epsilon)t_1 + B(\epsilon))(A(\epsilon)t_2 + B(\epsilon))]$$

$$= E[A^2(\epsilon)t_1t_2 + A(\epsilon)t_1B(\epsilon) + A(\epsilon)t_2B(\epsilon) + B^2(\epsilon)]$$

$$= t_1t_2 E[A^2(\epsilon)] + t_1 E[A(\epsilon)B(\epsilon)] + t_2 E[A(\epsilon)B(\epsilon)] + E[B^2(\epsilon)]$$

$$= \mu_A^2 t_1 t_2 + \mu_A \mu_B t_1 + \mu_A \mu_B t_2 + \mu_B^2$$

$$= (\mu_A t_1 + \mu_B)(\mu_A t_2 + \mu_B)$$

Since $\mu_X(t)$ is not constant, X is NOT WSS

(c) (14 points)

Find $M_X(s)$ $X(t) = A(t) + B(t)$

$$M_A(s) = e^{\mu_A s + \frac{\sigma_A^2 s^2}{2}}$$

$$M_B(s) = e^{\mu_B s + \frac{\sigma_B^2 s^2}{2}}$$

A, B are independent For $Z = X + Y$, $M_Z(s) = M_X(s) M_Y(s)$

$$\therefore M_X(s) = e^{\mu_A s + \frac{\sigma_A^2 s^2}{2}} e^{\mu_B s + \frac{\sigma_B^2 s^2}{2}}$$

(d) (5 points)

Find $f_X(t)$

$$F_X(t) = \int_{-\infty}^{\infty} A(\epsilon)t + B(\epsilon) d\epsilon$$

$$= \left[\frac{1}{2} A(\epsilon)t^2 + B(\epsilon)t \right]_{-\infty}^{\infty}$$

(e) (5 points)

Find $P[|X(t)| > \alpha]$ where $\alpha > 0$

Useful Identities

- $\sum_{k=0}^{\infty} r^k = 1 + r + r^2 + \dots = \frac{1}{1-r}$
- $\sum_{k=1}^n k = 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$
- $e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \dots$
- $\sum_{k=1}^{\infty} kr^{k-1} = 1 + 2r + 3r^2 + \dots = \frac{1}{(1-r)^2}$
- $\sum_{k=1}^n k^2 = 1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n^3}{3} + \frac{n^2}{2} + \frac{n}{6}$
- $(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$

Common Distributions

Bernoulli	$\mathbb{P}[X = 1] = p$	$\mathbb{E}[X] = p$	$\text{Var}[X] = p(1-p)$	$M_X(s) = 1 - p + pe^s$
Binomial	$p_X(k) = \binom{n}{k} p^k (1-p)^{n-k}$	$\mathbb{E}[X] = np$	$\text{Var}[X] = np(1-p)$	$M_X(s) = (1 - p + pe^s)^n$
Geometric	$p_X(k) = p(1-p)^{k-1}$	$\mathbb{E}[X] = \frac{1}{p}$	$\text{Var}[X] = \frac{1-p}{p^2}$	$M_X(s) = \frac{pe^s}{1-(1-p)e^s}$
Poisson	$p_X(k) = \frac{\lambda^k e^{-\lambda}}{k!}$	$\mathbb{E}[X] = \lambda$	$\text{Var}[X] = \lambda$	$M_X(s) = e^{\lambda(e^s-1)}$
Gaussian	$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	$\mathbb{E}[X] = \mu$	$\text{Var}[X] = \sigma^2$	$M_X(s) = e^{\mu s + \frac{\sigma^2 s^2}{2}}$
Exponential	$f_X(x) = \lambda \exp\{-\lambda x\}$	$\mathbb{E}[X] = \frac{1}{\lambda}$	$\text{Var}[X] = \frac{1}{\lambda^2}$	$M_X(s) = \frac{\lambda}{\lambda-s}$
Uniform	$f_X(x) = \frac{1}{b-a}$	$\mathbb{E}[X] = \frac{a+b}{2}$	$\text{Var}[X] = \frac{(b-a)^2}{12}$	$M_X(s) = \frac{e^{sb} - e^{sa}}{s(b-a)}$

Fourier Transform Table

$f(t) \longleftrightarrow F(w)$	$f(t) \longleftrightarrow F(w)$
1. $e^{-at}u(t) \longleftrightarrow \frac{1}{a+jw}, a > 0$	10. $\text{sinc}^2(\frac{Wt}{2}) \longleftrightarrow \frac{2\pi}{W} \Delta(\frac{w}{2W})$
2. $e^{at}u(-t) \longleftrightarrow \frac{1}{a-jw}, a > 0$	11. $e^{-at} \sin(w_0 t)u(t) \longleftrightarrow \frac{w_0}{(a+jw)^2 + w_0^2}, a > 0$
3. $e^{-a t } \longleftrightarrow \frac{2a}{a^2 + w^2}, a > 0$	12. $e^{-at} \cos(w_0 t)u(t) \longleftrightarrow \frac{a+jw}{(a+jw)^2 + w_0^2}, a > 0$
4. $\frac{a^2}{a^2 + t^2} \longleftrightarrow \pi a e^{-a w }, a > 0$	13. $e^{-\frac{t^2}{2\sigma^2}} \longleftrightarrow \sqrt{2\pi}\sigma e^{-\frac{\sigma^2 w^2}{2}}$
5. $te^{-at}u(t) \longleftrightarrow \frac{1}{(a+jw)^2}, a > 0$	14. $\delta(t) \longleftrightarrow 1$
6. $t^n e^{-at}u(t) \longleftrightarrow \frac{n!}{(a+jw)^{n+1}}, a > 0$	15. $1 \longleftrightarrow 2\pi\delta(w)$
7. $\text{rect}(\frac{t}{\tau}) \longleftrightarrow \tau \text{sinc}(\frac{w\tau}{2})$	16. $\delta(t - t_0) \longleftrightarrow e^{-jw t_0}$
8. $\text{sinc}(Wt) \longleftrightarrow \frac{\pi}{W} \text{rect}(\frac{w}{2W})$	17. $e^{jw_0 t} \longleftrightarrow 2\pi\delta(w - w_0)$
9. $\Delta(\frac{t}{\tau}) \longleftrightarrow \frac{\tau}{2} \text{sinc}^2(\frac{w\tau}{4})$	

Some definitions:

$$\text{sinc}(t) = \frac{\sin(t)}{t} \quad \text{rect}(t) = \begin{cases} 1, & -0.5 \leq t \leq 0.5, \\ 0, & \text{otherwise.} \end{cases} \quad \Delta(t) = \begin{cases} 1 - 2|t|, & -0.5 \leq t \leq 0.5, \\ 0, & \text{otherwise.} \end{cases}$$

Basic Trigonometry

$$e^{j\theta} = \cos \theta + j \sin \theta, \quad \sin 2\theta = 2 \sin \theta \cos \theta, \quad \cos 2\theta = 2 \cos^2 \theta - 1.$$

$$\begin{aligned} \cos A \cos B &= \frac{1}{2}(\cos(A+B) + \cos(A-B)) & \sin A \sin B &= -\frac{1}{2}(\cos(A+B) - \cos(A-B)) \\ \sin A \cos B &= \frac{1}{2}(\sin(A+B) + \sin(A-B)) & \cos A \sin B &= \frac{1}{2}(\sin(A+B) - \sin(A-B)) \end{aligned}$$