

## Genome analysis

# rCGH: a comprehensive array-based genomic profile platform for precision medicine

Frederic Commo<sup>1,2,\*</sup>, Justin Guinney<sup>2</sup>, Charles Ferté<sup>1,2</sup>, Brian Bot<sup>2</sup>,  
Celine Lefebvre<sup>1</sup>, Jean-Charles Soria<sup>1,3</sup> and Fabrice André<sup>1,3</sup>

<sup>1</sup>INSERM U981, Gustave Roussy, University Paris-Sud, Villejuif 94805, France, <sup>2</sup>Sage Bionetworks, Seattle, WA 98109, USA and <sup>3</sup>Department of Medical Oncology, Gustave Roussy, Villejuif 94805, France

\*To whom correspondence should be addressed.

Associate Editor: John Hancock

Received on 7 September 2015; revised on 2 December 2015; accepted on 3 December 2015

## Abstract

**Summary:** We present rCGH, a comprehensive array-based comparative genomic hybridization analysis workflow, integrating computational improvements and functionalities specifically designed for precision medicine. rCGH supports the major microarray platforms, ensures a full traceability and facilitates profiles interpretation and decision-making through sharable interactive visualizations.

**Availability and implementation:** The rCGH R package is available on bioconductor (under Artistic-2.0). The aCGH-viewer is available at [https://fredcommo.shinyapps.io/aCGH\\_viewer](https://fredcommo.shinyapps.io/aCGH_viewer), and the application implementation is freely available for installation at [https://github.com/fredcommo/aCGH\\_viewer](https://github.com/fredcommo/aCGH_viewer).

**Contact:** frederic.commo@gustaveroussy.fr

**Supplementary information:** Supplementary data are available at *Bioinformatics* online.

## 1 Introduction

Precision medicine aims at identifying individual cancer patients' molecular alterations—such as copy number alterations (CNAs), gene mutations or fusions and protein expressions—and matching these to targeted therapies (André *et al.*, 2014; Tsimberidou *et al.*, 2014). While next-generation sequencing is now widely used for identifying mutations, array-based comparative genomic hybridization (aCGH) is a common platform for detecting CNAs (André *et al.*, 2013; Laurent-Puig *et al.*, 2009): the advantages of aCGH over next-generation sequencing technologies include lower cost, rapid turn-around time and lower computational overhead. In the context of precision medicine, CNA analysis—alongside somatic mutation analysis—is critical for identifying clinically actionable genomic aberrations. However, significant technical challenges remain in the processing of aCGH data. Addressing these challenges requires new state-of-the-art tools for coordinating analysis and interpreting results.

## 2 Methods and implementation

An aCGH analysis can be decomposed into four distinct phases (Supplementary Fig. S1): (i) log<sub>2</sub> relative ratios (LRR) calculation (the sample DNA signals against a normal 2-copy DNA reference), (ii) profile centralization, (iii) profile segmentation and (iv) genomic profile interpretation to identify actionable genes affected by a CNA, to propose a matched therapeutic orientation.

The profile centralization defines a baseline—a neutral 2-copies level—from which CNA are estimated. We previously discussed the impact of centralization on aCGH analysis (Commo *et al.*, 2015), and rCGH implements the procedure described in the same article. Briefly, the vector of LRRs is considered as a mixture of Gaussian populations, and their respective proportion and parameters are estimated using an Expectation-Maximization algorithm. By default, the sub-population with a density peak higher than 50% of the highest density is considered as representing a neutral 2-copy state. Its mean is then used for centralizing the profile.



- Laurent-Puig,P. *et al.* (2009) Analysis of PTEN, BRAF, and EGFR status in determining benefit from cetuximab therapy in wild-type KRAS metastatic colon cancer. *J. Clin. Oncol.*, **27**, 5924–5930.
- Mermel,C.H. *et al.* (2011) GISTIC2.0 facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers. *Genome Biol.*, **12**, R41
- Olshen,A.B. *et al.* (2004) Circular binary segmentation for the analysis of array-based DNA copy number data. *Biostatistics*, **5**, 557–572.
- Tsimberidou,A.M. *et al.* (2014) Personalized medicine for patients with advanced cancer in the phase I program at MD Anderson: validation and landmark analyses. *Clin. Cancer Res.*, **20**, 4827–4836.
- Willenbrock,H. and Fridlyand,J. (2005) A comparison study: applying segmentation to array CGH data for downstream analyses. *Bioinformatics*, **21**, 4084–4091.
- Yuan,X. *et al.* (2012) Comparative analysis of methods for identifying recurrent copy number alterations in cancer. *PLoS One*, **7**, e52516.