


Gestion d'une base de donnée qualitatives et géométrique



Elina Marveaux

Master of Science in Biology
Biodiversity, Evolution and Ecology
Department of Biological Sciences
University of Bergen

12/05/2022
ANR DFG
AGENCE NATIONALE DE LA RECHERCHE
Deutsche Forschungsgemeinschaft

Cover page
Beckwithia glacialis on Snøhetta.

Table des matières

Preface	1
1. Summary	3
2. Introduction	5
3. “Penguins”	7
3.1. Plot	7
3.2. Data	8
References	11
4. Summary	13
Appendices	13
A. More results	15

List of Figures

1.	Chart Bar des NAS	1
A.1.	cover	15

List of Tables

Preface

This is a Quarto book.

To learn more about Quarto books visit <https://quarto.org/docs/books>.

[1] 2




Figure 1.: Chart Bar des NAS

1. Summary

In summary, this book has no content whatsoever.

[1] 2

2. Introduction

R table

```
1 data <- data.frame(year = c("1958", "1959", "1960"),
2                     max = c(317.51, 318.29, 320.04))
3
4 # ojs_define(tableR = data)
5
6 data
```

	year	max
1	1958	317.51
2	1959	318.29
3	1960	320.04

OJS line_chart from tutorial with ojs_define() and R table

```
1 yearlyChart = Plot.plot({
2   marks: [
3     Plot.line(tableR,
4       {x: "year", y: "max"},
5       { stroke: "red" }
6     )
7   ]}
8 )
```

Second bar chart with python table

```
1
2 Plot.rectY(tablePy, Plot.binX({y: "max"}, {x: "year", fill: "year"})).plot()
```


3. “Penguins”

A simple example based on Allison Horst’s Palmer Penguins dataset. Here we look at how penguin body mass varies across both sex and species (use the provided inputs to filter the dataset by bill length and island):

```
1  ///| echo = false
2  data = FileAttachment("data/palmer-penguins.csv").csv({ typed: true })

1  ///| echo = false
2  viewof bill_length_mm = Inputs.range(
3    [32, 50],
4    {value: 35, step: 1, label: "Bill length (min):"}
5  )
6  viewof islands = Inputs.checkbox(
7    ["Torgersen", "Biscoe", "Dream"],
8    { value: ["Torgersen", "Biscoe"],
9      label: "Islands:"
10   }
11 )

1  ///| echo: false
2  filtered = data.filter(function(penguin) {
3    return bill_length_mm < penguin.bill_length_mm &&
4           islands.includes(penguin.island);
5  })
```

3.1. Plot


```
1  ///| echo: false
2  ///| output: true
3  ///| label: fig-penguin-body-mass
4  ///| fig-cap: "Penguin body mass by sex and species"
5  Plot.rectY(filtered,
```

3. “Penguins”

```
6   Plot.binX(  
7     {y: "count"},  
8     {x: "body_mass_g", fill: "species", thresholds: 20}  
9   )  
10  .plot({  
11    facet: {  
12      data: filtered,  
13      x: "sex",  
14      y: "species",  
15      marginRight: 80  
16    },  
17    marks: [  
18      Plot.frame(),  
19    ]  
20  }  
21 )
```

3.2. Data

The penguins data from the **palmerpenguins** package contains size measurements for 344 penguins from three species observed on three islands in the Palmer Archipelago, Antarctica./ The plot below shows the relationship between flipper and bill lengths of these penguins.



References

4. Summary

In summary, this book has no content whatsoever.

[1] 2

A. More results

Some results that wouldn't fit into the main thesi

Some results that wouldn't fit into the main thesis




Figure A.1.: cover

