

Mean-Field Study of Kondo Phase Diagram

Elis Roberts

April 29, 2018

University of Cambridge

Plan

1. Introduction
2. Finite Temperature Study
3. Finite Field Study
4. Conclusion

Introduction

Kondo Model

Single localised spin interacting with conduction electrons $c_{k,\sigma}$

Model Hamiltonian

$$H_{\text{Kondo}} = \sum_{k,\sigma} \epsilon_k c_{k,\sigma}^\dagger c_{k,\sigma} + J \vec{S} \cdot \vec{S}(0)$$

Can map spin DOFs of impurity onto fermions f_σ

Path Integral → Mean-Field Theory

- Project framed in terms of path integral
- Treats Boltzmann sum as a functional integral over field configurations

Many-Body Path Integral

$$Z = \text{Tr} e^{-\beta H} = \int \mathcal{D}[c^\dagger, c] e^{-\int_0^\beta d\tau L}$$

- Similar to Feynman path integral approach, but with *imaginary time* $\tau = it/\hbar$ and associated Lagrangian L

Mean-Field Theory


Avoid this difficult integral by making stationary phase approximation i.e. *minimise*

New Fields

- Lagrangian similar to original Hamiltonian, with extra dynamical terms: $L = H_{\text{Kondo}} + c^\dagger \partial_\tau c$
- **But**, interaction term introduces difficult non-quadratic terms

Resolution

Introduce a new field V that mediates this interaction:



Constraints

- Making such variable changes sometimes requires **constraints**

e.g. Spin-1/2 \rightarrow Fermion Mapping

$$s_z = \frac{1}{2}(f_{\uparrow}^{\dagger}f_{\uparrow} - f_{\downarrow}^{\dagger}f_{\downarrow}), \quad s_+ = f_{\uparrow}^{\dagger}f_{\downarrow}, \quad s_- = f_{\downarrow}^{\dagger}f_{\uparrow}$$

Representation isn't faithful unless we impose the constraint:

$$f_{\uparrow}^{\dagger}f_{\uparrow} + f_{\downarrow}^{\dagger}f_{\downarrow} = 1$$

- These constraints are implemented by adding Lagrange multipliers $\{\lambda_i\}$ and extremizing wrt these too - e.g.

$$L \supset \lambda_{\text{RN}}(f_{\uparrow}^{\dagger}f_{\uparrow} + f_{\downarrow}^{\dagger}f_{\downarrow} - 1)$$

Soft-Constraint Approach

- Now consider rewriting the same constraint:

$$(1 - n_{\uparrow} - n_{\downarrow})^2 = n_{\uparrow}n_{\downarrow} + (1 - n_{\uparrow})(1 - n_{\downarrow}) = 0$$

- **But** this can't be done at mean-field level since

$$\langle (1 - n_{\uparrow} - n_{\downarrow})^2 \rangle \neq 0$$

Soft-Constraint

How about introducing a new fermion h obeying $h^{\dagger}h = 0$, but combining these constraints by instead imposing

$$(1 - n_{\uparrow} - n_{\downarrow})^2 - Kh^{\dagger}h = 0 \quad ?$$

Soft-Constraint Approach

- How could $(1 - n_{\uparrow} - n_{\downarrow})^2 - Kh^{\dagger}h = 0$ be any better?
- Have introduced a new free parameter K into description
- Provided $K > 0$ and $K \neq 1$, constraint operator now has positive *and* negative eigenvalues:

$$\{ 1, (1-K), 0, -K \}$$

Consequence

New composite constraint picks out both constraints, while still being possible at mean-field level

Finite Temperature Study

Solving Mean-Field Equations

Generate MF equations via minimisation:

$$\frac{\partial F_{\text{MF}}}{\partial \Delta} = 0, \quad \frac{\partial F_{\text{MF}}}{\partial \lambda_{\text{SC}}} = 0, \quad \dots$$

Deriving Heat Capacity

Unavoidable Phase Transition?

Behaviour of Order Parameter Δ

Finite Field Study

Incorporating $B \neq 0$

Increased Difficulty of Mean-Field Equations

Conclusion

Questions?

Backup

Final Lagrangian

$$\begin{aligned} L = & \sum_{k,\sigma} c_{k,\sigma}^\dagger \left(\frac{d}{d\tau} + \epsilon_k - \mu \right) c_{k,\sigma} + \sum_{\sigma} f_{\sigma}^\dagger \frac{d}{d\tau} f_{\sigma} + h^\dagger \frac{d}{d\tau} h \\ & + e^\dagger \frac{d}{d\tau} e + \sum_{\sigma} p_{\sigma}^\dagger \frac{d}{d\tau} p_{\sigma} + d^\dagger \frac{d}{d\tau} d \\ & + \sum_{\sigma} \lambda_{\sigma} (f_{\sigma}^\dagger f_{\sigma} - p_{\sigma}^\dagger p_{\sigma} - d^\dagger d) \\ & + \lambda_{\text{KR}} (e^\dagger e + \sum_{\sigma} p_{\sigma}^\dagger p_{\sigma} + d^\dagger d - 1) + \lambda_{\text{SC}} (e^\dagger e + d^\dagger d - K h^\dagger h) \\ & + 2 \frac{V V^*}{J} + \sum_{k,\sigma} \left(V^* c_{k,\sigma}^\dagger z_{\sigma} f_{\sigma} + V f_{\sigma}^\dagger z_{\sigma}^\dagger c_{k,\sigma} \right) \end{aligned} \tag{1}$$