

X-ray Vision: A Very Different View of the Universe

Elisea Jackson^a

^a*Institute for Computing in Research,*

Abstract

In the night sky there are many things that can't be seen with the naked eye. Visible light only makes up a small portion of the electromagnetic spectrum. Using satellites we are able to look at gamma rays and x-rays emissions for stars. Looking at these readings we can see what is happening hundreds of light years away. I created a model representing what the sky would look like if we could see several of the X-rays sources from earth with our own eyes. This includes soft gamma repeaters, X-ray bursters, and X-ray pulsars.

1. Introduction

In the model representing the X-ray sky there are three different dynamic sources of Energy: soft gamma repeaters, X-ray pulsars, and X-ray bursters. The one thing that all of these stars have in common is that they originate from Neutron Stars. When a giant star collapses on itself it can form either a black hole or a Neutron star. Neutron stars are formed from the smaller stars undergoing a supernova.

There are two types of Neutron stars: magnetars and pulsars. A magnetar has an extremely strong magnetic field. Its magnetic field can range from 10^{13} gauss to 10^{15} gauss. Magnetars have the most powerful magnetism of any object in the universe. The second type of neutron star is a pulsar. Another type of neutron star is a pulsar. The pulsars emit x-rays from their poles. The magnetic field of the neutron star accelerates the particles forming jets of light to appear from the pulsar's poles. As the poles go in and out of earth's view this creates a lighthouse effect and they appear to be pulsing. The crab pulsar is a famous pulsar with a steady rotation of 33 milliseconds. X-ray pulsars have a quicker rotation than magnetars however, they have a weaker magnetic field.

2. Methods

In my model of the are some steady X-ray emitting stars and some dynamic X-ray stars. The dynamic stars shown in my model are soft gamma repeaters, X-ray pulsars, and X-ray burst. One thing these stars have in common in that they all occur on a neutron star. Most Neutron stars are too dim to be detected. Only extremely active neutron stars with quick rotational period and a strong magnetic field can be detected. These are Magnetars which are Neutron stars with a strong magnetic field. Then there are pulsars which don't contain as strong of a magnetic field but, have a quicker rotational period usually a fraction of a second. In order to model the neutron star I used random probability distribution function to determine different attributes of the neutron star. These include the amplitude and frequency. What is Monte Carlo? Using Monte Carlo method we can create a sky with a similar view of the universe. When modeling high energy objects in space it is hard to account for every variable. In order to make up for lack of computational power and time we can use probability distributions to help. This helps create scenarios that are unique but probable based on what we currently know about these objects.

Using probability distribution – probability density function – I can simulate random situations using probability. Using probability distribution I can specify a mean and standard deviation: represented by μ and σ . If

the model were run several hundred times you would see a distribution of values similar to that of a gaussian bell curve with the majority of values around the mean and less and less values closer to the ends. This then can be translated into a cumulative distribution function. Using this function, when selecting a random number between 0 and 1 it will give you a value represented in the probability density function. The closer the random number is to one or zero the more extreme the value is. If a random number was picked and plotted using the probability density function several hundred times that plot would look similar to the graph created by the probability density function.

In my model to accomplish this I use the inverse error function in python. I select a random value between -1 and 1 and run the inverse error function on it. Then I add the mean and multiply it by the standard deviation.

The problem with this is that not everything is represented by a normal probability distribution. There could be more values on one side than the other. This is usually because the mean, median, and mode are not always the same value. In a normal probability distribution the mean, median, and mode are all the same value.

2.1. Optical Night Sky

In astronomy stars are shown using the Galactic Coordinate system. In this system the Latitude (b) goes from 90 to -90. Longitude (l) starts at 0 in the center and extends left to 180. This continues on the furthest right side of the graph going from 180 to 360 back at the center of the model. In my model I changed the longitude to extend from 180 to -180 to make it easier to pick values gathered around the center of the galaxy. In my model there is a drop down menu with a couple of options. The Static Sky option shows a model of the sky as we normally observe it. After sixty seconds you can switch the screen you are looking at by clicking the clear all button and viewing the X-ray sky. This shows the X-ray sources in the night sky. There are three different distributions of stars in the static star sky. There are stars that are randomly placed over the whole sky. The majority of stars are centered around the milky way. A large number of stars are centered around the galactic center. There is one dynamic star in the static night sky; there is a flare star located longitude (l) 100 and latitude (b) 30. This flare star is a very active star releasing lots of energy. As shown in the model we can't see the change in star size because the star is too far away and




Figure 1: Probability Density Function




Figure 2: Physics Letters B journal cover

the change is not enough. If you would like to see this star magnified in the menu there's a button called Flare star. This button shows another movie with the flare star magnified by a factor of ten so you can see the effects of the flare star. This gives us a general representation of the stars in the milky galaxy.

2.2. The Night Time Sky

2.3. Soft Gamma Ray Bursts

Soft Gamma Repeaters are Neutron Stars called Magnetars. Magnetars have an extraordinarily strong magnetic field ranging from 10^{14} to 10^{15} . Contrary to their name Soft Gamma Repeaters emit large amounts of energy number. Only when compared to a Gamma Ray Burst do Soft Gamma Repeaters seem dim. When these Neutron stars experience an (starquake) they produce a strong emission of energy. Starquakes are similar to that of an earth quake on earth. This is what satellites see as a Soft Gamma Repeater. Satellite reading will show a

pulsation in gamma readings. The neutron star itself isn't pulsing. After the initial eruption the energy disapates. When this is combined with the quick rotational period of magnatars it creates a light house effect. In my model I used the euation represented below to model soft gamma repeaters. AMP is the aplitude of the soft gamma repeater. f is the frequency or the rotational period. t is the start time or the time which the first soft gamma repeater apeared. t_1 is the current time.

2.4. X-ray pulsars

X-ray pulsars are formed because of the acceleration and (coliding) of matter at the poles of the neutron star. The poles of the netoron star rotate around an axis. We see neutron star only when the poles are pointed tword earth. Since to oples move twards and away form earth in a circle it looks like it is puling from earths point of view. In order to model this I used the equation below it is simmlar to the equation show in Neutron stars however it does not have a rate of decay. The burst on a soft gamma repeater only last for a short period of time with larger burst lasting (several minutes). X-ray pulasrs last for (around 10,000 years)(sitation). The time scale of my model is only 1 year makeing it unresonable to add a rate of decay to the X-ray pulsars observed in my model.

X Are binary star systems.

A random equation, the Toomre stability criterion:

2.5. X-ray Bursters

X-ray bursters occur when a neutron star accreates matter on the surface of the star causing the surface of the star to explode. This additional matter is usually acreated form a companion donner star. Thies star orbit around eachoter sloly matter is tranfered form the donner star to the Neutron star. This matter forms an acreation disk around the center of the star. In my model I show these stars as already apearing in the night sky. (As steady X-ray stars that sudenly burst) The burst happens randomly however thier is a correlation between how large the last burst was and when the next burst will occur. This is because the larger the burst mas the more energy(feul) it burned and the more time it will take to accumalate fuel back to the state it once was in. In my model I accomplish this by multiplying the amplitude of the burst times a random number to determine the time in which the next burst will happen.

2.6. Model of Sky in Optical Light

My model shows the view of the galaxy from earth using the galactic coordinate system. In the galatic coordinate system the center of the milky way is the center of the map. the majority of the stars are along the center of the map where the milky way is.(0 longitute) The stars which are futher from this center are stars that are closer to earth or bright object out side the milky way galaxy.

$$Q = \frac{\sigma_v \times \kappa}{\pi \times G \times \Sigma} \quad (1)$$

3. Summary and conclusions

3.1. Succuss

My model is a simple representation of some of the X-ray emitting stars in our galaxy. It shows how dynamic the universe is. Aside from the occasional passing comet our eyes fail to inform us of many of the changes occurring. It helps us to have a better understanding of the universe and interpret information received from satellites.

3.2. Limitations

One of the biggest constraints of this project was time. The model is relatively simple and gives a direct translation of what satellites receive to what it would look like in the sky. Models are important for helping us to improve our understanding of the universe. To improve the model it should model the actual mechanics of the stars and how different variables can affect the output – gamma and x-ray readings received by earth – to help us better interpret those readings. A monte carlo is a good model for something with so many variables; however, it then no longer is a view of our universe but a similar one. It is impossible to model every single atom in a neutron star. It would require too much computational power from a computer.

3.3. Futurework

In the future I would like to add different types of stars that also emit X-ray. I would like to experiment with different time frames. With my current time frame of a year within 60 seconds we miss out on some of the faster spinning objects. For example, millisecond pulsars have a spin speed faster than I could show in my presentation. I would also like to make my model more user friendly

by making the user able to end the movie when they like as well as make the model able to replay the X-ray pulsar which is currently impossible.

Acknowledgements

Thanks to ...

References

- Duncan, R. C. (2003). ‘magnetars’, soft gamma repeaters. ‘MAGNETARS’, SOFT GAMMA REPEATERS & VERY STRONG MAGNETIC FIELDS Robert C. Duncan, University of Texas at Austin. <http://solomon.as.utexas.edu/magnetar.html>
- Galactic coordinate system: Cosmos. Galactic Coordinate System — COSMOS. (n.d.). <https://astronomy.swin.edu.au/cosmos/G/Galactic+Coordinate+System>
- Jenner, L. (2016, July 7). NASA’s Hubble captures the beating heart of the Crab Nebula. NASA. <https://www.nasa.gov/feature/goddard/2016/nasa-hubble-captures-the-beating-heart-of-the-crab-nebula>
- Lewin, W. H., Paradijs, J. V., & Taam, R. E. (1993). X-Ray Bursts. 1993SSRv...62..223L page 227. <https://articles.adsabs.harvard.edu/full/1993SSRv...62..223L/0000227.000.html>
- Reddy, F. (2023, March 27). NASA missions study what may be a 1-in-10,000-year gamma-ray burst. NASA. <https://www.nasa.gov/feature/goddard/2023/nasa-missions-study-what-may-be-a-1-in-10000-year-gamma-ray-burst>
- Soft gamma repeater: Cosmos. Soft Gamma Repeater — COSMOS. (n.d.-a). <https://astronomy.swin.edu.au/cosmos/s/Soft+Gamma+Repeater#:~:text=Such%20a%20rapid%20spin%2Ddown,become%20anomal>
- Soft gamma repeater: Cosmos. Soft Gamma Repeater — COSMOS. (n.d.-b). <https://astronomy.swin.edu.au/cosmos/s/Soft+Gamma+Repeater>
- Sokol, J. (2021, April 8). Nature’s most magnetic objects, ripped apart in starquakes, can unleash ... A Star is Torn. <https://www.science.org/content/article/nature-s-most-magnetic-objects-ripped-apart-starquakes-can-unleash-powerful-flashes>
- Wikimedia Foundation. (2023a, April 4). Sagittarius A. Wikipedia. https://en.wikipedia.org/wiki/Sagittarius_A
- Wikimedia Foundation. (2023b, April 7). X-ray binary. Wikipedia. https://en.wikipedia.org/wiki/X-ray_binary
- Wikimedia Foundation. (2023c, July 22). Normal distribution. Wikipedia. https://en.wikipedia.org/wiki/Normal_distribution
- Wikimedia Foundation. (2023d, July 23). Gamma-ray burst. Wikipedia. https://en.wikipedia.org/wiki/Gamma-ray_burst
- Wikimedia Foundation. (2023e, July 24). Magnetar. Wikipedia. <https://en.wikipedia.org/wiki/Magnetar>
- Wikimedia Foundation. (2023f, July 25). Neutron star. Wikipedia. https://en.wikipedia.org/wiki/Neutron_star
- Wikimedia Foundation. (2023g, August 4). X-ray pulsar. Wikipedia. https://en.wikipedia.org/wiki/X-ray_pulsar#:~:text=X%2Dray%20pulsars%20or%20accretion,as%20much