



Predictive Modeling of Root-Zone Variables Using Greenhouse Data

Students: Eden Eliyahu, Oren Chaushu

Supervisors: Dr. Avner Priel, Dr. Alaa Jamal

01 Motivation

Root-zone pH and EC affect nutrient availability, salinity stress, and fertilizer efficiency. In greenhouses, they are typically monitored through accurate but infrequent manual samples, leaving gaps where changes from irrigation, fertigation, climate, and plant uptake may be missed. Predicting pH and EC from greenhouse data can give growers earlier insight, support faster corrections, and improve crop management.

02 Project Goal

- Develop a soft sensor that predicts future root-zone pH and EC from the latest measurement, forecasted climate, planned irrigation/fertigation and crop-state data.
- Estimate the $t_0 \rightarrow t_1$ change so growers can test a planned irrigation/fertigation strategy, see expected future pH/EC values, and adjust before conditions drift.

03 Data

16,682

10-min rows

109

pH/EC labels

20+

input features


Unified 10-minute master dataset, May 29-Sep 21, 2025:

- External weather & solar radiation — weather station / solar sensor
- Greenhouse micro-climate — internal sensors
- Irrigation & fertigation — management schedule
- Crop-state features — canopy-cover observations
- Manual pH & EC measurements — lab samples

04 Method

Two-stage prediction pipeline:

- Forecast internal greenhouse micro-climate from external weather and radiation.
- Predict $t_0 \rightarrow t_1$ root-zone pH/EC change using climate, crop state and planned events.



05 Algorithms & Model Choice

Hybrid model choice: nonlinear learning + robust modeling of rare extreme intervals.


Gradient Boosting

Main nonlinear model for tabular greenhouse data. Captures interactions between climate, irrigation, fertigation, crop state and baseline pH/EC.

Huber Regressor


Robust companion model for sharp pH/EC shifts. Reduces sensitivity to outliers caused by acid injection, fertigation and salinity events.

Most informative features



06 Validation Strategy


- Walk-forward validation matches deployment: future pH/EC is predicted from past labels, forecasted climate and planned irrigation/fertigation.
- Future pH/EC labels are never used for training until after that sample becomes available.



07 Micro-Climate Model

- Predicts internal greenhouse climate from external weather and radiation.
- Generates continuous climate inputs - temperature, RH, radiation, ET₀ and soil temperature - for the root-zone model.

Stage 1 — Micro-climate model accuracy



08 Root-zone Model Results - Walk-forward Test

0.33 pH MAE

0.93 pH R²


+62% pH MAE reduction

0.127 EC MAE

0.98 EC R²

+35% EC MAE reduction

MAE reduction is relative to a last-value baseline



09 Conclusions

- Prediction value** → The model improved over the last-measurement baseline (+62% pH, +35% EC) and supports earlier pH/EC drift detection.
- Decision-support platform** → built a user platform where growers can test planned irrigation/fertigation actions before applying them.
- Soft-sensor feasibility** → sparse samples can support useful forecasts when combined with dense operational data.

10 Challenges & Limitations

- Sparse labels** → modeled measured t_0 – t_1 changes using accumulated inputs
- Missing climate inputs** → filled greenhouse gaps with a micro-climate model
- Irregular sampling** → added time-gap and window-based aggregation features
- Low EC sensitivity** → used log-transformed EC change and absolute-error metrics
- Limited generalization** → validate across more seasons, crops, and conditions