

① If $t_1(n) \in O(g_1(n))$ and $t_2(n) \in O(g_2(n))$, then find $t_1(n) + t_2(n) \in O(\max\{g_1(n), g_2(n)\})$ prove the assertion.

Sol: We need to show that $t_1(n) + t_2(n) \in O(\max\{g_1(n), g_2(n)\})$. This means there exists a positive constant c and n_0 such that $t_1(n) + t_2(n) \leq c$

$$t_1(n) \leq c_1 g_1(n) \quad \forall n \geq n_0$$

$$t_2(n) \leq c_2 g_2(n) \quad \forall n \geq n_0$$

$$\text{let } n_0 = \max\{n_1, n_2\} \quad \forall n \geq n_0$$

$$\text{consider } t_1(n) + t_2(n) \quad \forall n \geq n_0$$

$$t_1(n) + t_2(n) \leq c_1 g_1(n) + c_2 g_2(n)$$

We need to relate $g_1(n)$ and $g_2(n)$ to $\max\{g_1(n), g_2(n)\}$:

$$g_1(n) \leq \max\{g_1(n), g_2(n)\} \text{ and } g_2(n) \leq \max\{g_1(n), g_2(n)\}$$

thus,

$$c_1 g_1(n) \leq c_1 \max\{g_1(n), g_2(n)\}$$

$$c_2 g_2(n) \leq c_2 \max\{g_1(n), g_2(n)\}$$

$$c_1 g_1(n) + c_2 g_2(n) \leq c_1 \max\{g_1(n), g_2(n)\} + c_2 \max\{g_1(n), g_2(n)\}$$

$$c_1 g_1(n) + c_2 g_2(n) \leq (c_1 + c_2) \max\{g_1(n), g_2(n)\}$$

$$t_1(n) + t_2(n) \leq (c_1 + c_2) \max\{g_1(n), g_2(n)\} \text{ for all } n \geq n_0$$

By the definition of Big O Notation

$$t_1(n) + t_2(n) \in O(\max\{g_1(n), g_2(n)\})$$

$$t_1(n) + t_2(n) \in O(\max\{g_1(n), g_2(n)\})$$

Thus, the assertion is proved

② Find the time complexity of the recurrence relation
Sol: let us consider such that Recurrence for merge sort

$$T(n) = 2T(n/2) + n$$

By using master's theorem

$$T(n) = aT(n/b) + f(n)$$

where $a \geq 1$, $b \geq 1$ and $f(n)$ is positive function

Ex: $T(n) = 2T(n/2) + n$

$$a=2, b=2, f(n)=n$$

By comparing of $f(n)$ with $n \log_b a$

$$\log_b a = \log_2 2 = 1$$

compare $f(n)$ with $n \log_b a$.

$$-f(n) = n$$

$$n \log_b a = n^1 = n$$

* $f(n) = O(n \log_b a)$, then $T(n) = O(n \log_b a \cdot \log n)$

In our case

$$\log_b a = 1$$

$$T(n) = O(n^1 \log n) = O(n \log n)$$

Then time complexity of Recurrence relation is

$$T(n) = 2T(n/2) + n \text{ is } O(n \log n)$$

$$③ \quad T(n) = \begin{cases} 2T(n/2) + 1 & \text{if } n > 1 \\ 1 & \text{otherwise} \end{cases}$$

Sol By applying of Master's Theorem

$$T(n) = aT(n/b) + f(n) \text{ where } a \geq 1, b \geq 1$$

$$T(n) = 2T(n/2) + 1$$

$$\text{Here } a=2, b=2, f(n)=1$$

By comparing of $f(n)$ and $n \log_b a$

$$\text{If } f(n) = O(n^c) \text{ where } c < \log_b a \text{ then } T(n) = O(n \log_b a)$$

$$\text{If } f(n) = O(n \log_b a) \text{ then } T(n) = O(n \log_b a \cdot \log n)$$

$$\text{If } f(n) = \Omega(n^c) \text{ where } c > \log_b a \text{ then } T(n) = O(f(n))$$

lets calculate $\log_b a$

$$\log_b a = \log_2 2 = 1$$

$$f(n) = 1$$

$$n \log_b a = n^1 = n$$

$$f(n) = O(n^c) \text{ with } c < \log_b a \text{ (case 1)}$$

$$\text{In this case } c=0 \text{ and } \log_b a=1$$

$$c < 1 \text{ so } T(n) = O(n \log_b a) = O(n^1) = O(n)$$

Time complexity of Recurrence problem

$$T(n) = 2T(n/2) + 1 \text{ is } O(n)$$

$$④ \quad T(n) = \begin{cases} 2T(n-1) & \text{if } n > 0 \\ 1 & \text{otherwise} \end{cases}$$

Sol Here, where $n=0$

$$T(0) = 1$$

Recurrence relation analysis

for $n > 0$.

$$T(n) = 2T(n-1)$$

$$T(n) = 2T(n-1)$$

$$T(n-1) = 2T(n-2)$$

$$T(n-2) = 2T(n-3)$$

$$T(1) = 2T(0)$$

from this partition

$$T(n) = 2 \cdot 2 \cdot 2 \cdots 2 \cdot T(0) = 2^n \cdot T(0)$$

Since $T(0) = 1$ we have

$$T(n) = 2^n$$

The Recurrence relation is

$$T(n) = 2T(n-1) \text{ for } n > 0 \text{ and } T(0) = 1 \text{ is } T(n) = 2^n$$

⑤ Big O notation show that $f(n) = n^2 + 3n + 5$ is $O(n^2)$

Sol: $f(n) = O(g(n))$ means $c > 0$ and $n_0 \geq 0$

$$f(n) \leq c \cdot g(n) \text{ for } n \geq n_0$$

$$\text{given is } f(n) = n^2 + 3n + 5$$

$$c > 0, n_0 \geq 0 \text{ such that } f(n) \leq cn^2$$

$$f(n) = n^2 + 3n + 5$$

lets choose $c = 2$

$$f(n) \leq 2 \cdot n^2$$

$$f(n) = n^2 + 3n + 5 \leq n^2 + 3n^2 + 5n^2 = 9n^2$$

$$\text{so } c = 9, n_0 = 1, f(n) \leq 9n^2 \text{ for } n \geq 1$$

$$f(n) = n^2 + 3n + 5 \text{ is } O(n^2)$$