


Linear Algebra Basics: Matrices


Foundations of Data Analysis

February 26, 2019



Review: Linear Regression

Model the data as a line:



$$y_i = \alpha + \beta x_i + \epsilon_i$$

α : intercept

β : slope

ϵ_i : error

Reminder: Centering Data

Means:

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad \bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$$

Centered data:

$$\tilde{y}_i = y_i - \bar{y}$$

$$\tilde{x}_i = x_i - \bar{x}$$

Review: Least Squares

Goal: minimize **sum-of-squared error**:


$$\text{SSE}(\alpha, \beta) = \sum_{i=1}^n \epsilon_i^2 = \sum_{i=1}^n (y_i - \alpha - \beta x_i)^2$$

Solution:

$$\hat{\beta} = \frac{\langle \tilde{x}, \tilde{y} \rangle}{\|\tilde{x}\|^2}$$
$$\hat{\alpha} = \bar{y} - \hat{\beta} \bar{x}$$

Review: Linear Regression as Projection

$$\begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \\ \vdots \\ \tilde{y}_n \end{bmatrix} = \beta \begin{bmatrix} \tilde{x}_1 \\ \tilde{x}_2 \\ \vdots \\ \tilde{x}_n \end{bmatrix} + \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{bmatrix}$$



$$\hat{\beta} = \frac{\langle \tilde{x}, \tilde{y} \rangle}{\|\tilde{x}\|^2}$$

Review: Correlation vs. Regression Slope

Correlation:

$$\text{corr}(x, y) = \frac{\langle \tilde{x}, \tilde{y} \rangle}{\|\tilde{x}\| \|\tilde{y}\|} = \cos \theta$$

Regression Slope:

$$\hat{\beta} = \frac{\langle \tilde{x}, \tilde{y} \rangle}{\|\tilde{x}\|^2} = \text{corr}(x, y) \frac{\|\tilde{y}\|}{\|\tilde{x}\|}$$

R^2 Statistic

R^2 statistic gives the **proportion of explained variance**:

$$\begin{aligned} R^2 &= \frac{\text{explained variance in } y}{\text{total variance of } y} \\ &= 1 - \frac{\text{unexplained variance in } y}{\text{total variance of } y} \end{aligned}$$

$$R^2 = 1 - \frac{\text{var}(\epsilon)}{\text{var}(y)} = 1 - \frac{\|\epsilon\|^2}{\|\tilde{y}\|^2}$$

Range: R^2 is always between 0 and 1

Data Tables

	ID	M.F	Hand	Age	Educ	SES	MMSE	CDR	eTIV	nWBV	ASF	Delay	RightHippoVol	LeftHippoVol
0	OAS1_0002_MR1	F	R	55	4	1.0	29	0.0	1147	0.810	1.531	NaN	4230	3807
1	OAS1_0003_MR1	F	R	73	4	3.0	27	0.5	1454	0.708	1.207	NaN	2896	2801
2	OAS1_0010_MR1	M	R	74	5	2.0	30	0.0	1636	0.689	1.073	NaN	2832	2578
3	OAS1_0011_MR1	F	R	52	3	2.0	30	0.0	1321	0.827	1.329	NaN	3978	4080
4	OAS1_0013_MR1	F	R	81	5	2.0	30	0.0	1664	0.679	1.055	NaN	3557	3495
5	OAS1_0015_MR1	M	R	76	2	NaN	28	0.5	1738	0.719	1.010	NaN	3052	2770
6	OAS1_0016_MR1	M	R	82	2	4.0	27	0.5	1477	0.739	1.188	NaN	3421	3119
7	OAS1_0018_MR1	M	R	39	3	4.0	28	0.0	1636	0.813	1.073	NaN	4496	4283
8	OAS1_0019_MR1	F	R	89	5	1.0	30	0.0	1536	0.715	1.142	NaN	3760	3167
9	OAS1_0020_MR1	F	R	48	5	2.0	29	0.0	1326	0.785	1.323	NaN	3557	3394
10	OAS1_0021_MR1	F	R	80	3	3.0	23	0.5	1794	0.765	0.978	NaN	3715	3019
11	OAS1_0022_MR1	F	R	69	2	4.0	23	0.5	1447	0.757	1.213	NaN	3258	3566
12	OAS1_0023_MR1	M	R	82	2	3.0	27	0.5	1420	0.710	1.236	NaN	3217	2160
13	OAS1_0026_MR1	F	R	58	5	1.0	30	0.0	1235	0.820	1.421	NaN	3783	3535
14	OAS1_0028_MR1	F	R	86	2	4.0	27	1.0	1449	0.738	1.211	NaN	3452	3100
15	OAS1_0030_MR1	F	R	65	2	3.0	29	0.0	1392	0.764	1.261	NaN	3969	3406

Row: individual data point

Column: particular dimension or feature

Matrices

A matrix is an $n \times d$ array of real numbers:

$$X = \begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1d} \\ x_{21} & x_{22} & \cdots & x_{2d} \\ \vdots & & & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nd} \end{pmatrix}$$

Notation: $X \in \mathbb{R}^{n \times d}$

A **data matrix** is n data points, each with d features

Row and Column Vectors

Given an $n \times d$ matrix

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1d} \\ a_{21} & a_{22} & \cdots & a_{2d} \\ \vdots & & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nd} \end{pmatrix}$$

Row vectors: $a_{i\bullet} = (a_{i1} \ a_{i2} \ \cdots \ a_{id})$

Column vectors: $a_{\bullet j} = \begin{pmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{nj} \end{pmatrix}$

Matrix Addition

We can add two matrices of the same size: $A, B \in \mathbb{R}^{n \times d}$

$$\begin{aligned} A + B &= \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1d} \\ a_{21} & a_{22} & \cdots & a_{2d} \\ \vdots & & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nd} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & \cdots & b_{1d} \\ b_{21} & b_{22} & \cdots & b_{2d} \\ \vdots & & & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{nd} \end{pmatrix} \\ &= \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1d} + b_{1d} \\ a_{21} + b_{12} & a_{22} + b_{22} & \cdots & a_{2d} + b_{2d} \\ \vdots & & & \vdots \\ a_{n1} + b_{n1} & a_{n2} + b_{n2} & \cdots & a_{nd} + b_{nd} \end{pmatrix} \end{aligned}$$

Matrix-Scalar Multiplication

We can multiply a matrix $X \in \mathbb{R}^{n \times d}$ by a scalar $s \in \mathbb{R}$:

$$sX = s \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1d} \\ a_{21} & a_{22} & \cdots & a_{2d} \\ \vdots & & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nd} \end{pmatrix} = \begin{pmatrix} sa_{11} & sa_{12} & \cdots & sa_{1d} \\ sa_{21} & sa_{22} & \cdots & sa_{2d} \\ \vdots & & & \vdots \\ sa_{n1} & sa_{n2} & \cdots & sa_{nd} \end{pmatrix}$$

Matrix-Vector Multiplication

We can multiply an $n \times d$ matrix A with a d vector v :

$$Av = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1d} \\ a_{21} & a_{22} & \cdots & a_{2d} \\ \vdots & & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nd} \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_d \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^n a_{1j}v_j \\ \sum_{j=1}^n a_{2j}v_j \\ \vdots \\ \sum_{j=1}^n a_{nj}v_j \end{pmatrix}$$

The result is a d vector.

Each entry is a dot product between a row of A and v :

$$Av = \begin{pmatrix} \langle a_{1\bullet}, v \rangle \\ \langle a_{2\bullet}, v \rangle \\ \vdots \\ \langle a_{n\bullet}, v \rangle \end{pmatrix}$$


Matrices as Transformations

Consider a 2D matrix and coordinate vectors in \mathbb{R}^2 :

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \quad v_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Then Av_1 and Av_2 result in the columns of A :

$$Av_1 = \begin{pmatrix} a_{11} \\ a_{21} \end{pmatrix}, \quad Av_2 = \begin{pmatrix} a_{12} \\ a_{22} \end{pmatrix}$$



Linearity of Matrix Multiplication

Given a matrix $A \in \mathbb{R}^{n \times d}$, vectors $v, w \in \mathbb{R}^d$, and scalars $s, t \in \mathbb{R}$:

$$A(sv + tw) = sAv + tAw$$

Matrix-Matrix Multiplication

Given matrices $A \in \mathbb{R}^{m \times d}$ and $B \in \mathbb{R}^{d \times n}$,
their **product** is:

$$(AB)_{ij} = \sum_{k=1}^d a_{ik} b_{kj}$$

Or, using dot products:

$$(AB)_{ij} = \langle a_{i\bullet}, b_{\bullet j} \rangle$$

Note: A column dimension = B row dimension

Note: resulting product AB is $m \times n$

Multilinear Regression

What if x_i has two features (x_{i1}, x_{i2}) ?

$$y_i = x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

Written as matrix-vector multiplication:

$$\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \\ \vdots & \vdots \\ x_{n1} & x_{n2} \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} + \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{pmatrix}$$

Or, in matrix notation:

$$y = X\beta + \epsilon$$

Multilinear Regression: d features

What if x_i has d features $(x_{i1}, x_{i2}, \dots, x_{id})$?

$$y_i = x_{i1}\beta_1 + x_{i2}\beta_2 + \dots + x_{id}\beta_d + \epsilon_i$$

Written as matrix-vector multiplication:

$$\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1d} \\ x_{21} & x_{22} & \cdots & x_{2d} \\ \vdots & \vdots & & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nd} \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_d \end{pmatrix} + \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{pmatrix}$$

Or, in matrix notation:

$$y = X\beta + \epsilon$$

Multilinear Regression: Adding an Intercept

Use β_1 for the intercept:

$$y_i = \beta_1 + x_{i2}\beta_2 + \cdots + x_{id}\beta_d + \epsilon_i$$

First column of X is ones:

$$\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & x_{12} & \cdots & x_{1d} \\ 1 & x_{22} & \cdots & x_{2d} \\ \vdots & \vdots & & \vdots \\ 1 & x_{n2} & \cdots & x_{nd} \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_d \end{pmatrix} + \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{pmatrix}$$

Or, in matrix notation:

$$y = X\beta + \epsilon$$

Matrix Inverse

Given a square matrix $A \in \mathbb{R}^{d \times d}$, it's **inverse** is a matrix A^{-1} such that:

$$A^{-1}A = AA^{-1} = I$$

The inverse exists if and only if A has linearly independent columns.

Transpose

The transpose of a matrix $A \in \mathbb{R}^{n \times d}$ is a matrix $A^T \in \mathbb{R}^{d \times n}$ that “flips” row and column indices:

$$(A^T)_{ij} = A_{ji}$$

Note: we can do inner products with transpose:

$$\langle v, w \rangle = v^T w$$

Least Squares Problem

Regression equation:

$$y = X\beta + \epsilon$$

Minimize the sum-of-squared error:

$$\begin{aligned}\text{SSE}(\beta) &= \|\epsilon\|^2 \\ &= \|y - X\beta\|^2 \\ &= (y - X\beta)^T (y - X\beta)\end{aligned}$$

LS Solution

Derivative:

$$\frac{\partial}{\partial \beta} \text{SSE}(\beta) = -X^T(y - X\beta) = -X^T y + X^T X \beta$$

$$X^T X \beta = X^T y$$

set derivative to zero

$$(X^T X)^{-1} (X^T X) \beta = (X^T X)^{-1} X^T y$$

multiply by $(X^T X)^{-1}$

$$\hat{\beta} = (X^T X)^{-1} X^T y$$

solve for β