

School of Geography and Geology  
McMaster University

---

Applied Spatial Statistics

**Point Pattern Analysis I & II**

# This session:

---

- Point Pattern Analysis I
  - Definitions
  - First and Second Order Properties
  - Visualizing Point Patterns
  - Exploring Point Patterns
    - Quadrat counts
    - Quadrat Analysis

# Definitions


---

- Region
- Point
- Event
- Attributes
- Patterns
  - Random
  - Clustered
  - Dispersed/Regular

# Definitions: Region

---

- **Region ( $R$ )** – Specific area over the surface of the earth that is of interest



# Definitions: Region

---

- **Examples of regions - Continent**



# Definitions: Region

---

- **Examples of regions - Province**



# Definitions: Region

---

- **Examples of regions**
  - **City**



# Definitions: Region

---

- **Examples of regions**
  - **Land plot**





# Definitions: Region

---


- **Examples of regions**
  - **Computer chip**



# Definitions: Region

---



- Examples of regions
  - Section of tissue




# Definitions: Region

---

- *Regular* region




- *Irregular* region



# Definitions: Point

---


- Point – An arbitrary location in region  $R$



# Definitions: Event


---

- Event – An “interesting” occurrence in region  $R$




$\mathbf{s}=(s_1,s_2)$   
"x" and "y"  
coordinates

# Definitions: Event – Examples



# Definitions: Event – Examples



Note:  
What about the  
underlying population?

# Definitions: Event – Examples

---

- Event – Further examples




# Definitions: Attributes

---

- $s_1, s_2$ : the “x” and “y” coordinates of the event
- Other values attached to event
  - Type of occurrence
    - Type of building
    - Tree species
    - ...
  - Time of occurrence
  - Other characteristics
    - Depth of borehole
    - Age
    - ...


# Definitions: Patterns

---




# Definitions: Patterns

---



# Definitions: Patterns


---



# Definitions: Population

---


- *Mapped* point pattern – A complete map of events in  $R$



# Definitions: Sample

---


- ***Sampled* point pattern – Some events in  $R$  have not been recorded**



# Definitions: Sample

---


## ○ Limitations - Example



# Definitions: Edge Effects

---

## ○ Edge Effects






# Definitions: Edge Effects

---


- **Possible Solution – Creating a torus**



# Definitions: Edge Effects


---

- **Possible Solution – Guard Area**



# First & Second Order Properties

## ○ Number of Events per Unit Area



# First & Second Order Properties

---

- **Number of Events per Unit Area**
  - $Y(\mathcal{A}_i)$ : Number of events per unit area  $i$
  - $E[Y(\mathcal{A})]$ : Mean number of events per unit area
  - $\text{COV}[Y(\mathcal{A}_1), Y(\mathcal{A}_2)]$ : Covariance between  $\mathcal{A}_1$  and  $\mathcal{A}_2$

# First & Second Order Properties

---

- **Number of Events per Unit Area**

**Depends on the size of  $\mathcal{A}$**

# First order properties

---

- **Intensity – Mean number of events per unit area at point  $s$**

$$\lambda(\mathbf{s}) = \lim_{ds \rightarrow 0} \left\{ \frac{E[Y(\mathbf{ds})]}{ds} \right\}$$

- **Stationary process:  $\lambda(\mathbf{s})$  is a constant  $\rightarrow \lambda$**

# Second order properties

---

- **Second Order Intensity**


$$\gamma(\mathbf{s}_i, \mathbf{s}_j) = \lim_{ds_i, ds_j \rightarrow 0} \left\{ \frac{E[Y(\mathbf{ds}_i)Y(\mathbf{ds}_j)]}{ds_i ds_j} \right\}$$

- **Stationary, isotropic process:**  
 **$\gamma(\mathbf{s})$  depends on distance only**

# Visualizing Point Patterns

---


- Dot Map





# Example: Shops in Hamilton, ON

---



# Visualizing Point Patterns

---

- Dot Map with Attributes

# Example: Shops in Hamilton, ON

---

# Example: Cities in Eastern Asia

---

- Dot Map with Attributes


# Example: Cities in Eastern Asia

---

- Dot Map with Attributes

# Visualizing Point Patterns

- Dot Map with Attributes
  - Represent underlying population




# Exploring Point Patterns

---

- Methods
  - Quadrat Counts/Analysis
  - Kernel Estimation

# Quadrat Analysis

**Figure 5.5** Illinois Tornado Pattern with Quadrats Superimposed





# Quadrat Analysis




Figure 2.1 Location of Settlements of 300 or Greater Population in 1971 in Southern Saskatchewan

# Quadrat Counts – Moving Window





Figure 2.1 Location of Settlements of 300 or Greater Population in 1971 in Southern Saskatchewan


# Quadrat Counts – Moving Window

- Smooth variation of intensity  $\lambda(\mathbf{s})$   
(Good)



# Quadrat Counts – Moving Window


- Relative location of points within window?



# Quadrat Counts – Moving Window

---


- Size of window?




# Quadrat Counts – Moving Window

---

- Large window






- Small window



# Quadrat Analysis

---

- Based on the frequency of events occurring in various parts of the region



# Quadrat Analysis

---

- Mean cell frequency

- $MEAN = \# \text{ of events } (n) / \# \text{ of cells } (m)$

- Variance of the cell frequencies

- $$VAR = \frac{\sum f_i X_i^2 - (\sum f_i X_i)^2 / m}{m - 1}$$

- Variance–mean ratio

- $VMR = VAR / MEAN$



# Case 1


---

# of Events $X$	$X^2$	# of Cells with $X$ Events ( $f$ )	$fX$	$fX^2$
$X=0$	0	0	0	0
$X=1$	1	0	0	0
$X=2$	4	0	0	0
$X=3$	9	9	27	81
$X=4$	16	0	0	0
$X=5$	25	0	0	0
		Sum=	27	81

$$MEAN = 27/9 = 3 \quad VAR = \frac{81 - 27^2/9}{9-1} = 0 \quad VMR = \frac{0}{3} = 0$$


# Case 2

---



# Case 3

---



# Quadrat Analysis

---

- Variance-mean ratio
  - $VMR < 1$ : The pattern is more dispersed than random
  - $VMR > 1$ : The pattern is more clustered than random
  - $VMR = 1$ : The pattern is random

# Next...

---

- Kernel estimation
- Second order properties
- Nearest neighbor analysis
- K Functions