

Stars and Planets Problem Set I

Qingru Hu

March 9, 2023

Exercise I.1 Sun's efficiency

We can calculate the energy density of the Sun from the solar constant, that is the energy recieved by the Earth in unit area in unit time. The solar constant is $P = 1367 \text{ Wm}^{-2}$, and we know that the distance from the Sun to the Earth is $r = 1 \text{ au}$, so we can calculate the luminosity of the Sun:

$$L = 4\pi r^2 P = 3.845 \times 10^{26} \text{ W} \quad (1)$$

Therefore, the energy output of the Sun per unit mass $\rho_{\text{E, Sun}}$ is:

$$\rho_{\text{E, Sun}} = \frac{L}{M_{\odot}} = 1.93 \times 10^{-4} \text{ W kg}^{-1} \quad (2)$$

The radiation energy density of a human being can be estimated from a human being's temperature, which is normally $T = 300\text{K}$. And we can assume that the skin area of a human being is about $A = 1\text{m}^2$, and the mass of a human being is $m = 50 \text{ kg}$. Therefore, we can calculate the radiation energy of a human being to be:

$$\rho_{\text{E, Human}} = \frac{\sigma T^4 A}{m} = 9.1854 \text{ W kg}^{-1} \quad (3)$$

Exercise I.2 Color-Magnitude diagram

(a)

I choose the "hipparcos-bright.txt" dataset to draw the color-magnitude diagram Fig.1. The star that has the greatest parallax is highlighted with a blue dot and the brightest one with a red dot.

(b)

That's because for a specific star the apparent and the absolute differ only by a constant.

$$m - M = 5 \log_{10} \left(\frac{d}{10 \text{ pc}} \right) \quad (4)$$

The $B - V$ color index is defined as the subtraction of magnitudes at the B and V bands, so the constant cancels out.




Figure 1: The color-magnitude diagram of "hipparcos-bright.txt"

(c)

According to the Wien's displacement law:

$$\lambda_{\max} = \frac{0.2898 \text{ cm K}}{T} \quad (5)$$

the lower the effective temperature T_{eff} , the more radiation in the "visual" band (V) and the less radiation in the "blue" band (B). Therefore, the star has larger magnitude in the "blue" band (B) and smaller magnitude in the "visual" band (V), and their subtraction $B - V$ color is higher. That's why stars of higher $B - V$ have lower T_{eff} .

(d)

The "hipparcos-bright" sample doesn't contain white dwarfs. The main sequence stars and the RGB stars are labeled as blue and orange respectively in the Fig.3. I only does a rough classification of these two groups of stars.

(e)

Exercise I.3 Transits

(a)

We assume a circular orbit and uniform random probability distribution of orbital revolution axes on the unit sphere.

The probability of the transit not being observed by us is the fraction of the solid angle expanded by θ_0 in Fig.4 and 2π (the solid angle of the half hemisphere):

$$\cos \theta_0 = \frac{R_{\star} + R_{\text{p}}}{a} \quad (6)$$

$$P_{\text{not-trans-orbit}} = \frac{\int_0^{\theta_0} \sin \theta d\theta \int_0^{2\pi} d\phi}{2\pi} = 1 - \cos \theta_0 \quad (7)$$

Therefore, the probability that a planet will transit at some point in its orbit is:

$$P_{\text{trans-orbit}} = 1 - P_{\text{not-trans-orbit}} = \cos \theta_0 = \frac{R_{\star} + R_{\text{p}}}{a} \quad (8)$$




Figure 2: The color-magnitude diagram with classification




Figure 3: The Hertzsprung-Russell diagram of the hipparcos-bright samples




Figure 4: The orbit demenstration of a critical situation

(b)




Figure 5: The transit configuration

The side view of the planet transit is given in Fig.5, and only the planets whose orbital direction are in the shaded region can transit. From the geometry and the above problem(a) we can get that:

$$\sin \theta_m = \frac{R_p + R_\star}{a}$$

$$z = a \sin \theta$$

As shown in Fig. , the length of the transit path Δl can be approximated as:

$$\Delta l = 2\sqrt{(R_p + R_\star)^2 - z^2} = 2(R_p + R_\star)\sqrt{1 - \left(\frac{a \sin \theta}{R_p + R_\star}\right)^2} \quad (9)$$

The ratio of the transit duration and the orbital period is:

$$\frac{\Delta t}{T} = \frac{\Delta l}{2\pi a} = \frac{1}{\pi} \frac{R_p + R_\star}{a} \sqrt{1 - \left(\frac{a \sin \theta}{R_p + R_\star}\right)^2} \quad (10)$$

The fraction of possible orbits in the range $(\theta, d\theta)$ is $\frac{2\pi \cos \theta d\theta}{4\pi} = \frac{1}{2} \cos \theta d\theta$, so the probability of current transits can be written as:

$$\begin{aligned} P &= \int_{-\theta_m}^{\theta_m} P_{\text{trans-orbit}} \cdot \frac{\Delta t}{T} \cdot \frac{1}{2} \cos \theta d\theta \\ &= \frac{1}{2\pi} \left(\frac{R_p + R_\star}{a} \right)^2 \int_{-\theta_m}^{\theta_m} \sqrt{1 - \left(\frac{\sin \theta}{\sin \theta_m} \right)^2} d(\sin \theta) \\ &= \frac{1}{4} \left(\frac{R_p + R_\star}{a} \right)^2 \end{aligned}$$

(c)

Assuming that all of the $n = 100,000$ stars harbor such an Earth-like planet, so $R_\star = R_\odot = 7 \times 10^{10}$ cm, $R_p = R_\oplus = 6.4 \times 10^8$ cm and $a = 1 \text{ AU} = 1.5 \times 10^{13}$ cm. Pluge into the equation (5) and we can get the number N of the planets that will eventually show a transit is:

$$N = P_{\text{trans-orbit}} n = \frac{R_\odot + R_\oplus}{a} n \approx 471 \quad (11)$$

(d)

The maximum percentage change Δ in brightness of an Earth-like planet is:

$$\Delta = \frac{\Delta F}{F} = \frac{\pi R_p^2}{\pi R_\star^2} = \frac{R_\oplus^2}{R_\odot^2} = 0.01\% \quad (12)$$

The relation between the magnitude m and the flux F is :

$$m = -2.5 \log F + \text{const} \quad (13)$$

Take derivatives on both sides and we get:

$$\Delta m = -\frac{2.5}{\ln(10)} \frac{\Delta F}{F} \approx -1.086 \Delta = -1.086 \times 10^{-4} \quad (14)$$

Therefore, the drop in magnitude is $\Delta m = -1.086 \times 10^{-4}$.

(e)

I think Bernie is right. The white dwarf is as bright as the host star, so the white dwarf will not cause a dip in the their joint brightness when it orbits before the host star. However, when the white dwarf orbits behind the host star, light emitted by the white dwarf will be blocked by the host star, resulting in a dip in their joint brightness, which is very similar to a transit signal. This effect is shown in Fig.6. In fact, eclipsing binary systems like this make up most of the falses positives in trasit planet detections.

(f)

The radial-velocity amplitude v_\star can be calculated from the following equations:

$$m_\star v_\star = m_p v_p \quad (15)$$

$$G(m_\star + m_p) = \frac{4\pi^2 a^3}{P^2} \quad (16)$$




Figure 6: The orbit and light curve of an eclipsing binary, credit: Wikimedia, NASA

We notice that $vP = 2\pi a$, where $v = v_\star + v_p$, so we can cancel P and v_p :

$$G(m_\star + m_p) = a(v_\star + \frac{m_\star v_\star}{m_p})^2$$

$$v_\star = \frac{m_p}{m_p + m_\star} \sqrt{\frac{G(m_\star + m_p)}{a}}$$

For an Earth-like planet, we can get that $v_\star = 8.95 \text{ cm s}^{-1}$; for a white dwarf companion, we can get that $v_\star = 2106.12 \text{ km s}^{-1}$.

(g)

Due to the huge radial velocity amplitude of a white dwarf, we can rule out that the transiting body is a white dwarf by taking an RV measurement.

(h)

The precision of the current RV measurements is too low to detect the extreme small radial velocity amplitude of an Earth-like planet. Therefore it is not useful to ensure that the transiting body is an Earth-mass planet using RV measurement.

Exercise I.4 Microlensing

(a)

Set the origin of x axis at the periapsis of the trajectory, and denote the periapsis-passing time as $t = 0$. At any time t , the y component of the equation of motion is given by:

$$\frac{dv_y}{dt} = \frac{GM}{b^2 + x^2} = \frac{GM}{b^2 + (vt)^2}. \quad (17)$$

The above equation can be integrated directly and gives:

$$v_y = \int_0^{v_y} dv'_y = \int_{-\infty}^{\infty} \frac{GM}{b^2 + (vt)^2} dt = \frac{GM}{bv} \int_{-\infty}^{\infty} \frac{GM}{1 + u^2} du = \frac{\pi GM}{bv}. \quad (18)$$

(b)

The scattering angle can easily be derived by:

$$\psi \approx \frac{v_y}{v_x} = \frac{\pi GM}{bv^2}. \quad (19)$$

(c)

Assume that we have a source object at distance d_s from the observer, and a lens object at distance $d_l < d_s$. We further assume that these two objects seen from the observer are aligned. Now we have the following geometric relation

$$\psi(d_s - d_l) = \theta_E d_s \quad (20)$$

and

$$b = \theta_E d_l. \quad (21)$$

Therefore we have:

$$\frac{\pi GM}{\theta_E d_l c^2} (d_s - d_l) = \theta_E d_s. \quad (22)$$

And it turns out to be:

$$\theta_E = \sqrt{\frac{\pi GM}{c^2} \frac{d_s - d_l}{d_s d_l}}. \quad (23)$$

(d)

The Einstein ring can be calculated as:

$$\theta_E = \sqrt{\frac{4GM_L}{c^2} \frac{d_s - d_l}{d_l d_s}} \quad (24)$$

Pluge in the values and we can get:

$$\theta_E = 3.46 \times 10^{-9} \text{ radian} = 0.713 \text{ mas}$$

The Einstein ring radius in the lens plane r_{lE} is:

$$r_{lE} = \theta_E d_l = 2.854 \text{ au}$$

The Einstein ring radius in the source plane r_{sE} is:

$$r_{sE} = \theta_E d_s = 5.708 \text{ au}$$

(e)

The magnification is the same when $\beta = +/ - \theta_E$, so we only need to calculate one situation. When $\beta = \theta_E$, the image θ satisfies

$$\theta_E = \theta - \frac{\theta_E^2}{\theta} \quad (25)$$

The two solutions to the above equation is:

$$\begin{aligned} \theta_1 &= 0.118\theta_E \\ \theta_2 &= -2.118\theta_E \end{aligned}$$

The magnification factor m can be calculated:

$$m = \sum_{i=1}^2 \frac{\theta_i^4}{\theta_i^4 - \theta_E^4} = 1.052 \quad (26)$$

The source will be brighter in magnitude by $2.5 \times \log_{10} m = 0.055$

(f)

From (e), we can see that when the source cross the Einstein ring, it is slightly magnified. Therefore, from the figure we can get that the duration of the Einstein ring crossing time is:

$$t_{\text{E, cross, star}} = 8725 - 8650 = 75 \text{ days} \quad (27)$$

(g)

Assuming that the proper motion speed between the source and the lens is the same during the microlensing event and the caustic of the planet and the star does not affect each other very much (still perserve the Einstein ring shape), the mass ratio of the planet and the host star is proportional to the square root of the Einstein ring crossing time.

$$\frac{m_{\text{p}}}{m_{\star}} = \sqrt{\frac{t_{\text{E, cross, planet}}}{t_{\text{E, cross, star}}}} \quad (28)$$

If $m_{\star} = 0.5M_{\odot}$, then the mass of the planet is:

$$m_{\text{p}} = m_{\star} \sqrt{\frac{t_{\text{E, cross, planet}}}{t_{\text{E, cross, star}}}} = 9611 \text{ } M_{\oplus} \quad (29)$$