

Causal Recreational Cannabis

Patrick Massey and Elliott Metzler

5/13/2022

Abstract

Can legalizing the recreational use of cannabis reduce rates of drug poisoning deaths? This paper seeks to study the relationship between changes in state-level policy toward recreational cannabis use and drug poisoning death rates. Specifically, we evaluate the impact of such policy change on the first state to legalize, Colorado. We implement a synthetic control design in which we use demographic data for each state in the United States, including Washington D.C., and estimate the causal impact of Colorado's legalization policy shift in 2012. We find [[insert findings and explanation of implications]].

1 Introduction and Background

[[Need to do]]

Table 1: Marijuana Legalization by State

State	Year
Colorado	2012
Washington	2012
Alaska	2014
Oregon	2014
California	2016
Maine	2016
Massachusetts	2016
Nevada	2016
Michigan	2018
Vermont	2018

Figure 1 shows the average death rate across the United States and the average death rate for Colorado between 2000 and 2018.




Figure 1: Drug Poisoning Death Rates Trend (2000-2018)

2 Data

The data employed for this paper originates from two sources. First, we source data on drug poisoning deaths, population, and an estimate of the crude drug poisoning death rate from the United States Center for Disease Control (“CDC”). Second, we source state-level demographic data from IPUMS USA, which is a database harmonizing, organizing, and structuring data from various population surveys in the United States over time.

The first data set contains summary statistics by state and by year from 1999 to 2018. For each state in each year, the data reports the number of deaths due to drug poisoning, the population, and the ratio of the two previous variables as the crude death rate. Additionally, this data reports an estimate of the standard error for the crude death rate, and a lower and upper confidence interval around the point estimate. In addition to these base estimates, the data originally includes age-adjusted rates with standard errors and confidence intervals and the United State-wide crude death rate and United State-wide age-adjusted death rate.

From this data set, we focus on the crude death rate for the state and extract this variable along with the state and year variables for data merging purposes. We focus our attention on the crude death rate statistic for a couple of important reasons. First, using the death rate as opposed to the total number of deaths allows us to immediately control for differences in state population when estimating our model. Second, we exclude the age-adjusted rate because our model also accounts for age demographics explicitly as covariates, so it would be inappropriate to use this as our outcome variable of interest.

The second data set contains nearly 28 million entries where each entry represents a person included in [[a survey]] in a particular year. Importantly, this data includes the year and state in which that entry resided along with extensive demographic data. Thus, we are able to use the groups of these entries associated with each year and each state and calculate demographic summary information representative of each year in each state. We include in our demographic summaries the proportion of the population that is listed as male as compared to female [[should footnote simplicity of coding here (I don’t remember seeing info about non-binary or choose not to identify)]]], a summary of a few significant

age buckets, proportions of the population based on race, on marital status, on education, and on employment status. Additionally, we summarize the mean number of hours worked, the median income, the mean number of children, and the mean number of children under 5 years old. We present the mean and standard deviation of each of these demographic variables in Table 2.

To construct our final data set for use in analysis, we combine the two data sources. Since we have summarized the latter data set at the state and year level and the first data set is already at this level, we merge on the state and year. Thus, we have data for each state, inclusive of Washington D.C., for each year between 2000 and 2018 with data on deaths associated with drug poisoning, population, crude death rate, and all of our descriptive demographics data.

Table 2: Variable Summary

Variable	Mean	Std. Dev
Drug Poisonings		
Deaths	777.179	887.743
Population	5995451.840	6730237.227
Crude Death Rate	13.381	7.082
Gender Population Proportions		
Male	0.489	0.012
Female	0.511	0.012
Age Population Proportions		
Under 30	0.176	0.022
Under 50	0.450	0.042
Over 50	0.374	0.045
Race Population Proportions		
American Indian	0.018	0.039
Asian	0.039	0.070
Black	0.088	0.094
Other Race	0.045	0.039
White	0.810	0.131
Hispanic Population Proportions		
Hispanic	0.076	0.081
Not Hispanic	0.924	0.081
Marital Status Population Proportions		
Married	0.609	0.062
Not Married	0.391	0.062
Education Population Proportions		
Less Than High School	0.084	0.029
High School	0.372	0.050
Some College	0.247	0.036
College	0.191	0.034
More Than College	0.106	0.041
Employment Population Proportions and Summary Statistics		
Employed	0.724	0.045
Not Employed	0.276	0.045
Mean Hrs Worked	32.222	2.269
Median Income	22380.650	5043.007
Children Summary Statistics		
Mean Children	0.823	0.107
Mean Children Under 5 years old	0.168	0.029

Please see the appendix for detailed descriptions of our procedure to re-code and summarize variables.

3 Methodology

We utilize the synthetic control method first developed by Abadie and Gardeazabal (2003) and then expanded upon in Abadie et. al (2010)(2015). Similar to Abadie et. al (2010) we are analyzing the impact of a government policy shock. In this analysis we are estimating the impact of Colorado’s legalization of recreation cannabis in 2012. For our synthetic control, Colorado is the treatment state of interest, and we have a donor pool of 41[Need to double check that’s correct] states. The synthetic control method [MORE INFO HERE]

Suppose that we have $1, 2 \dots S + 1$ states and let $s = 1$ be the treated state, additionally we have $t = 1, 2 \dots T$ time periods where T_0 represents the number of pretreatment periods, and $T_0 + 1 \dots T$ are the posttreatment periods. With that let Y_{st} be the crude death rate for state s at time t . In a posttreatment time line we are estimating

$$\hat{\alpha} = Y_{1t} - \sum_{s=2}^{S+1} w^* Y_{st}$$

w_s^* is a vector of optimally chosen weights for states from our donor pool to represent our synthetic Colorado. Again deriving from Abadie (2010) w_s^* is the vector of weights that minimizes the following equation

$$(X_1 - X_0 w)' V (X_1 - X_0 w)$$

X_1 is a $(K \times 1)$ vector of state population variables and X_0 is a $(K \times S)$ matrix of state population variables for S donor states. Note that $w_s \geq 0$ and $\sum w_s = 1$.

We utilize the weights generated to create a synthetic Colorado, where we examine the counterfactual of a Colorado that never legalized recreational cannabis. Additionally we follow Abadie (2010) and use their placebo technique as a robustness check. The placebo technique consists of assigning each state in our donor pool as the treated state and then creating an optimal w_s^* , from the other donor states as well as the treated state, in the pretreatment period. Utilizing the weights we calculate the posttreatment root mean squared prediction

error (RMSPE) for each state. We then take a ratio between the post/pretreatment RMSPE for each state. We expect this value to be high for our treated state, which would suggest that the posttreatment RMSPE is large due to the differences between the synthetic and the observed Colorado.

Table 3: Synthetic Weights

Unit	Weight
Alabama	0.0000000
Arizona	0.4537880
Arkansas	0.0000000
Connecticut	0.0000016
Delaware	0.0000000
District of Columbia	0.0946779
Florida	0.0000000
Georgia	0.0000000
Hawaii	0.0000001
Idaho	0.0000000
Illinois	0.0000000
Indiana	0.0000000
Iowa	0.0000000
Kansas	0.0000001
Kentucky	0.0000000
Louisiana	0.0000000
Maryland	0.0000000
Minnesota	0.0000603
Mississippi	0.0000000
Missouri	0.0000000
Montana	0.0000000
Nebraska	0.0000001
New Hampshire	0.3754123
New Jersey	0.0000001
New Mexico	0.0000000
New York	0.0000000
North Carolina	0.0000000
North Dakota	0.0000005
Ohio	0.0000000
Oklahoma	0.0000000
Pennsylvania	0.0000000
Rhode Island	0.0000000
South Carolina	0.0000000
South Dakota	0.0000000
Tennessee	0.0000000
Texas	0.0000005
Utah	0.0760580
Virginia	0.0000001
West Virginia	0.0000000
Wisconsin	0.0000000
Wyoming	0.0000000

Table 4: Balance Table

Variable	Colorado	Synthetic Colorado	Donor Sample
american_indian_prop	9.350500e-03	2.247720e-02	1.359220e-02
asian_prop	2.643560e-02	2.504760e-02	3.465280e-02
black_prop	2.950610e-02	5.946740e-02	9.825980e-02
college_prop	2.448619e-01	2.038708e-01	1.840564e-01
employed_prop	7.532957e-01	7.324926e-01	7.260995e-01
female_prop	5.075856e-01	5.140958e-01	5.147653e-01
hispanic_prop	1.387230e-01	1.202302e-01	6.652670e-02
hs_prop	3.035144e-01	3.409944e-01	3.835767e-01
less_than_hs_prop	6.840350e-02	8.375930e-02	9.042740e-02
male_prop	4.924144e-01	4.859042e-01	4.852347e-01
married_prop	6.315082e-01	6.044073e-01	6.271986e-01
mean_children	8.130376e-01	8.475448e-01	8.435551e-01
mean_children_u5	1.828829e-01	1.788849e-01	1.739943e-01
mean_hrs_worked	3.388178e+01	3.267719e+01	3.270569e+01
median_income	2.452308e+04	2.347387e+04	2.132795e+04
more_than_college_prop	1.297666e-01	1.217832e-01	9.962340e-02
not_employed_prop	2.467043e-01	2.675074e-01	2.739005e-01
not_hispanic_prop	8.612770e-01	8.797698e-01	9.334733e-01
not_married_prop	3.684918e-01	3.955927e-01	3.728014e-01
other_race_prop	6.431370e-02	5.621730e-02	4.057560e-02
over_50_prop	3.368499e-01	3.481782e-01	3.596021e-01
some_college_prop	2.534536e-01	2.495922e-01	2.423161e-01
under_30_prop	1.803345e-01	1.797119e-01	1.745255e-01
under_50_prop	4.828157e-01	4.721099e-01	4.658724e-01
white_prop	8.703941e-01	8.367905e-01	8.129196e-01




Figure 2: Trends in Crude Death Rate: Colorado vs. Synthetic Colorado




Figure 3: Differences in Crude Death Rate: Colorado vs. Synthetic Colorado

4 Discussion

5 Conclusion

6 Appendix

6.1 Variable Re-coding Procedure Detail

- Gender (*sex*)

Original coding:

- 1 = Male
- 2 = Female

Summarized by calculating proportion of total that is male and proportion of total that is female.

- Age (*age*)

Original coding:

- Actual age of entry

Summarized by categorizing into the following age groups and calculating the proportion of the total that fall into that age group.

- Under 30 years old
- Between 30 years old and under 50 years old
- Over 50 years old

- Need to do - finish transferring notes from the markdown file into here.